

# Biological Monitoring Heavy Metals in Fingernails and Scalp Hair of Autoworkers in Saudi Arabia

Eida Mohammed Alshammari

Received: 10 January 2022 / Received in revised form: 24 March 2022, Accepted: 25 March 2022, Published online: 28 March 2022

## Abstract

Biological materials have gained an irreplaceable position as a bio-indicator for environmental pollution with heavy metals. Human hair and nails have come to play an important role in this subject according to the recommendations by the World Health Organization (WHO). The levels of selected toxic and heavy metals (Pb, Zn, Cd, and Ni) in the fingernails and scalp hair of autoworkers in Saudi Arabia were estimated to study exposure to chemical pollutants metals. The samples were collected from different ages of male autoworkers (car mechanics, auto welding, repair car tires washing, and oil change services center) and office workers (control) volunteers in Hail city. The analysis was performed using inductively coupled plasma mass spectrometry (ICP-MS) after microwave acid digestion at their recommended conditions. The data obtained from office workers (control) samples were used for this purpose. A systematic analysis investigation was done for all samples taking into consideration the age and experience per year. As a result of this research, it has been revealed that there was a statistically significant relationship between the type of work (autoworkers) or workplace and the heavy metal concentration detected in fingernails and scalp hair samples, which represent a very important guide for the labor and health organizations. This result proved that fingernails and scalp hair could be used as a good biological indicator for the assessment of toxic heavy metal pollution.

**Keywords:** Fingernails, Scalp hair, Autoworkers, Saudi Arabia, Heavy metal pollution

## Introduction

Environmental pollution has an important role in increasing the levels of toxic and heavy metals in the human body, which lead to various chronic diseases (Rajfur *et al.*, 2018; Silva *et al.*, 2018; Ocelić Bulatović *et al.*, 2019; Cabar *et al.*, 2020; Noreen *et al.*, 2020; Almeida-da-Silva *et al.*, 2021; Ikese, *et al.*, 2021; Wu *et al.*, 2021; Lotah, 2022). The working environment has come to play an important role in this subject, due to the direct relationship between man and the degree of hazards chemical agents present in their workplace environment. In addition, each workplace environment

is not unique but also in the degree of hazards it poses to the workers. Furthermore, the pollution of the workplace environment with toxic heavy metals is a worldwide problem (Lemos & De Carvalho, 2010; Massadeh *et al.*, 2010; Amartei *et al.*, 2011; Peter *et al.*, 2012; Vincenti *et al.*, 2013; Momčilović *et al.*, 2016; Alrobaian & Arida, 2019; Al-Muzafar *et al.*, 2021; Ikese, *et al.*, 2021; Lotah *et al.*, 2022). In this context, toxic and heavy chemicals come out in the workplace environment from various uses in the form of spray or dust, which can get into the body of workers mainly either through inhalation and/or dermal contact. Therefore, accurate assessment of toxic and heavy elements in the human tissues is extremely important. Therefore, analyzed with some biological materials such as blood, teeth, nails, and hair; it is possible to monitor changes in the body. Various analytical techniques have been considered for the measurement of heavy and toxic metals in different biological samples (Geier *et al.*, 2016; Mehra & Thakur, 2016; Ocelić Bulatović *et al.*, 2019; Noreen *et al.*, 2020; Almeida-da-Silva *et al.*, 2021). Fingernails and scalp hair can be considered metabolic end products, as well as they, are long-term with many trace elements into their structure during the growth process, the trace element contents of which reflect mineral metabolism in the body (Rajfur *et al.*, 2018; Silva *et al.*, 2018; Cabar *et al.*, 2020; Ikese, *et al.*, 2021; Wu *et al.*, 2021; Lotah, 2022). Therefore, the measurement of toxic and heavy metals content in nails and hair plays an important role to understand as well as monitoring the impact of chemical environmental pollution on various inhabitants of a community. Furthermore, many researchers recommended the use of human hair for monitoring heavy metals. They found that fingernails and scalp hair have the useful as disclosing historical details on the dietary state and chronic diseases as well as exposure of individuals to chemical pollutants, especially toxic and heavy metals, including the effects of the workplace. In this context, fingernails and scalp hair have gained an irreplaceable position in monitoring the impact of chemical environmental pollution than other biological materials due to their easy collection without injuring the donor, elements accumulate over a long period without any changes, and the capacity of hair and nails to accumulate metals during extended periods, reflecting at least 1 year of exposure (Geier *et al.*, 2016; Mehra & Thakur, 2016; Ullah *et al.*, 2017; Rajfur *et al.*, 2018; Ocelić Bulatović *et al.*, 2019; Cabar *et al.*, 2020; Shin *et al.*, 2020; Ikese, *et al.*, 2021; Lotah, 2022). In addition, they found higher concentrations of residues are present in hair samples at a rate more than 10 times at least in urine or blood serum samples. For this reason, fingernails and scalp hair have been suggested as suitable measures of toxic and heavy elements in polluted areas. The purpose of the present study was to find out baseline information

Eida Mohammed Alshammari\*

Department of Chemistry, College of Science, University of Hail, Hail, Saudi Arabia.

\*E-mail: eida.alshammari@uoh.edu.sa



about the level of toxic and heavy elements (Pb, Zn, Cd, and Ni) in fingernails and scalp hair of autoworkers in Saudi Arabia using ICP-MS. No information has been reported and/ or available to evaluate the exposure of these metals in autoworkers in our area. Within this context, our study mainly aimed at the relationship between the workplace, such as autoworkers, and the presence of toxic and heavy metal levels in the hair and nail. Therefore, this study focused on the effect of autoworker's workplace and years of working on the accumulation of toxic and heavy metals in the hair and nails of workers. On the other hand, the auto workers chosen for this study were in direct contact with thinner, grease, petrol, diesel, etc. The obtained data can be lead to develop and help to find an easy and better diagnostic assessment for heavy metals toxicity particularly as a screening program for autoworkers and others.

## Materials and Methods

All reagents used in this work are AR grade and there is no need for further purification. Deionized water with  $18\text{ M}\Omega\text{ cm}^{-1}$  was used to prepare all standard solutions, dilute samples, and wash all tools and glassware throughout. Hydrogen peroxide ( $\text{H}_2\text{O}_2$ , 35%), and nitric acid ( $\text{HNO}_3$ , 69%) were obtained from Sigma-Aldrich. All plastic and glassware used were washed and cleaned with dilute nitric acid, and air-dried before use.

### *Samples Collection*

Samples in this study were collected from different sites in Hail city, the northern province of Saudi Arabia. The fresh samples (fingernails and scalp hair) were collected from 83 exposed subjects (autoworkers) and 24 office workers (control) using sterilized scissors and nail clippers of volunteers across several workplaces at the age of 18 to 60 years old males. All the individuals are not allowed to cut their fingernails and scalp hair for about 60 days before sampling. The study was conducted in accordance with the Declaration of Helsinki and approved by the Research Ethics Committee of Hail University, Hail, Saudi Arabia. Scalp hair samples were obtained from workers who did not have colored or treated hair and taken from the nape of the head as near as possible to the scalp. Nail samples were collected from all 10 fingers. In this context, all samples were cut into pieces as small as possible. Samples were classified according to the type of workplace, stored at about  $5^\circ\text{C}$  until digestion and analysis processes. In this context, samples were collected from healthy volunteers who unpolluted workplace sites (control), for purpose of comparison. Before samples collection, the hands of all autoworkers were craftily washed with medicated soap metal-free and then dried to remove any external contamination by using sterile tissue paper. In order not to affect the determination data, care was taken not to have the hair dyed while collecting the samples for study. The collected samples were kept to seal tightly coded polythene bags in our laboratory for pre-treatment. Before sample collection, questionnaire surveys were performed for all participants containing highlights of information such as age, home address, nutritional habits, smoking water source, total monthly income, current health with regard to chronic diseases or consequently drugs treatment, smoking habits, and occupational

exposure to heavy metals. Information was obtained through multiple-choice questions.

### *Washing Procedure*

The key aim of the washing process is to remove the external pollutants from hair and nails pre-digestion technique to have a true level of endogenous metals. In the other words, it will eliminate only the surface external pollutants without extracting elements from the samples or depositing elements on them. The results of existing and previous research have indicated that detergent washing is not suitable, where it reduced the metal concentrations of samples compared with the organic solvent washing. However, washing procedures do not essentially affect the heavy metals concentrations due to the strong complex of disulfide groups in the keratin proteins with these metals (Pozebon *et al.*, 2017; Alrobaian & Arida, 2019). Fingernails and scalp hair samples were initially washed with no-ionic detergent, Triton X-100, and then by  $\text{CH}_3\text{COCH}_3$  (acetone), after which washed by deionized  $\text{H}_2\text{O}$ . After washing processes, the fingernails and scalp hair samples were dried in an electric oven at about  $60^\circ\text{C}$  and protected in polyethylene bags until analysis. According to the International Atomic Energy Agency (Gönener *et al.*, 2020; Lotah *et al.*, 2022), this method has been come to be one of the most successful washing procedures in removing external trace elements and preserving internal trace elements.

### *Sample Preparation for Trace Element Analysis*

After the drying process, each investigated sample (0.5 g of fingernails or scalp hair) placed in a Teflon container was individually digested in the microwave digestion system with a mixture of 7 mL of concentrated  $\text{HNO}_3$  and 3 mL of  $\text{H}_2\text{O}_2$ . In this system, the temperature ramped to  $200^\circ\text{C}$  for 10 minutes, after which held for 10 minutes, and then cooled to reach room temperature. This method of digestion includes minimum sample handling steps which lead to minimizing the sample contamination besides a high large number of samples, which lead to improving the overall throughput and the reliability of the suggested analysis protocol. The microwave procedure operated according to the report by the US Environmental Protection Organization (USEPA) (Alrobaian & Arida, 2019). After cooling of the digestion vessel, the resulting clear solution was quantitatively diluted with deionized water and an aliquot was inserted into the cell for analysis using ICP-MS. A blank sample, containing deionized water was also used under the same conditions. The investigated toxic and heavy metals were measured by inductively coupled Plasma mass spectrometer- ICPMS2030 (Shimadzu- Japan) under optimized plasma conditions in all digested samples and blanks as well. The concentration of the investigated metals was measured automatically using the standard graph, which was prepared under the same plasma conditions. The system was adjusted to triplicate determined of the samples as well as the automatically calculated both correlation coefficient and the relative standard deviation, which found to be 0.99998 and  $< 2\%$ , respectively. The measurement method (ICP-MS) was validated using working calibration solutions of the under-investigated metal ions (Pb, Zn, Cd, and Ni). Standard solutions of all the metals investigated were prepared using stepwise dilutions of certified standard stock

solutions. These solutions for calibration of all the metals investigated were prepared from ultra-grade a 1000 mg/L ICP multielement solution (Merck, Milan, Italy), using stepwise dilutions of certified standard stock solutions. The procedure validation included linearity, accuracy, and precision.

## Results and Discussion

Each work environment is unique and the degree of hazards it poses to the workers. This study revealed the risks of toxic and heavy metals coming from highly polluted environments and the workers in these places are directly in contact with various pollutants and have different health problems. Heavy metal pollution in the workplace environment has come to be a worldwide problem. In this context, the use of nails and hair analysis for toxic and heavy metals has been recommended by the World Health Organization (WHO), due to the dependence of toxic and heavy metals levels in them on environmental exposure. In this work, we determine the concentration of toxic and heavy metals (Pb, Zn, Cd, and Ni) in fingernails and scalp hair samples of volunteer workers and controls (no direct metal exposure) by ICP-MS technique to estimate any relation between heavy metals concentration and environmental and/or occupational exposure. The ICP-MS measurement was validated by analysis of various standard samples and spiking are known concentrations of toxic and heavy metals, which showed good accuracy, linearity, precision, and lower limit of detection. The measurements were performed in triplicate and the mean was automatically calculated. Many researchers reported that there is a relationship between the concentration of trace elements in human tissues and many dangerous human diseases such as cancer, mutagenicity, neurobehavioral decrements, cardiovascular diseases, gastrointestinal irritation, and immune dysfunction which maybe lead to deaths (Pozebon *et al.*, 2017; Gönener *et al.*, 2020).

### Nails and Hair Samples Analysis Results

In the current study, bio-monitoring of four heavy metals was assessed by ICP-MS after microwave-assisted sample digestion with HNO<sub>3</sub> in the whole nails and hair samples (total n= 107) volunteer workers and controls (no direct metal exposure). The volunteers were divided into two groups: control (n=24, 22.4 %), and workers (n=83, 77.6 %). In this context, the volunteer workers were categorized according to the types of occupations into four sets, namely, car mechanics workers (n= 21), and auto welding workers (n=20), repair car tires workers (n= 22), and washing and oil change services center workers (n=20). For comparison, the tested metals (Pb, Cd, Zn, and Ni) were also measured in whole fingernails and scalp hair samples (n=24) of control. The control volunteers were classified according to the ages as well as exposure duration, Year (y) as shown in **Table 1**. The concentration ranges and averages of Cd, Ni, Zn, and Pb in fingernails of unpolluted workplace site (control) are 0.28 - 0.56(0.38), 4.67 -10.12(7.04), 57.34 - 107.44 (81.87), and 8.45 - 12061(10.27), while for scalp hair samples are 0.10 - 0.43 (0.23), 3.12 - 9.14(6.01), 50.56 - 105.61(73.84), and 6.54 - 13.90(9.41), respectively, **Table 1**. These obtained results are consistent with that reported for other parallel studies and are they are agreement the international tolerance levels (Nakaona *et al.*, 2020; Yin *et al.*, 2021). In this context, the levels of these toxic heavy metals in a biological specimen such as fingernails and scalp hair should be below the international tolerance levels as far as possible. However, environmental Pollutants have come to play an important role to introduce these elements into human biological systems, and therefore the term tolerant levels should be used instead of reference values (Alrobaian & Arida, 2019; Nakaona *et al.*, 2020). On the other hand, the obtained results show that the levels of the studied metals of fingernails and scalp hair in unpolluted workplace site volunteers reasonably fluctuated within a relatively narrow range for a given element. Indeed, the relative fluctuation in the obtained values in each studied metal level reflects the variety in culture, age, environment, and habits (Nakaona *et al.*, 2020).

**Table 1.** Heavy metals levels ( $\mu\text{g/g}$ ) in fingernails and scalp hair samples of unpolluted workplace site (control)

No	Exposure duration, Year(y)	Age*	Cd		Ni		Zn		Pb	
			Nails	Hairs	Nails	Hairs	Nails	Hairs	Nails	Hairs
1	1	a	0.39	0.18	5.23	4.76	61.47	72.31	8.45	6.54
2	2	a	0.32	0.20	4.87	3.12	57.34	50.56	9.33	11.30
3	4	b	0.28	0.19	7.24	6.68	78.35	69.57	8.97	6.86
4	4	b	0.29	0.19	4.67	3.42	94.24	80.57	11.43	8.59
5	5	b	0.31	0.24	6.45	4.96	63.26	47.83	9.47	11.45
6	7	b	0.35	0.17	6.86	5.39	78.35	80.56	9.45	11.98
7	9	b	0.40	0.13	8.32	8.21	81.34	69.58	10.34	8.45
8	9	b	0.32	0.11	6.78	7.12	90.85	83.35	9.46	7.67
9	10	b	0.28	0.19	5.91	4.35	84.74	68.34	8.85	10.79
10	10	c	0.42	0.31	6.47	3.67	90.07	78.48	12.23	7.56
11	10	c	0.39	0.25	5.86	6.33	107.44	90.35	10.36	8.98
12	10	c	0.48	0.32	8.33	5.95	88.25	93.47	8.58	5.97
13	12	c	0.29	0.21	7.45	5.92	78.91	56.79	9.85	7.93
14	12	c	0.28	0.10	4.79	4.23	68.36	54.88	10.47	8.85
15	15	c	0.44	0.22	7.60	6.87	70.34	78.63	8.49	10.76
16	15	c	0.51	0.43	6.44	4.86	90.36	78.57	11.86	9.34

17	16	c	0.31	0.15	7.55	5.49	75.85	80.98	10.38	12.87
18	18	d	0.42	0.23	8.98	9.12	87.34	60.86	11.84	8.77
19	20	d	0.46	0.35	8.95	7.23	69.37	54.61	9.83	7.85
20	22	d	0.28	0.11	5.49	4.87	79.48	64.84	11.62	9.97
21	27	d	0.34	0.19	6.15	5.98	90.35	76.84	9.37	10.89
22	30	e	0.56	0.41	8.58	9.14	98.56	75.30	12.54	8.56
23	37	e	0.49	0.31	9.94	7.69	92.36	105.61	10.31	13.90
24	38	e	0.51	0.36	10.12	8.93	87.98	99.44	12.97	9.93

\* Age: a = ≤ 20 y, b = 21-30 y, c = 31-40 y, d = 41-50 y, e = 51-60 y.

**Table 2** shows the concentration of toxic and heavy metal (Pb, Cd, Zn, and Ni) in fingernails and scalp hair samples of 21 car mechanics workers volunteers. It was found that the concentration ranges and averages of Cd, Ni, Zn, and Pb in fingernails of car mechanics workers were 0.41-0.80 (0.63), 4.12- 11.32 (7.96), 62.8-142.21 (95.66), 6.75-19.43(13.29), while the concentration ranges and averages of Cd, Ni, Zn and Pb in scalp hair of car mechanics workers were 0.12 - 0.76 (0.38), 4.10-13.20 (9.73), 72.65-142.21 (112.43), and 7.12-25.32(17.93), respectively.

From **Table 3**, it was found that the concentration ranges and averages of Cd, Ni, Zn, and Pb in fingernails of car mechanics workers were 0.35- 0.72 (0.49), 7.94 - 14.96 (11.07), 83.65 - 150.70 (109.78), and 7.10 - 17.54(12.27), while the concentration ranges and averages of Cd, Ni, Zn and Pb in scalp hair of car mechanics workers were 0.39 - 0.65 (0.50), 8.21-13.27 (10.90), 88.71-16121 (126.95), and 7.82 - 19.21(14.27), respectively.

The investigated toxic heavy elements were also determined fingernails and scalp hair samples of 22 repair car tire workers

volunteers. The data obtained are summarized in **Table 4**. It was found that the concentration ranges and averages of Cd, Ni, Zn, and Pb in fingernails of car mechanics workers were 0.38-0.80 (0.61), 9.20-15.96 (12.88), 61.5-119.61 (97.78), and 6.98-16.86(11.65), while the concentration ranges and averages of Cd, Ni, Zn and Pb in scalp hair of car mechanics workers were 0.41-0.91 (0.73), 8.79-16.88 (12.99), 66.21-126.51 (101.09), and 7.54-15.94(12.03), respectively.

**Table 5** summarizes the concentration of the investigated toxic and heavy metals fingernails and scalp hair samples of 20 washing and oil change services center workers volunteers. The concentration ranges and averages of Cd, Ni, Zn and Pb in fingernails of car mechanics workers were 0.43 -0.71 (0.55), 5.43 - 17.95 (11.81), 72.6 - 144.6 (104.37), and 7.98 - 18.76(12.07), while the concentration of these metals hair samples were 0.48 - 0.87 (0.66), 6.32 - 18.79 (12.49), 77.43 - 133.62 (108.58), and 8.32 - 20.21(14.68), respectively.

**Table 2.** Heavy metals levels (µg/g) in fingernails and scalp hair samples of car mechanics workers

No	Exposure duration, Year(y)	Age	Cd		Ni		Zn		Pb	
			Nails	Hairs	Nails	Hairs	Nails	Hairs	Nails	Hairs
1	2	a	0.41	0.21	5.81	4.77	62.8	81.30	8.21	7.12
2	3	b	0.44	0.12	4.88	4.10	71.24	79.73	6.75	7.79
3	3	b	0.41	0.14	5.76	5.13	78.94	72.65	7.42	10.34
4	4	b	0.53	0.16	4.12	6.75	70.57	86.77	7.89	8.98
5	5	b	0.57	0.14	5.72	6.81	90.23	90.55	9.31	13.43
6	6	b	0.47	0.27	6.75	8.92	79.89	88.94	8.51	16.43
7	6	b	0.62	0.26	5.53	8.44	90.21	102.45	13.75	15.90
8	7	b	0.54	0.25	5.98	7.99	86.94	99.64	11.86	19.76
9	7	b	0.71	0.23	7.32	10.32	79.73	112.64	9.86	18.43
10	9	b	0.65	0.32	6.88	9.66	85.87	107.93	13.72	21.67
11	10	c	0.46	0.39	8.71	11.05	108.12	123.54	12.89	19.67
12	10	c	0.48	0.28	7.56	10.55	94.55	115.80	12.64	22.35
13	10	c	0.61	0.43	9.14	11.79	88.58	119.50	16.87	22.89
14	11	c	0.58	0.39	8.31	10.66	112.06	128.40	14.51	20.12
15	13	c	0.79	0.45	9.54	11.88	94.67	132.40	16.21	18.34
16	16	d	0.68	0.54	10.12	11.69	109.32	134.75	15.95	17.98
17	17	d	0.80	0.63	9.69	12.32	98.52	125.95	19.43	19.37
18	22	d	0.76	0.76	11.32	12.90	126.41	134.98	17.86	23.43

19	20	e	0.89	0.72	10.31	12.33	122.58	140.20	19.31	22.93
20	24	e	0.85	0.70	10.88	13.20	129.82	140.77	17.54	24.21
21	29	e	0.88	0.74	10.63	13.10	127.93	142.21	18.76	25.32

\* Age: a = ≤ 20 y, b = 21-30 y, c = 31-40 y, d = 41-50 y, e = 51-60 y

**Table 3.** Heavy metals levels ( $\mu\text{g/g}$ ) in fingernails and scalp hair samples of auto welding workers

No	Exposure duration, Year(y)	Age	Cd		Ni		Zn		Pb	
			Nails	Hairs	Nails	Hairs	Nails	Hairs	Nails	Hairs
1	1	a	0.37	0.39	8.89	9.45	83.65	90.43	8.21	8.97
2	1	a	0.43	0.42	7.94	8.86	99.64	88.71	9.71	7.82
3	3	a	0.35	0.39	8.61	8.21	91.63	112.23	7.10	9.63
4	5	b	0.39	0.46	7.89	10.43	113.7	126.91	9.42	7.96
5	5	b	0.46	0.46	9.71	9.23	94.95	107.68	8.75	10.55
6	7	b	0.38	0.49	10.43	9.78	89.91	99.65	9.07	12.43
7	10	b	0.47	0.43	8.57	8.97	99.72	130.66	12.15	11.85
8	11	b	0.43	0.40	8.94	9.97	114.71	125.56	10.13	13.73
9	13	c	0.59	0.47	11.89	8.95	98.95	114.83	9.54	15.74
10	14	c	0.39	0.54	10.75	10.55	105.87	109.59	13.81	14.86
11	14	c	0.47	0.51	12.70	11.23	112.80	125.65	12.43	15.97
12	17	c	0.55	0.46	9.95	10.65	98.86	114.84	9.96	14.85
13	18	c	0.47	0.58	13.71	12.16	96.06	132.88	14.83	16.72
14	20	c	0.46	0.48	10.86	11.85	122.75	140.32	12.66	16.79
15	20	c	0.57	0.54	12.08	12.76	102.24	134.86	16.70	15.55
16	21	c	0.63	0.58	14.06	13.53	90.98	156.98	15.12	17.89
17	20	d	0.51	0.58	13.81	11.92	141.81	149.98	15.79	17.43
18	24	d	0.62	0.65	12.95	13.27	138.9	161.21	17.54	18.89
19	25	d	0.58	0.61	14.00	13.08	150.70	155.90	15.85	18.52
20	28	e	0.72	0.63	13.75	13.14	147.90	160.13	16.75	19.21

\* Age: a = ≤ 20 y, b = 21-30 y, c = 31-40 y, d = 41-50 y, e = 51-60 y

The data of unpolluted workplace sites (**Table 1**) and autoworkers (**Tables 2-5**) were compared. In general, it was noted that the concentrations of toxic and heavy metals were higher in autoworkers samples than that in control samples, and this is in agreement with that reported by others for similar studies (Al-Easawi *et al.*, 2017; Wang *et al.*, 2009). The impact of age and exposure duration per year on element concentrations was investigated. It was found that clear relation between the concentrations of toxic and heavy metals influence age and exposure duration. Consequently, many factors contributed to introducing toxic and heavy metals into biological systems; this means the term of tolerant levels suitable for use instead of reference values (Pozebon *et al.*, 2017; Alrobaian & Arida, 2019; Lotah *et al.*, 2022). In this context, the average levels of Cd were found to be slightly increased upon repair car tires workers (**Table 4**) compared with other autoworkers (**Tables 2, 3 and 5**). On the other hand, the concentrations of Zn in auto welding workers (**Table 3**) were found to be significantly higher compared with that in other autoworkers (**Tables 2, 4 and 5**). The concentration average levels of Pb and Ni in the washing and oil change services center workers' fingernails and scalp hair volunteers (**Table 5**) were found to be slightly increased compared with other

autoworkers (**Tables 2-4**). A comparison of the toxic and heavy elements' concentrations in fingernails and scalp hair of autoworkers and control volunteers was carried out by using Microsoft Excel of Office 365 to calculate the T-test. It was determined that didn't normal distribution, a value  $p < 0.05$  was considered to be significant. These results are consistent with other similar studies as well as below the international tolerance levels (Gönerer *et al.*, 2020; Nakaona *et al.*, 2020). As a result of this study, a statistically significant relationship has been found between autoworkers and the toxic and heavy metal levels in the fingernails and scalp hair of volunteers. Indeed, once a toxic and heavy element is absorbed, it distributes in tissues and organs, later on, excretion typically occurs primarily through kidneys and digestive tract, some elements tend to persist in some storage sites, like the liver, bones, hair, and kidneys, for many years (Pozebon *et al.*, 2017; Alrobaian & Arida, 2019; Lotah *et al.*, 2022). In general, the sequences for the concentration of investigated metals in groups (unpolluted workplace site (control), car mechanics workers, car mechanics workers, and washing and oil change services center workers) are the following:  $\text{Zn} > \text{Pb} > \text{Ni} > \text{Cd}$ , as shown in (**Tables 1-3 and 5**) while the repair car tires workers are  $\text{Zn} > \text{Ni} \geq \text{Pb} > \text{Cd}$ , as shown in **Table 4**. As a result of this research,

it has been revealed that there is a statistically significant relationship between environmental including the effects of the

workplace of autoworkers, and the toxic and heavy metal levels detected in their fingernails and scalp hair.

**Table 4.** Heavy metals levels ( $\mu\text{g/g}$ ) in fingernails and scalp hair samples of repair car tires workers

No	Exposure duration, Year(y)	Age	Cd		Ni		Zn		Pb	
			Nails	Hairs	Nails	Hairs	Nails	Hairs	Nails	Hairs
1	2	a	0.38	0.45	9.15	8.79	70.14	69.37	6.98	7.20
2	2	a	0.43	0.51	10.12	9.43	61.5	71.22	8.43	6.54
3	5	b	0.58	0.41	11.62	8.95	83.52	66.21	8.98	8.41
4	7	b	0.61	0.64	9.20	10.68	98.72	81.80	11.21	11.25
5	7	b	0.55	0.59	12.61	9.87	85.74	79.77	10.95	10.55
6	8	b	0.64	0.68	12.84	12.33	78.91	91.87	11.79	11.41
7	10	b	0.49	0.73	10.77	12.82	89.63	94.71	9.75	10.60
8	11	b	0.61	0.68	14.08	11.85	95.11	98.70	13.06	11.14
9	11	b	0.40	0.72	11.83	13.22	90.66	91.85	8.89	9.69
10	13	b	0.54	0.79	13.70	12.14	113.70	99.62	12.85	11.85
11	12	c	0.71	0.72	11.72	11.91	102.45	105.53	10.76	13.82
12	18	c	0.49	0.88	13.05	13.76	88.69	108.80	9.63	12.86
13	19	c	0.68	0.82	12.92	12.60	90.63	98.97	11.85	12.52
14	19	c	0.57	0.78	15.48	14.81	104.76	113.23	14.21	14.10
15	20	c	0.74	0.85	13.38	13.93	98.06	106.60	12.09	13.61
16	22	c	0.64	0.84	13.87	14.11	109.66	114.85	14.94	11.76
17	23	c	0.80	0.78	15.47	15.21	112.46	119.65	14.87	14.85
18	21	d	0.64	0.79	14.50	14.47	99.64	113.82	16.86	12.98
19	26	d	0.77	0.77	13.76	15.48	119.58	121.85	16.65	13.64
20	30	d	0.69	0.91	14.20	16.67	119.61	126.31	15.08	15.94
21	29	e	0.72	0.88	13.21	15.93	109.60	122.89	13.93	15.20
22	31	e	0.78	0.84	15.96	16.88	113.87	126.51	14.08	14.77

\* Age: a =  $\leq 20$  y, b = 21-30 y, c = 31-40 y, d = 41-50 y, e = 51-60 y

**Table 5.** Heavy metals levels ( $\mu\text{g/g}$ ) in fingernails and scalp hair samples of washing and oil change services center workers

No	Exposure duration, Year(y)	Age	Cd		Ni		Zn		Pb	
			Nails	Hairs	Nails	Hairs	Nails	Hairs	Nails	Hairs
1	3	a	0.53	0.49	6.08	6.64	80.06	75.81	7.98	8.08
2	3	b	0.48	0.48	7.35	6.32	72.6	77.23	8.63	9.11
3	2	b	0.50	0.57	5.43	6.77	81.16	77.43	10.64	8.32
4	4	b	0.48	0.49	7.75	6.94	79.95	82.32	9.72	8.87
5	7	b	0.43	0.62	8.53	8.21	88.65	88.64	10.61	8.62
6	8	b	0.51	0.59	7.84	7.93	97.43	94.72	7.88	12.28
7	10	b	0.48	0.54	9.07	10.21	90.66	112.85	8.91	14.77
8	10	b	0.56	0.68	14.57	9.88	112.71	99.53	12.14	14.54
9	11	b	0.47	0.63	12.73	12.32	105.32	116.64	9.05	15.82
10	12	b	0.61	0.7.2	10.70	13.21	86.94	120.21	11.09	16.81
11	12	b	0.49	0.74	8.59	12.89	99.06	112.62	9.74	18.60
12	13	b	0.54	0.62	13.82	14.11	92.94	118.51	13.73	17.67
13	9	c	0.63	0.77	10.84	13.54	131.71	108.98	11.82	15.65
14	12	c	0.54	0.67	15.82	15.88	112.61	116.73	13.79	12.39
15	13	c	0.61	0.65	17.95	18.79	98.79	124.71	12.77	17.11
16	13	c	0.56	0.74	17.09	16.76	133.71	121.43	14.81	16.87

17	21	d	0.71	0.82	13.85	15.89	116.86	129.64	14.89	19.21
18	18	d	0.64	0.79	14.71	16.84	136.82	133.62	17.63	18.53
19	19	d	0.70	0.85	15.80	17.54	144.65	129.66	16.88	20.21
20	28	e	0.60	0.87	17.73	18.54	124.70	130.41	18.76	20.14

\* Age: a = ≤ 20 y, b = 21-30 y, c = 31-40 y, d = 41-50 y, e = 51-60 y

## Conclusion

The levels of certain toxic and heavy metals (Pb, Zn, Cd, and Ni) were determined by using ICP-MS in fingernails and scalp hair of autoworkers in Saudi Arabia to study exposure to chemical pollutants metals. It was found that the concentrations of toxic and heavy metals were higher in autoworkers samples than that in control samples Toxic and heavy metals levels increase with increase the age years of autoworkers, epically for Pb and Cd. The obtained result proved that fingernails and scalp hair could be used as a good biological indicator for the assessment of toxic heavy metal pollution. On the other hand, this research shows that there is a statistically significant relationship between the type of work (autoworkers) or workplace and the heavy metal concentration detected in fingernails and scalp hair samples, which represent a very important guide for the labor and health organizations.

**Acknowledgments:** The author extends her appreciation to individuals that filled my questionnaire and volunteered their fingernails and scalp hair samples to be used for this study without any reservation. I wish also to express my thanks to anonymous reviewers, for their constructive suggestions.

**Conflict of interest:** None

**Financial support:** None

**Ethics statement:** None

## References

- Almeida-da-Silva, C. L. C., Dakafay, H. M., O'Brien, K., Montierth, D., Xiao, N., & Ojcius, D. M. (2021). Effects of electronic cigarette aerosol exposure on oral and systemic health. *Biomedical Journal*, *44*(3), 252-259.
- Al-Muzafar, H. M., & Al-Hariri, M. T. (2021). Elements alteration in scalp hair of young obese Saudi females. *Arab Journal of Basic and Applied Sciences*, *28*(1), 122-127. doi:10.1080/25765299.2021.1911070
- Alrobaian, M., & Arida, H. (2019). Assessment of heavy and toxic metals in the blood and hair of Saudi Arabia smokers using modern analytical techniques. *International Journal of Analytical Chemistry*, *2019*.
- Amartey, E. O., Asumadu-Sakyi, A. B., Adjei, C. A., Quashie, F. K., Duodu, G. O., & Bentil, N. O. (2011). Determination of heavy metals concentration in hair Pomades on the Ghanaian market using atomic absorption spectrometry technique. *British Journal of Pharmacology and Toxicology*, *2*(4), 192-198.
- Cabar, H. D., Ersoy Karaçuha, M., & Yilmaz, M. (2020). The Interaction Between Concentration of Heavy Metal-Trace Elements and Non-Smoking Status of Adolescents in Sinop (Turkey). *Biological Trace Element Research*, *194*(1), 105-114.
- Geier, D. A., Kern, J. K., Hooker, B. S., Sykes, L. K., & Geier, M. R. (2016). Thimerosal-preserved hepatitis B vaccine and hyperkinetic syndrome of childhood. *Brain Sciences*, *6*(1), 9.
- Gönener, A., Karaçuha, M. E., Cabar, H. D., Yilmaz, M., & Gönener, U. (2020). The relationship between dietary habits of late adolescent individuals and the heavy metal accumulation in hair. *Progress in Nutrition*, *22*, 146-155.
- Ikese, C. O., Adie, P. A., Adah, C., Amokaha, R., Abu, G., & Yager, T. (2021). Heavy metal levels in spent engine oils and fingernails of auto-mechanics. *Ovidius University Annals of Chemistry*, *32*(1), 28-32.
- Lemos, V. A., & de Carvalho, A. L. (2010). Determination of cadmium and lead in human biological samples by spectrometric techniques: a review. *Environmental Monitoring and Assessment*, *171*(1), 255-265.
- Lotah, H.N.A., Agarwal, A. K., & Khanam, R. (2022). Heavy metals in hair and nails as markers of occupational hazard among welders working in United Arab Emirates. *Toxicological Research*, *38*, 63-68.
- Massadeh, A., Gharibeh, A., Omari, K., Alomary, A., Tumah, H., & Hayajneh, W. (2010). Simultaneous determination of Cd, Pb, Cu, Zn, and Se in human blood of Jordanian smokers by ICP-AES. *Biological Trace Element Research*, *133*(1), 1-11. doi:10.1007/s12011-009-8405-y.
- Mehra, R., & Thakur, A. S. (2016). Relationship between lead, cadmium, zinc, manganese and iron in hair of environmentally exposed subjects. *Arabian Journal of Chemistry*, *9*, S1214-S1217.
- Mončilović, B., Prejac, J., Višnjević, V., Brundić, S., Skalny, A. A., & Mimica, N. (2016). High hair selenium mother to fetus transfer after the Brazil nuts consumption. *Journal of Trace Elements in Medicine and Biology*, *33*, 110-113.
- Nakaona, L., Maseka, K. K., Hamilton, E. M., & Watts, M. J. (2020). Using human hair and nails as biomarkers to assess exposure of potentially harmful elements to populations living near mine waste dumps. *Environmental Geochemistry and Health*, *42*(4), 1197-1209.
- Noreen, F., Sajjad, A., Mahmood, K., Anwar, M., Zahra, M., & Waseem, A. (2020). Human biomonitoring of trace elements in scalp hair from healthy population of Pakistan. *Biological Trace Element Research*, *196*(1), 37-46.
- Ocelić Bulatović, V., Mandić, L., Turković, A., Kučić Grgić, D., Jozinović, A., Zovko, R., & Govorčin Bajsić, E. (2019). Environmentally friendly packaging materials based on thermoplastic starch. *Chemical and Biochemical Engineering Quarterly*, *33*(3), 347-361.

- Peter, O. O., Eneji, I. S., & Sha'Ato, R. (2012). Analysis of heavy metals in human hair using atomic absorption spectrometry (AAS). *American Journal of Analytical Chemistry*, 3, 770-773.
- Pozebon, D., Scheffler, G. L., & Dressler, V. L. (2017). Elemental hair analysis: A review of procedures and applications. *Analytica Chimica Acta*, 992, 1-23.
- Rajfur, M., Świsłowski, P., Nowainki, F., & Śmiechowicz, B. (2018). Mosses as biomonitor of air pollution with analytes originating from tobacco smoke. *Chemistry-Didactics-Ecology-Metrology*, 23(NR 1-2), 127-136. doi:10.1007/s12011-019-01769-5
- Shin, W. J., Jung, M., Ryu, J. S., Hwang, J., & Lee, K. S. (2020). Revisited digestion methods for trace element analysis in human hair. *Journal of Analytical Science and Technology*, 11(1), 1-5.
- Silva, L. A., Robazzi, M. L., Assuncao, H. F., Dalri, R., Maia, L. G., Silverira, S., Mendonca, G. S., Rabahi, M. F., & Porto, C. C. (2018). Impact of environmentally pollution on carboxyhemoglobin levels among smoking and nonsmoking motorcycle taxi drivers. *Bioscience Journal*, 34(2), 477-485.
- Ullah, H., Noreen, S., Rehman, A., Waseem, A., Zubair, S., Adnan, M., & Ahmad, I. (2017). Comparative study of heavy metals content in cosmetic products of different countries marketed in Khyber Pakhtunkhwa, Pakistan. *Arabian Journal of Chemistry*, 10(1), 10-18.
- Vincenti, M., Salomone, A., Gerace, E., & Pirro, V. (2013). Role of LC-MS/MS in hair testing for the determination of common drugs of abuse and other psychoactive drugs. *Bioanalysis*, 5(15), 1919-1938.
- Wu, I. P., Liao, S. L., Lai, S. H., & Wong, K. S. (2021). The respiratory impacts of air pollution in children: Global and domestic (Taiwan) situation. *Biomedical Journal*, S2319-4170(21)00176-1. doi:10.1016/j.bj.2021.12.004
- Yin, J., Hou, W., Vogel, U., Li, X., Ma, Y., Wang, C., Wang, H., & Sun, Z. (2021). TP53 common variants and interaction with PPP1R13L and CD3EAP SNPs and lung cancer risk and smoking behavior in a Chinese population. *Biomedical Journal*, S2319-4170(21)00005-6. doi:10.1016/j.bj.2021.01.006