

# *Ocimum basilicum* L.: A Systematic Review on Pharmacological Actions and Molecular Docking Studies for Anticancer Properties

Islam Boulaares, Samir Derouiche\*, Janetta Niemann

Received: 02 November 2023 / Received in revised form: 01 March 2024, Accepted: 04 March 2024, Published online: 15 March 2024

## Abstract

Basil (*Ocimum basilicum* L) is a plant that belongs to the Lamiaceae family and is known for its pharmacological and therapeutic properties, as it is rich in active biological substances, which endow it with antioxidant and anti-inflammatory activities. Many people from all over the world use basil in traditional medicine, including North African countries, especially Algeria, which they use to treat many organic and infectious diseases. On the other hand, according to scientific reports, the *O. basilicum* plant contains most of the major and biologically active molecules such as flavonoids, phenols, tannins, steroids, glycosides, and reduced sugars. Pharmacological studies also indicate that the plant has antioxidant, anti-inflammatory, analgesic and antibacterial activities. In silico tools, a compound's ability to block a receptor can be successfully studied by combining many in silico approaches. There have been reports of natural chemicals of *O. basilicum* having anticancer action. In conclusion, *O. basilicum* may be one of the best sources of plant medicines that can be used to treat many acute and chronic diseases.

**Keywords:** Basil, Systematic review, Phytochemistry, Botany, Pharmacological activities, Molecular docking study

## Introduction

Research on herbal medicines has grown to be one of the most pressing scientific issues in the previous two decades (Chetehouna *et al.*, 2020). A variety of ailments has historically been prevented or treated using medicinal herbs. More than 1200 plants are utilized in traditional medicine across the globe, according to ethnopharmacological investigations (Derouiche, 2020). The use of medical and food plants is a major part of the medication management of so-called chronic diseases in some

### Islam Boulaares, Samir Derouiche\*

Department of Cellular and Molecular Biology, Faculty of Natural Sciences and Life, University of El-Oued, El-Oued 39000, Algeria.

Laboratory of Biodiversity and Application of Biotechnology in the Agricultural Field, Faculty of Natural Sciences and Life, University of El-Oued, El-Oued 39000, Algeria.

### Janetta Niemann

Department of Genetics and Plant Breeding, Poznań University of Life Sciences, Dojazd 11, 60-632 Poznań, Poland.

\*E-mail: dersamebio@gmail.com



© 2024 The Author(s). This is an **Open Access** article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0).  
<https://creativecommons.org/licenses/by/4.0/deed.en>

traditional non-industrialized civilizations (China, and several African and Latin American nations), according to Chetehouna *et al.* (2023). Bioactive compounds found in medicinal plants represent a variety of interests used in various fields. The secondary metabolites of these compounds are those that are most clearly demonstrated in the therapeutic sector (Tungmunthum *et al.*, 2018).

The public's historical usage of plants as medicine and confirmation of their pharmaceutical effectiveness encouraged their medical use. While their active compounds might treat illnesses and alleviate symptoms, However, the development of study into the elements in such plants demonstrates that their use extends beyond its use in folk medicine (Kalamartzis *et al.*, 2020). *Ocimum basilicum*, sometimes known as sweet basil, is one of the therapeutic plants. Many plants of the genus *Ocimum* have been utilized throughout history to cure a variety of illnesses and ailments. *O. basilicum* plays a significant role in this genus because of its many therapeutic characteristics (Purushothaman *et al.*, 2018).

In light of these data, this review aimed to identify the phytochemical contents and pharmacological effect of *Ocimum basilicum* L.

## Materials and Methods

For this review, the literature on the biological properties, secondary metabolites, and botanical description of basil was collected, examined, and summarized. All articles that have been published concerning this species have been collected using scientific search engines including Springer Link, Scopus, Science Direct, Wiley Online, Web of Science, PubMed, Scinder, and Google Scholar (e.g., WIPO, CIPO, USPTO). These search engines, as well as numerous patient offices, used to use Scopus, Wiley Online, Scifinder, and PubMed. It's common to hear the phrase "*Ocimum basilicum*," either by itself or in conjunction with the phrases "chemical substances" and "pharmacological activity." There were no restrictions on languages. The obtained data were identified and modified using their titles, abstracts, and contents. To discover if any other papers were pertinent, the reference lists of the papers that were retrieved were also examined.

## Results and Discussion

*Generalities and Geographic Distribution of Ocimum + L*

*Ocimum basilicum* L., also called sweet basil (Ahmed *et al.*, 2019), is among the aromatic plants (**Figure 1**) as it belongs to the *Lamiaceae* family. As it is characterized by its richness in chemicals (Rumengan *et al.*, 2019), it is known to be a plant used in medicine and food (Taha *et al.*, 2020). *Ocimum basilicum* L. is grown in soil and clearings Basil (*Ocimum basilicum* L.), is an herb that has been cultivated since ancient times, making it very well-known (Georgiadou *et al.*, 2018). It is also a plant that grows in warm regions, Africa, tropical regions, and some regions in Asia and South America, and it is very perennial (Choi *et al.*, 2019).



**Figure 1.** Basil leaf shape (Nadeem *et al.*, 2020)

#### Taxonomy and Botanical Description

*Ocimum basilicum* is an aromatic plant that grows all year round, reaching 0.3 to 0.5 meters in length, reaching 1 meter in some cases. 'Dark Garnet' is a cultivar of basil that has dark purple leaves and stems. Its leaves are oval, but its flowers are often wrinkled and white and pink. Its fruits contain small nuts, and when wet they are gummy (Burkil, 1995).

The branched herbaceous plant *Ocimum basilicum* L. has square, glabrous stems and branches that range in color from green to light purple. It typically stands between 0.6 and 0.9 meters tall. Basil leaves are simple, oppositely oriented, and range in length

from 2.5 to 5 cm on average. These longer leaves feature an oval form, a sharp apex, and a lobed or serrated edge (Nadeem *et al.*, 2020). Basil's taxonomy is regarded as being extensive and complicated (**Table 1**). According to Chowdhury *et al.* (2017), genetic diversity driven by cross-pollination and several environmental conditions is thought to be the cause of this taxonomic complexity. *Ocimum basilicum* L., sometimes known as Labiatae, is a member of the Lamiaceae family of mints, which is part of the Lamiales order (Azoz *et al.*, 2016; Idowu & Ozlegbe, 2017). One of the most significant species in the *Ocimum* genus, which has more than 60 species, is basil (Murali & Prabakaran, 2018).

**Table 1.** Taxonomy of Basil (Romanus *et al.*, 2020)

| Kingdom     | Plantae         |
|-------------|-----------------|
| Clade       | Asteroids       |
| Clade       | Eudicots        |
| Clade       | Tracheophytes   |
| Family      | Lamiaceae       |
| Genus       | Lamiales        |
| Species     | O. basilicum L  |
| Order       | Ocimum          |
| Common Name | Sweet basil     |
| Local Names | Ibibio/Efik-Iko |
| Yoruba      | Efinrin wewe    |
| Igbo        | Nchu- anwu      |
| Hausa-      | Daidoya tagida  |

#### Chemical Composition

According to Falowo *et al.* (2019), The basil plant contains variable amounts of nutrients such as glucides, protein, vitamins, fats, and some minerals such as potassium, iron, calcium, and magnesium. In addition, the basil plant contains several secondary metabolites such as saponins, alkaloids, flavonoids, and terpenoids. Terpenoids, alkaloids, flavonoids, and ascorbic acid (**Table 2**) have been identified (Rumengan *et al.*, 2019).

**Table 2.** Major constituents of *O. basilicum*

| Major constituents                                                                                                                                    | References                                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------|
| $\beta$ -Myrcene, 1,8-cineol, methyl cinnamate, Myrtenol, $\alpha$ -cubebene, $\beta$ -ocimene, caryophyllene, eugenol, et $\alpha$ -farnesene.       | (Abou El-Soud <i>et al.</i> , 2015)       |
| carvone and <i>iso</i> -pinocampnone.                                                                                                                 | (De-Martino <i>et al.</i> , 2009)         |
| Linalool and Methyl chavicol,                                                                                                                         | (Sajjadi, 2006)                           |
| $\alpha$ -cardinole, Linalool, $\delta$ -caridnene, $\alpha$ -selinène, $\alpha$ -bergamotène, $\beta$ -selinene, $\gamma$ guaiene, $\alpha$ -guaiene | (Kostic <i>et al.</i> , 2008)             |
| estragole and Linalool                                                                                                                                | (Martinez-Velazquez <i>et al.</i> , 2011) |

#### Therapeutic Effect and Economic Value of Sweet Basil


According to Bahcesular *et al.* (2020), The plant (*Ocimum basilicum* L.) can be considered one of the most important aromatic plants that have medicinal importance that can be grown and marketed and culinary relevance in various regions of the world because it includes significant components (Kalamartzis *et al.*, 2020). Due to a high level of secondary metabolites (Dörr *et al.*, 2020), basil leaves have substantial antioxidant and

antibacterial properties that are used in folk medicine to treat a wide range of illnesses (Ahmed *et al.*, 2019). According to numerous studies (Akbari *et al.*, 2019), basil essential oil possesses fungistatic and insecticidal properties. It can also be utilized as a flavoring agent in cosmetics, medicine, and food (Rezzoug *et al.*, 2019). All parts of Basil have historically been used as medication to treat digestive issues, kidney dysfunction, warts, worms, diarrhea, and migraines (Falowo *et al.*, 2019). Several studies conducted by various researchers have also

demonstrated that the essential oils isolated from *Oryza basilicum* exhibit outstanding antibacterial action against a variety of microbes, including fungi, bacteria, and Gram-positive and Gram-negative bacteria (Matasyoh *et al.*, 2008).

In the present review, we indicate that the *Ocimum basilicum* is richer in secondary metabolites like flavonoids, steroids, Phenols, Catechic Tannin, Saponoside, Carbohydrates, and Alkaloids. This is in the same line with the study of Tariq *et al.* (2016), *O.*


*basilicum* is considered a plant rich in active substances such as phenolic compounds, glycosides, terpenes, saponins, flavonoids, and many other substances, makeup flavonoids (**Figure 2**). Numerous advantageous physiological activities are exhibited by flavan-3-ols, flavonoids, and other derivatives (Panche *et al.*, 2016). Several flavonoid characteristics, such as antioxidant, anti-inflammatory, antiplatelet activity, and cardio-protective effects (Corti *et al.*, 2009).



**Figure 2.** Some flavonoid basic skeletons with hydroxyl groups (Katz *et al.*, 2011)

It has been proven that secondary metabolites have effective anti-inflammatory activity, such as terpenes, and antioxidant activity, such as flavonoids (**Figure 3**), which qualifies them to be a treatment for many cancers and osteoporosis, and the effects of coronary diseases can also be reduced by using phenols. (Derouiche *et al.*, 2020). Regarding the antioxidant activity, several studies by El-Dakar *et al.* (2015) show that Basil has a high-power antioxidant. A study by Harnafi *et al.* (2009) suggested that phenolic compounds are predominant in *O. basilicum* such as tannins and flavonoids. On the other hand, Due

to its phenolic and aromatic components, *O. basilicum* has demonstrated antioxidant activity (Gebrehiwot *et al.*, 2016). The aqueous extracts of Basil had a very high level of free radical scavenging antioxidant activity. Basil is a plant that is frequently used as medicine (Constantinescu & Adriana, 2019). In 2020, Pistelli *et al.* conducted another study and discovered that basil has phenolic antioxidant chemicals. Flavonoids have antioxidant properties that prevent oxidative damage through a variety of processes (Rameshrad *et al.*, 2015).




**Figure 3.** Reaction of antioxidant activity of flavonoids (Katz *et al.*, 2011)

Basil was additionally widely mentioned as a typical anti-inflammatory. We conducted several studies on human erythrocytes exposed to a pro-inflammatory substance to confirm this activity and assess the mechanism at play. The study highlights the significance of *O. basilicum* leaves' anti-inflammatory properties (Ahmad *et al.*, 2015). Pro-inflammatory cytokines play essential roles in the inflammatory response like IL-1 antagonist receptor, IL-6, IL-10, and IL-112 are antiinflammatory cytokines (Güez *et al.*, 2017). The main mechanism of anti-inflammatory action is the inhibition of prostaglandin formation resulting from the inactivation of cyclooxygenase, the enzymes of arachidonic acid metabolism, and lipoxygenase (Barbalho *et al.*, 2011).


Sweet basil showed strong antibacterial resistance to a variety of pathogens. Moderate antibacterial activity was displayed by *O. basilicum*. In contrast, Gebrehiwod *et al.* (2015) revealed that the Basil and its oil essential primary component, linalool, have antifungal properties and that Gram-positive bacterial strains were more sensitive to them than their counterparts.

In silico tools, a compound's ability to block a receptor can be successfully studied by combining many in-silico approaches. There have been reports of natural chemicals having anticancer capabilities, but it's still unclear how they work. To shed light on potential mechanisms of the anticarcinogenic interaction between receptor proteins and enzymes, theoretical docking studies targeting lipoxygenase-5 (LOX) and cyclooxygenase-2 (COX-2) have been conducted. In the production of prostaglandin E2, the primary enzyme is COX-2 (Bourzikat *et al.*, 2022). Given the clear association between inflammation and carcinogenesis, prospective COX-2 inhibitors may also qualify as chemopreventive medicines for cancer (Gurpinar *et al.*, 2013). According to docking research, isoeugenol (PDB ID: 4COX) sits in the active site of COX-2 with an orientation comparable to that of indomethacin (Figure 4). The pocket is composed of some amino acids. The terminal carbon atom is traveling in the direction of Leu384 (Redzicka *et al.*, 2023).




**Figure 4.** Interaction between the active site of COX-2 and Isoeugenol (Zarlaha *et al.*, 2014).

According to a different study, the molecular docking method was used to evaluate the compounds' propensity for binding to DNA gyrase B's active site. The photos of the 2D and 3D ligand-AA interactions for the two compounds under study are displayed in Figure 5. When the residues interacting with these compounds are compared to those in the active site. This supports the theory that these chemicals have the solidest inhibitory influence on *E. coli* (Eswaramoorthy *et al.*, 2021). Van der Waals connections from the AAs Glu50 and Val120 surround the trans-anethole, while the compound's benzene ring interacts with an alkyl bond from Val 71 through a pi-pi cationic link formed by Gly77. The lone hydrogen bond in the molecule is formed by Asn46 with the oxygen atom; Asp73 forms the electron-accepting carbon bond with a carbon-hydrogen. Furthermore, there are notable interactions between methyl chavicol and trans-anethole. The hydrophobic chains comprising the amino acids, as well as the alkyl bonds with Val167 and Ile94, envelop this molecule. By completing a hydrogen bonding contact with the crucial AAs Thr165, the molecule is expanded into the active site outwards (Kiessling & Diehl, 2021).



a)



**Figure 5.** Representation by 3D and 2D of the docking pose of DNA gyrase B and *trans*-anethol (Qasem *et al.*, 2023).

Isoeugenol's significant cytotoxic action can be partially explained by its ability to inhibit the COX and LOX enzymes, as well as by *trans*-anetholes at the active site of DNA gyrase B of *E. Coli*, according to a study of (Das Chagas Pereira and Mendes, 2020). In tests on cell lines, caffeine performs on par with or better than cisplatin (Teng *et al.*, 2020).

## Conclusion

The fact that the plant's aqueous extract contains a large number of bioactive molecules, as well as anti-inflammatory, anti-oxidants, and anticancer compounds, shows that basil has a strong biological impact on lowering several diseases associated with these mechanisms.

**Acknowledgments:** None

**Conflict of interest:** None

**Financial support:** None

**Ethics statement:** This study was approved by the Faculty of natural and life Sciences, El-Oued University.

## References

- Ahmad, Ch. M., Naz, S. B., Sharif, A., Akram, M., & Saeed, M. A. (2015). British biological and pharmacological properties of the sweet basil (*Ocimum basilicum*). *Journal of Pharmaceutical Research*, 7(5), 330-339.
- Ahmed, A. F., Attia, F. A. K., Liu, Z., Li, C., Wei, J., & Kang, W. (2019). Antioxidant activity and total phenolic content of essential oils and extracts of sweet basil (*Ocimum basilicum* L.) plants. *Food Science and Human Wellness*, 8(3), 299-305.
- Akbari, A. G., Binesh, S., Ramshini, H., Soltani, E., Amini, F., & Marieh, S. (2019). Mirfazeli selection of basil (*Ocimum*

*basilicum* L.) full-sib families from diverse landraces. *Journal of Applied Research on Medicinal and Aromatic Plants*, 12, 66-72.

- Azoz, S. N., El-Taher, A. M., Boghdady, M. S., & Nassar, D. M. A. (2016). The impact of foliar spray with ascorbic acid on growth, productivity, anatomical structure and biochemical constituents of volatile and fixed oils of basil plant (*Ocimum basilicum* L.). *Middle East Journal of Agriculture*, 5(4), 549-565.
- Bahcesular, B., Yildirim, E. D., Karaçocuk, M., Kulak, M., & Karaman, S. (2020). Seed priming with melatonin effects on growth, essential oil compounds, and antioxidant activity of basil (*Ocimum basilicum* L.) under salinity stress. *Industrial Crops and Products*, 146, 1-9.
- Barbalho, S. M., Machado, F. M. V. F., Rodrigues, J. D. S., Silva, T. H. P. D., & Goulart, R. D. A. (2011). Sweet Basil (*Ocimum basilicum*): Much more than a condiment. *Cellmed Orthocellular Medicine and Pharmaceutical Association*, 2(1), 1-5.
- Bourzikat, O., El Abbouchi, A., Ghammaz, H., El Brahmi, N., El Fahime, E., Paris, A., Daniellou, R., Suzenet, F., Guillaumet, G., & El Kazzouli, S. (2022). Synthesis, anticancer activities and molecular docking studies of a novel class of 2-Phenyl- 5,6,7,8- tetrahydroimidazo [1,2-b]pyridazine derivatives bearing sulfonamides. *Molecules (Basel Switzerland)*, 27(16), 5238.
- Burkill, H. (1995). The useful plants of West Tropical Africa. 2nd ed. Kew: Royal Botanic Gardens 3,19-27.
- Chetehouna, S., Atoussi, O., Boulaares, I., Guemari, I. Y., & Derouiche, S. (2020). The effect of Chronic Tobacco smoking on Atherogenic index and Cardiovascular disease risk in El-Oued (Algeria) men. *Asian Journal of Research in Chemistry*, 13(6), 1-6.
- Chetehouna, S., Derouiche, S., & Réggami, Y. (2023). Green synthesis of SeNPs using *Sonchus maritimus* based nanosized metal oxides for in vitro biological applications and in vivo acute toxicity evaluation. *Kragujevac Journal of Science*, (45), 65-78.
- Choi, J. Y., Heo, S., Bae, S., Jiyeon, K., & Moon, K. D. (2019). Discriminating the origin of basil seeds (*Ocimum basilicum* L.) using hyperspectral imaging analysis. *LWT, Food Science and Technology*, 118(5), 108715.
- Chowdhury, T., Mandal, A., Roy, S. C., & De Sarker, D. (2017). Diversity of the genus *Ocimum* (Lamiaceae) through morpho-molecular (RAPD) and chemical (GC-MS) analysis. *Journal of Genetic Engineering and Biotechnology*, 15(1), 275-286.
- Constantinescu, A. M. & Adriana, M. (2019). Antioxidant and antiseptic properties of volatile oils from different medicinal plants: A review. *International Journal of Pharmacognosy and Chinese Medicine*, 3(3).
- Corti, R., Flammer, A. J., Hollenberg, N. K., & Lüscher, T. F. (2009). Cocoa and cardiovascular health. *Circulation*, 119(10), 1433-1441.
- Das Chagas Pereira de Andrade, F., & Mendes, A. N. (2020). Computational analysis of eugenol inhibitory activity in lipoxygenase and cyclooxygenase pathways. *Scientific Reports*, 10(1), 16204.

- De Martino, L., De Feo, V., & Nazzaro, F. (2009). Chemical composition and in vitro antimicrobial and mutagenic activities of seven Lamiaceae essential oils. *Molecules (Basel, Switzerland)*, 14(10), 4213-4230.
- Derouiche, S. (2020). Current review on herbal pharmaceutical improve immune responses against COVID-19 infection. *Research Journal of Pharmaceutical Dosage Forms and Technology*, 12(3), 181-184.
- Derouiche, S., Guemari, I. Y., & Boulaares, I. (2020). Characterization and acute toxicity evaluation of the MgO nanoparticles synthesized from aqueous leaf extract of *Ocimum basilicum* L. *Algerian Journal of Biosciences* 01(01), 001-006.
- Dörr, O. S., Brezina, S., Rauhut, D., & Mibus, H. (2020). Plant architecture and phytochemical composition of basil (*Ocimum basilicum* L.) under the influence of light from microwave plasma and high-pressure sodium lamps. *Journal of Photochemistry and Photobiology. B, Biology*, 202, 111678.
- El-Dakar, A. Y., Shalaby, S. M., Nemetallah, B. R., Saleh, N. E., Sakr, E. M., & Toutou, M. M. (2015). Possibility of using basil (*Ocimum basilicum*) supplementation in Gilthead sea bream (*Sparus aurata*) diet. *The Egyptian Journal of Aquatic Research*, 41(2), 203-210.
- El-Soud, N. H., Deabes, M., El-Kassem, L. A., & Khalil, M. (2015). Chemical Composition and antifungal activity of *ocimum basilicum* L. Essential Oil. *Open Access Macedonian Journal of Medical Sciences*, 3(3), 374-379.
- Eswaramoorthy, R., Hailekiros, H., Kadir, F., & Endale, M. (2021). In silico molecular docking, DFT analysis and ADMET studies of carbazole alkaloid and coumarins from roots of *Clausena anisata*: A potent inhibitor for quorum sensing. *Advances and Applications in Bioinformatics and Chemistry: AABC*, 14, 13-24.
- Falowo, A. B., Mukumbo, F. E., Idamokoro, E. M., Afolayan, A. J., & Muchenje, V. (2019). Phytochemical constituents and antioxidant activity of sweet basil (*Ocimum basilicum* L.) essential oil on ground beef from Boran and Nguni cattle. *International Journal of Food Science*, 2019, 2628747.
- Gebrehiwot H., Dekebo A., & Bachheti R. K. (2016). Characterization of some compounds isolated from Sweet basil (*Ocimum basilicum* L.) leaf extract. *International Journal of Scientific Reports*, 2(7), 159-164.
- Gebrehiwot, H., Bachetti, R. K., & Dekebo, A. (2015). Chemical composition and antimicrobial activities of leaves of sweet basil (*Ocimum basilicum* L.) herb. *International Journal of Basic & Clinical Pharmacology*, 4(5), 869-875.
- Georgiadou, E. C., Kowalska, E., Patla, K., Kulbat, K., Smolińska, B., Leszczyńska, J., & Fotopoulos, V. (2018). Influence of heavy metals (Ni, Cu, and Zn) on nitro-oxidative stress responses, proteome regulation and allergen production in basil (*Ocimum basilicum* L.) plants. *Frontiers in Plant Science*, 9, 862.
- Güez, C. M., Souza, R. O. D., Fischer, P., Leão, M. F. D. M., Duarte, J. A., Boligon, A. A., Athayde, M. L., Zuravski, L., Oliveira, L. F. S. D., & Machado, M. M. (2017). Evaluation of basil extract (*Ocimum basilicum* L.) on oxidative, anti-genotoxic, and anti-inflammatory effects in human leukocyte cell cultures exposed to challenging agents. *Brazilian Journal of Pharmaceutical Sciences*, 53(1), 1-12.
- Gurpinar, E., Grizzle, W. E., & Piazza, G. A. (2013). COX-Independent Mechanisms of Cancer Chemoprevention by Anti-Inflammatory Drugs. *Frontiers in Oncology*, 3, 181.
- Harnafi, H., Aziz, M., & Amrani, S. (2009). Sweet basil (*Ocimum basilicum* L.) improves lipid metabolism in hypercholesterolemic rats. *e-SPEN, the European e-Journal of Clinical Nutrition and Metabolism*, 4(4), e181-e186.
- Idowu, J. A., & Oziegbe, M. (2017). Mitotic and meiotic studies on two species of *Ocimum* (Lamiaceae) and their F1 hybrids. *Botanica Lithuanica*, 23(1), 59-67.
- Kalamartzis, I., Dordas, C., Georgiou, P., & Menexes, G. (2020). The use of appropriate cultivar of basil (*Ocimum basilicum*) can increase water use efficiency under water stress. *Agronomy*, 10(1), 70.
- Katz, D. L., Doughty, K., & Ali, A. (2011). Cocoa and chocolate in human health and disease. *Antioxidants & Redox Signaling*, 15(10), 2779-2811.
- Kiessling, L. L., & Diehl, R. C. (2021). CH-  $\pi$  interactions in glycan recognition. *ACS Chemical Biology*, 16(10), 1884-1893.
- Kostić, M., Popović, Z., Brkić, D., Milanović, S., Sivčev, I., & Stanković, S. (2008). Larvicidal and antifeedant activity of some plant-derived compounds to *Lymantria dispar* L. (Lepidoptera: Limantriidae). *Bioresource Technology*, 99(16), 7897-7901.
- Martinez-Velazquez, M., Castillo-Herrera, G. A., Rosario-Cruz, R., Flores-Fernandez, J. M., Lopez-Ramirez, J., Hernandez-Gutierrez, R., & del Carmen Lugo-Cervantes, E. (2011). Acaricidal effect and chemical composition of essential oils extracted from *Cuminum cyminum*, *Pimenta dioica* and *Ocimum basilicum* against the cattle tick *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae). *Parasitology Research*, 108, 481-487.
- Matasyoh, L. G., Matasyoh, J. C., Wachira, F. N., Kinyua, M. G., Muigai, A. W., & Mukiana, T. K. (2008). Antimicrobial activity of essential oils of *Ocimum gratissimum* L. From different populations of Kenya. *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, 5(2), 187-193.
- Murali, M., & Prabakaran, G. (2018). Effect of Different Solvents System on Antioxidant Activity and Phytochemical Screening in Various Habitats of *Ocimum basilicum* L. (Sweet basil) Leaves. *International Journal of Zoology and Applied Biosciences*, 3, 375-381.
- Nadeem, F., Hanif, M. A., Bhatti, I. A., Jilani, M. I., & Al-Yahyai, R. (2020). Basil. *Medicinal Plants of South Asia*, 47-62.
- Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: An overview. *Journal of Nutritional Science*, 5, e47.
- Purushothaman, B., Srinivasan, R. P., Purushothaman, S., Ranganathan, B., Gimbut, J., & Shanmugam, K. (2018). A comprehensive review on *Ocimum basilicum*. *Journal of Natural Remedies*, 18(3), 2320-3358.

- Qasem, A., Assaggaf, H., Mrabti, H. N., Minshawi, F., Rajab, B. S., Attar, A. A., Alyamani, R. A., Hamed, M., Mrabti, N. N., Baaboua, A. E., et al. (2023). Determination of chemical composition and investigation of biological activities of *Ocimum basilicum* L. *Molecules (Basel, Switzerland)*, 28(2), 614.
- Rameshrad, M., Salehian, R., Fathiazad, F., Hamedeyazdan, S., Garjani, M., Maleki-Dizaji, N., & Vosooghi, R. (2014). The effects of *Ocimum basilicum* ethanol extract on carrageenan induced paw inflammation in rats. *Pharmaceutical Sciences*, 20(4), 149-156.
- Redzicka, A., Wiatrak, B., Jeřkowiak-Kossakowska, I., Kochel, A., Placzek, R., & Czyżnikowska, Ż. (2023). Design, synthesis, biological evaluation, and molecular docking study of 4, 6-Dimethyl-5-aryl/alkyl-2-[2-hydroxy-3-(4-substituted-1-piperazinyl) propyl] pyrrolo [3, 4-c] pyrrole-1, 3 (2 H, 5 H)-diones as Anti-inflammatory agents with dual inhibition of COX and LOX. *Pharmaceuticals*, 16(6), 804.
- Rezzoug, M., Bakchiche, B., Gherib, A., Roberta, A., FlaminiGuido, Kilinçarslan, Ö., Mammadov, R., & Bardaweel, S. K. (2019). Chemical composition and bioactivity of essential oils and Ethanolic extracts of *Ocimum basilicum* L. and *Thymus algeriensis* Boiss. & Reut. from the Algerian Saharan Atlas. *BMC Complementary and Alternative Medicine*, 19(1), 146.
- Rumengan, I. F. M., Mandey, L., Citraningtiyas, G., & Luntungan, A. H. (2019). Antihyperglycemic capacity of basil (*Ocimum basilicum* L.) Leaves extracts coated with the marine fish scales derived nanochitosan. *IOP Conference Series: Materials Science and Engineering*, 56, 1-7.
- Sajjadi, S. E. (2006). Analysis of the essential oils of two cultivated basil (*Ocimum basilicum* L.) from Iran. *DARU Journal of Pharmaceutical Sciences*, 14(3), 128-130.
- Taha, R. S., Alharby, H. F., Bamagoos, A. A., Medani, R. A., & Rady, M. M. (2020). Elevating tolerance of drought stress in *Ocimum basilicum* using pollen grains extract; A natural biostimulant by regulation of plant performance and antioxidant defense system. *South African Journal of Botany*, 128, 42-53.
- Tariq, J., Shumaila, S., Riaz, S., Afzal, S., Hussain, S., Riaz, S., Uzair, M. & Chaudhary, B. (2016). Phytochemical and biological evaluation of *Ocimum basilicum* roots. *International Journals of Pharmacy and Pharmaceutical Research Human*, (2), 292-301.
- Teng, Y. N., Wang, C. C. N., Liao, W. C., Lan, Y. H., & Hung, C. C. (2020). Caffeic acid attenuates multi-drug resistance in cancer cells by inhibiting efflux function of human P-glycoprotein. *Molecules (Basel, Switzerland)*, 25(2), 247.
- Tungmunthum, D., Thongboonyou, A., Pholboon, A., & Yangsabai, A. (2018). Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. *Medicines (Basel, Switzerland)*, 5(3), 93.
- Umoh, R. A., Johnny, I. I., Udoh, A. E., Elijah, A. A., Umoh, O. T., & Essiet, L. E. (2020). Comparative evaluation of the larvicidal properties of methanol extracts and fractions of *Ocimum gratissimum* L. and *Ocimum basilicum* L. leaves (Lamiaceae) on the fourth instar larvae of *Culex quinquefasciatus* L. and control of filariasis. *Journal of Complementary and Alternative Medical Research*, 11(3), 24-31.
- Zarlaha, A., Kourkoumelis, N., Stanojkovic, T. P., & Kovala-Demertzi, D. (2014). Cytotoxic activity of essential oil and extracts of *Ocimum basilicum* against humancarcinoma cells. Molecular docking study of isoeugenol as a potent cox and loxinhibitor. *Digest Journal of Nanomaterials and Biostructures*, 9(3), 907-917.