

Effect of Fasciola Hepatica Invasion on Cow Productivity

Igor Gennadievich Glamazdin, Ilya Nikolaevich Medvedev*, Ilnura Ibragimovna Fayzullina, Natalya Yuryevna Sysoeva, Marina Mikhailovna Goryacheva, Galina Mikhailovna Kryukovskaya, Tatyana Olegovna Maryushina

Received: 12 June 2022 / Received in revised form: 03 September 2022, Accepted: 07 September 2022, Published online: 12 September 2022
© Biochemical Technology Society 2014-2022
© Sevas Educational Society 2008

Abstract

In the study, the level of antibodies to fasciolas in the milk of cows in the Moscow region of Russia was assessed and the relationship between their number and the productive characteristics of these animals was traced. In October 2021, a milk sample was taken from a dairy farm in the Moscow region of Russia, and the number of antibodies to Fasciola hepatica (ODRf) and Ostertagia ostertagi (ODRo) was assessed. Their level was determined using multivariate linear regression models and their relationship with economically significant indicators was traced: the amount of protein in milk, the percentage of fat in milk, and the interval between past calving. An increase in the ODRf quartile from 25% (0.412) to 75% (0.976) was associated with a decrease in the volume of milk produced per year by 0.8 kg (day of the lactation process) ($p=0.002$), with a decrease in fat content in milk by 0.07% ($p<0.001$), with an increase in the interval between sections by 4.6 days ($p=0.04$) in the absence of a significant relationship with the level of protein in milk. In the case of simultaneous infection with both parasites, the effect of ODRf and ODRo on the volume of milk produced was additional, rather than synergistic.

Keywords: Cows, Parasitism, Fasciola hepatica, Ostertagia ostertagi, Milk production, ELISA

Introduction

Modern science continues to search for approaches to improve health (Karpov *et al.*, 2021; Zavalishina *et al.*, 2022) and increase the viability of the mammalian organism (Zavalishina, 2018b; Kočović *et al.*, 2021). For this purpose, direct observations are carried out (Zavalishina, 2018c; Zavalishina *et al.*, 2021), and various schemes of experimental work are set up (Vorobyeva *et al.*, 2018; Kulikov *et al.*, 2020). At the same time, despite all the

Igor Gennadievich Glamazdin, Ilnura Ibragimovna Fayzullina, Natalya Yuryevna Sysoeva, Marina Mikhailovna Goryacheva, Galina Mikhailovna Kryukovskaya, Tatyana Olegovna Maryushina

Department of Veterinary Medicine, Moscow State University of Food Production, 125080, Moscow, Russia.

Ilya Nikolaevich Medvedev*
Faculty of Medicine, Russian State Social University, 129226, Moscow, Russia.

*E-mail: ilmedv1@yandex.ru

efforts made by science and practice, the Fasciola hepatica parasite is still common and very common in cattle. The situation is complicated by the fact that, as a rule, fascioliasis in cattle is mostly asymptomatic (Zavalishina, 2018d; Yıldız *et al.*, 2021), but it causes serious physiological changes that lead to significant economic losses (Torgerson & Claxton, 1999). Despite this fact, few studies related to the impact of the disease on productivity levels have been performed on this parasitic disease (Vercruysse & Claerebout, 2001; Drljača *et al.*, 2020). At the same time, in some cases, there were questions about the scheme or design of the study (Burden *et al.*, 1978; Dargie, 1987).

The presence of *F. hepatica* in the body, as a rule, is determined using scatological methods. At the same time, in recent years, specific tests have been increasingly used to determine its presence, which is gradually replacing scatological research methods. Newer tests are associated with the detection of specific fasciola antigens in blood and feces (Leclipteux *et al.*, 1998; Mezo *et al.*, 2004) or the detection of the presence of specific antibodies to fasciola in the blood (Farrell *et al.*, 1981; Salimi-Bejestani *et al.*, 2005b; Korobov & Glamazdin, 2010) and in milk (Boulard *et al.*, 1985; Reichel *et al.*, 2005; Salimi-Bejestani *et al.*, 2005a). The available reagent kits for antibodies to fasciola have already been sufficiently developed and allow detecting the presence of the parasite with an unmistakable accuracy, which makes it possible to judge the degree of its spread. Given the impossibility of the complete eradication of *F. hepatica*, it is important to direct diagnostic control to reduce its frequency of occurrence and prevalence, in which the invasion would have a minimal impact on the overall lactation capacity of the dairy herd of cows (Vercruysse & Claerebout, 2001; Mai *et al.*, 2022).

Previously, the prevalence of invasion in the amount of 30 different flukes with a prevalence in the herd of 25% of the total population of the herd was supposed to be considered as threshold values for optimum productivity (Vercruysse & Claerebout, 2001; Roest Henk *et al.*, 2021). However, these figures either cannot be determined prior to slaughter or are based on questionable information. Recording the amount of highly specific antibodies to the parasite present in marketed milk is now considered a valid method of tracking the overall level of parasite infestation in a dairy herd. Also, this method is recognized as reliable for the diagnosis of parasites in the herd, where the level of infestation of cows causes a decrease in productivity (Sanchez & Dohoo, 2002; Charlier *et al.*, 2005a). Scientists have elucidated the relationship between the content of antibodies specific to fasciola in milk and



the number of antibodies in the blood of lactating cows (Salimi-Bejestani *et al.*, 2005a; Sloan *et al.*, 2021).

The purpose of the study: is to find out the relationship between the number of antibodies specific to fasciola in milk and indicators reflecting the productivity of dairy cows in the Moscow region of Russia to assess possible economic losses.

Materials and Methods

The study was conducted in dairy farms of the Moscow region of Russia with a total number of livestock of more than 2000 dairy cows. Milk samples were taken from the milk collection containers. All taken milk samples were delivered to the laboratory of the Department of Veterinary Medicine of the Moscow State University of Food Production in Moscow within 6 hours after milking. After delivery to the laboratory, all milk samples were subjected to centrifugation (16000 g for 5 minutes). Fat was removed from them, the supernatant was obtained and frozen at -20°C until the scheduled analyzes were performed. All selected milk samples were tested using *F. hepatica* ELISA and *O. ostertagi* ELISA.

Cows that gave milk, samples of which were taken for analysis, were kept in farms in the Moscow region of Russia. They were purebred for the Black-and-White breed or had different bloodlines for the Black-and-White and Holstein breeds. In general, the cows that gave milk for sampling had 2-4 lactations. The study was completed in March 2021.

After defrosting all milk samples, they were examined using 2 indirect ELISAs to detect the content of specific antibodies capable of reacting with *Fasciola hepatica* and *Ostertagia ostertagi* antigens.

The amount of antibodies capable of reacting with *F. hepatica* was recorded by the traditional method by Salimi Bejestani *et al.* (2005b). For this, 96-well microplates with a flat bottom were used. They were loaded per well with 0.5 mg/ml excretory and secretory (ES) antigen in 0.05 M carbonate-bicarbonate buffer (pH 9.6). An overnight incubation followed by washing the well six times with phosphate-buffered saline containing 0.05% tween 80 (PBST). Then blocking was done with 100 ml of 2% skimmed milk powder (Marvel, Premier Beverages, Stafford, UK) in PBST (SMP/PBST) and the incubation process was carried out for 1 hour at 37.0°C. The plates used in the work were traditionally washed and undiluted samples of the studied milk (100 ml) were added to their wells. Evaluated serum samples, considered as negative and positive controls, were diluted at 1:800 in SMP/PBST and added (100 ml/well) to six wells of each plate. Tablets are traditionally incubated and washed.

Rabbit anti-bovine IgG conjugated with horseradish peroxidase (Jackson Immunoresearch Laboratories) was used. It was diluted 1/6000 in SMP/PBST and added to wells per 100 ml. The substrate solution consisted of 0.1% OPD (orthophenylenediamine) in citric acid/phosphate buffer (pH 5.0) with 1/2000 H₂O₂ and was applied to wells at a rate of 100 ml. The plates thus prepared were incubated at room temperature in the absence of light for 10 min. Blocking of the reaction was carried out by introducing 50 ml of

2.5 M HCl and then the optical density was evaluated at a wavelength of 492 nm. The amount of antibodies was expressed as the ratio of the detected optical density (ODRf) according to the formula $ODRf = (OD\ NC)/(PC\ NC)$ where OD is the value of the optical density of the assessed image, and NC and PC OD are negative and positive levels. The sensitivity of this enzyme immunoassay makes it possible to detect infection in milk in a herd of more than 25% of cows with an accuracy of 95% (95% CI 89–100%) and 80% (95% CI 66–94%), respectively.

It has been noted that the ES antigen is unable to bind to *O. ostertagi* (Salimi-Bejestani *et al.*, 2005a; Salimi-Bejestani *et al.*, 2005b). Anti-*O. ostertagi* antibodies were quantified using a known ELISA method (Charlier *et al.*, 2005b). The ELISA used crude antigen from mature *O. ostertagi*. The result of this test is reported as an optical density ratio (ODRo). The expected 95% scatter difference between two ODRo values of a particular sample using separate plates and on different days reaches 0.16 (Charlier *et al.*, 2005b) The adult worm crude antigen interacts very well with other infections of the gastrointestinal tract, including *Cooperia* spp. (Keus *et al.*, 1981) and with other worms, including *F. hepatica* (Eysker & Ploeger, 2000).

Differences in ODR values were noted between dairy farms for which production situation data were and were not obtained by applying a two-tailed Mann-Whitney U-test ($\alpha = 0.05$) for ODRf and a two-tailed t-test ($\alpha = 0.05$) for ODRo. The level of Spearman's rank correlation between ODRf and ODRo was determined.

The initial relationship of ODRf with the number of economically significant indicators (the amount of milk produced, the percentage of protein in milk, the fat content of milk, and the time between the onset of calving) was estimated using linear regression models. The dairy herd was considered as a whole, and the significance of the parameters was assessed using two-tailed F-tests ($\alpha=0.05$). The average indicators of the amount of milk received, the protein content in milk, and the level of the fat content of milk over the past year before taking milk samples for the study were taken into account and their relationship with ODRf was traced. Several covariates were introduced into the model as independent variables because they were economically important factors: mean number of lactating cows, mean number of lactations, mean duration of lactation, mean logarithm [(number of somatic cells/1000)/mL], and ratio miscarriages to the number of calving for the whole year. Since a negative correlation is known between the value of the average annual milk yield and % of its fat content (apparently due to the effect of dilution of milk), milk yield was considered as a covariant in the course of studying the relationship of ODRf with % of milk fat. The effect of simultaneous parasitism in the body of the cow *F. hepatica* and gastrointestinal nematodes on the amount of milk received was considered an independent factor in the model that evaluates the relationship between ODRf and the volume of milk received and reveals the relationship between ODRf and ODRo. The relationship between the number of ODRf and the time between calving was determined using a model that included covariates in the form of the distribution of calving, the bulk of cows in the herd, the average annual number of productive animals in the herd, and the number of lactations in cows. The

adequacy of the models was determined by evaluating the quadratic terms for any independent variable. The normality of the individual considered models was determined using standard plots of residuals and plots of probability compared to the predicted values.

Results and Discussion

The results of the assessment of antibody levels in relation to ODRf and ODRo are shown in **Table 1**. The values of the ODRf content

were shifted to the right side, while the distribution of ODRo values showed a normal character (**Figure 1**). The numbers of ODRf and ODRo were lower in holdings reporting performance data than in herds for which such information was available. The differences found were not high, but statistically significant (ODRf: $p < 0.001$; ODRo: $p = 0.003$). A positive correlation was found between the level of ODRf and the amount of ODRo ($R = 0.35$; $p < 0.001$).

Table 1. General information on the levels of antibodies to *F. hepatica* (ODRf) and *O. ostertagi* (ODRo) in milk samples taken from tanks, in October 2021 in the Moscow region of Russia

Characteristics	Minimum	Quantile value	Maximum	25%	50%	75%
ODRf						
Information received from the farm	52	0.248	0.386	0.510	0.931	1.756
Farm data not received	36	0.165	0.437	0.632	1.012	2.117
General level	88	0.165	0.412	0.559	0.976	2.117
ODRo						
Production information received	54	0.063	0.305	0.418	0.508	0.962
Production data not received	31	0.085	0.327	0.436	0.542	1.157
General level	85	0.085	0.315	0.422	0.530	1.157




Figure 1. Frequency distribution of levels of antibodies to *F. hepatica* (ODRf) and to *O. ostertagi* (ODRo) in milk from reservoirs in the Moscow Region collected in March 2021

The overall performance statistics collected in the year prior to sampling are summarized in **Table 2**.

Table 2. Performance characteristics at the herd level from April 2020 to March 2021

Index	Minimum	quartiles			Maximum
		25%	50%	75%	
The volume of milk received, kg / cow day					
Median ODRf	11.2	21.3	23.6	25.4	32.8
Median ODRf	12.7	18.5	21.1	23.6	30.5

Lactation days	72	183	194	205	293
Median ODRf					
Median ODRf	155	183	195	210	480
Milk fat %	3.4	4.2	4.4	4.6	5.2
Median ODRf					
Median ODRf	3.4	4.2	4.4	4.5	5.0
Protein content in milk, %	3.0	3.2	3.3	3.4	3.7
Median ODRf					
Median ODRf	3.0	3.2	3.3	3.4	3.5
Number of lactation	1.4	2.3	2.5	2.7	4.1
Median ODRf					
Median ODRf	1.9	2.3	2.5	2.8	3.9
Number of breeding animals	4	31	40	51	123
Median ODRf					
Median ODRf	15	28	37	49	127
The ratio of miscarriages and annual calving	0.08	0.71	0.78	0.84	0.99
Median ODRf					
Median ODRf	0.01	0.60	0.76	0.82	0.91

A significant negative relationship was noted between the value of ODRf and the volume of average annual milk produced (Table 3). The increase in ODRf within the interquartile interval was associated with a decrease in the volume of milk supplied during the year by 0.8 kg/(cow day) (p=0.002), which is associated with a decrease in the volume of milk supplied per year by 3%.

It was not possible to find a statistically significant relationship between the level of ODRf and the percentage of protein in milk [slope = 0.013; 95% CI = (0.035; 0.007); p = 0.24]. An increase in ODRf within the interquartile range was associated with a 0.07% drop in milk fat [slope = 0.095; 95% CI = (0.145; 0.046); p<0.001],

and when the range between calvings increases by 4.6 days [slope = 7.6; 95% CI = (0.78; 14.3); p=0.04].

When looking for a relationship between ODRf and ODRo on the one hand, and with milk yield on the other hand, the negative relationship for ODRo [slope index = 4.47; 95% CI = (6.41; 2.61); p<0.001]. An unexpressed negative relationship was found for the ODRf level [slope index = 0.64; 95% CI = (1.32; 0.09), p=0.07]. At the same time, there was no significant relationship between the levels of ODRf and ODRo in relation to their influence on the level of milk yield (p=0.69).

Table 3. Linear regression model data to find the relationship between ODRf in milk and average annual milk yield

Variable	δ	95% CI		n- meaning
		LL	UL	
Meaning ODRf	1.11	1.80	0.40	0.003
Average number of cows	0.025	0.007	0.010	0.002
Average number of lactated cows	5.5	0.5	10.6	0.028
Average number of lactations squared	1.2	1.9	0.3	0.018
Average duration of lactation	0.014	0.025	0.005	0.009

The existence of an organism is always associated with many biological interactions (Zavalishina, 2018a). This largely determines the success of the development of the entire ontogeny and the adaptation of the organism to the external environment (Zavalishina, 2020b). A very common interaction in wildlife is parasitism. It has a varying degree of impact on the animal's body, and many of its aspects still need to be clarified.

The milk samples analyzed in the study were taken in October 2021. The activity of the intermediate hosts of fasciola begins in

April-May. It is known that antibodies against F. hepatica ES antigens are registered 2-4 weeks after the parasite enters the body (Salimi-Bejestani *et al.*, 2005a) and can remain for a long time after its complete removal from the body (Boulard *et al.*, 1995; Chauvin *et al.*, 1997). In view of the fact that the presence of antibodies to F. hepatica in body fluids is a consequence of past interactions between the parasite and the organism, it has been suggested that a negative impact on productivity is associated with migratory young flukes (Zavalishina, 2020a). This occurs, as a rule, during

the stay of animals on the pasture, and by adult flukes, as a rule, during the winter maintenance. In the work, the available production characteristics of the year preceding the collection of milk samples were used, which made it possible to evaluate the relationship between ODRf and productivity.

The presence of a somewhat distorted distribution of the ODRf level and an approximately normal distribution of the ODRo level was quite expected, in view of the fact that the ODRf distribution includes non-infected and infected groups of animals, while for *O. ostertagi* the entire population should be considered as infected with it (Agneesssens *et al.*, 2000).

ELISA data and farm worker data cannot be obtained from all randomly selected herds. At the same time, the differences in ODR levels between herds, in which the level of milk yield was taken into account or not, turned out to be fully statistically significant. In this regard, the conclusions obtained using models that traced the relationship between production parameters and parasitism cannot be considered representative of the entire sample of animals considered. Although several possible confounding factors have been considered, it should be recognized that the associations found may be partly due to the influence of other factors that can reduce the amount of milk produced, but are caused by parasitism in the body of *F. hepatica*. At the same time, the conducted study contains an assessment of economic losses traced in many livestock farms, and the patterns found for the simultaneous presence of several parasites will be true for various dairy herds kept in areas endemic to *F. hepatica*. Also, the obtained information can be applied in studies, when it turns out the cost or economic efficiency of individual measures to control *F. hepatica*. ELISA on *F. hepatica* is strictly species-specific, while ELISA on *O. ostertagi* can also bind to *F. hepatica*. In this regard, the impact of ODRf on productivity levels was assessed, without considering ODRo in the course of the study. ODRf and ODRo were used simultaneously as part of the model to determine the possibility of a synergistic effect of the presence of concomitant parasites.

There is concern that the presence of *F. hepatica* in the body of cows reduces the overall level of nutrients in milk (Black & Froyd, 1972). The results obtained in the work show that infection of the body of a cow with liver fluke does not affect the amount of protein present in milk, contributing to a decrease in the concentration of fat molecules in milk. This is in contrast to nematode infestations, which have little effect on milk fat content and milk protein levels (McPherson *et al.*, 2001; Nødvedt *et al.*, 2002; Charlier *et al.*, 2005a).

It was previously noted that the presence of liver fluke in the body of a cow reduces the likelihood of conception and complicates the course of pregnancy (Dargie, 1987). The data obtained by the authors confirm this in view of the fact that farms with a low level of ODRf in cows showed the greatest intervals between calving. In this regard, in the course of assessing the damage from the invasion of *F. hepatica*, it is rational to take into account the upcoming reproductive pathology.

It was previously noted that the simultaneous invasion of *F. hepatica* and *O. Ostertagi* leads to more negative changes than either of them separately. The work performed by the authors

allows us to assume the presence of a total, rather than mutually aggravating, the influence of each parasite. This is quite consistent with the data (Dargie, 1987), which did not reveal a relationship between *Ostertagia* and *Fasciola* invasions concerning the general condition of the animals or concerning changes in the body weight of the young born.

The negative relationship between the number of ODRf and economically important characteristics of animals makes it possible to determine the "economic threshold" that allows identifying herds with liver fluke invasion, which affects the level of milk secretion. Estimation of the level of antibodies at its cost is quite affordable and should be done to record their number, capable of predicting a positive response to the productivity of animals from interventions against *Fasciola* infestations. More observations are required to elucidate the dynamics of the level of productivity with the use of fasciolicide among dairy cows with different levels of ODRf and to determine the relationship between the number of ODRf and productive capacity in some dairy cows.

Conclusion

Invasion of the parasite *Fasciola hepatica* is still common in cattle throughout the world. Even though fasciolosis is often asymptomatic in cattle, it causes a significant decrease in the productivity of cows. In this regard, the broad diagnostics of this parasite in the livestock of cows is very relevant. The presence of *F. hepatica* in the body is traditionally determined using scatological methods. Nowadays, highly sensitive immunological tests are increasingly being used to detect it. Tests for antibodies to *fasciola* are well developed and can accurately detect the presence of the parasite. In modern farms, it is important to monitor the frequency of occurrence of this parasite, given that with its prevalence, a minimal decrease in the productivity of a herd of dairy cows up to 25% of their livestock is possible. In the work carried out in October 2021, the level of antibodies to *fasciolas Fasciola hepatica* (ODRf) and *Ostertagia ostertagi* (ODRo), which are in the composition of milk, which was obtained from healthy cows in the Moscow region of Russia, was assessed and a relationship was traced with the productive characteristics of dairy cows. The number of antibodies was estimated using multivariate linear regression models, tracing their relationship with protein content in milk, percentage of fat in milk, and intervals between births in cows. An increase in the ODRf level from the 25% quantile (0.412) to the 75% quantile (0.976) in the Moscow region of Russia was associated with a decrease in the amount of milk supplied during the year by 0.8 kg (lactation day) ($p=0.002$), with a decrease in the value the average amount of fat in the composition of milk by 0.07% ($p<0.001$), with an increase in the intervals between sections by 4.6 days ($p=0.04$) and in the absence of a significant relationship with the level of protein in milk. It turned out that the relationship between ODRf and ODRo was complementary rather than synergistic in terms of co-infection.

Acknowledgments: The team of authors thanks the administration of the Moscow State University of Food Production for the opportunity to research its basis.

Conflict of interest: None

Financial support: None

Ethics statement: The study was approved by the local ethics committee of the Moscow State University of Food Production on September 15, 2018 (protocol №11).

References

- Agneessens, J., Claerebout, E., Dorny, P., Borgsteede, F. H., & Vercruyse, J. (2000). Nematode parasitism in adult dairy cows in Belgium. *Veterinary Parasitology*, *90*(1-2), 83-92.
- Black, N. M., & Froyd, G. (1972). The possible influence of liver fluke infestation on milk quality. *Veterinary Record*, *90*(3), 71-72.
- Boulard, C., Carreras, F., & Van Gool, F. (1995). Evaluation of nitroxylin and closantel activity using ELISA and egg counts against *Fasciola hepatica* in experimentally and naturally infected cattle. *Veterinary Research*, *26*(4), 249-256.
- Boulard, C., Bouvry, M., Argente, G., Villejoubert, C., Galliene, C., & Lévêque, F. (1985). Comparaison de la détection des foyers de fasciolose par test ELISA sur lactosérum et sérum et par coproscopie. In *Annales de Recherches Vétérinaires* (Vol. 16, No. 4, pp. 363-368).
- Burden, D. J., Hughes, D. L., Hammet, N. C., & Collis, K. A. (1978). Concurrent daily infection of calves with *Fasciola hepatica* and *Ostertagia ostertagi*. *Research in Veterinary Science*, *25*(3), 302-306.
- Charlier, J., Claerebout, E., Duchateau, L., & Vercruyse, J. (2005a). A survey to determine relationships between bulk tank milk antibodies against *Ostertagia ostertagi* and milk production parameters. *Veterinary Parasitology*, *129*(1-2), 67-75.
- Charlier, J., Duchateau, L., Claerebout, E., & Vercruyse, J. (2005b). Assessment of the repeatability of a milk *Ostertagia ostertagi* ELISA and effects of sample preparation. *Preventive Veterinary Medicine*, *68*(2-4), 277-288.
- Chauvin, A., Moreau, E., & Boulard, C. (1997). Diagnosis of bovine fascioliasis using serology of pools of sera. Interpretation in field conditions. *Veterinary Research*, *28*(1), 37-43.
- Dargie, J. D. (1987). The impact on production and mechanisms of pathogenesis of trematode infections in cattle and sheep. *International Journal for Parasitology*, *17*(2), 453-463.
- Driljača, J. N., Vejnović, A. M. T., Miljković, D. M., Popović, M. J., Rakić, D. B., Sekulić, S. R., Čapo, I. Đ., & Petković, B. B. (2020). Changes in mouse thymus after exposure to tube-restraint stress. *Archives of Biological Sciences*, *72*(1), 5-11.
- Eysker, M., & Ploeger, H. W. (2000). Value of present diagnostic methods for gastrointestinal nematode infections in ruminants. *Parasitology*, *120*(7), 109-119.
- Farrell, C. J., Shen, D. T., Wescott, R. B., & Lang, B. Z. (1981). An enzyme-linked immunosorbent assay for diagnosis of *Fasciola hepatica* infection in cattle. *American Journal of Veterinary Research*, *42*(2), 237-240.
- Karpov, V. Y., Zavalishina, S. Y., Bakulina, E. D., Dorontsev, A. V., Gusev, A. V., Fedorova, T. Y., & Okolelova, V. A. (2021). The physiological response of the body to low temperatures. *Journal of Biochemical Technology*, *12*(1), 27-31.
- Keus, A., Kloosterman, A., & Van den Brink, R. (1981). Detection of antibodies to *Cooperia* ssp. and *Ostertagia* ssp. in calves with the enzyme linked immunosorbent assay (ELISA). *Veterinary Parasitology*, *8*(3), 229-236.
- Kočović, D. M., Bajuk-Bogdanović, D., Maslovarić, I., Božić-Nedeljković, B., Andus, P. R., & Daković, M. (2021). Raman spectral analysis of the brainstem and responses of neuroglia and cytokines in whole-body gamma-irradiated rats after administration of aminothioliol-based radioprotector GL2011. *Archives of Biological Sciences*, *73*(2), 161-173.
- Korobov, A. I., & Glamazdin, I. G. (2010). Elaboration of diagnostic test-system on the basis of dot enzyme-linked immunosorbent assay (dot-elisa) at cattle fasciolosis. *Russian Parasitological Journal*, *3*, 88-92.
- Kulikov, E. V., Zavalishina, S. Y., Vatikov, Y. A., Seleznev, S. B., Parshina, V. I., Voronina, Y. Y., Popova, I. A., Bondareva, I. V., Petrukhina, O. A., Troshina, N. I., et al. (2020). The effects of meldonium on microrheological abnormalities of erythrocytes in rats with obesity: An experimental study. *Bali Medical Journal*, *9*(2), 444-450.
- Leclipteux, T., Torgerson, P. R., Doherty, M. L., McCole, D., Protz, M., Farnir, F., & Losson, B. (1998). Use of excretory/secretory antigens in a competition test to follow the kinetics of infection by *Fasciola hepatica* in cattle. *Veterinary Parasitology*, *77*(2-3), 103-114.
- Mai, J., Wang, K., Liu, C., Xiong, S., & Xie, Q. (2022). $\alpha\beta 3$ -targeted sEVs for efficient intracellular delivery of proteins using MFG-E8. *BMC Biotechnology*, *22*(1), 1-10. doi:10.1186/s12896-022-00745-7
- McPherson, W. B., Gogolewski, R. P., Slacek, B., Familton, A. S., Gross, S. J., Maciel, A. E., & Ryan, W. G. (2001). Effect of a peri-parturient eprinomectin treatment of dairy cows on milk production. *New Zealand Veterinary Journal*, *49*(3), 106-110.
- Mezo, M., González-Warleta, M., Carro, C., & Ubeira, F. M. (2004). An ultrasensitive capture ELISA for detection of *Fasciola hepatica* coproantigens in sheep and cattle using a new monoclonal antibody (MM3). *Journal of Parasitology*, *90*(4), 845-852.
- Nødtvedt, A., Dohoo, I., Sanchez, J., Conboy, G., DesCôteaux, L., & Keefe, G. (2002). Increase in milk yield following eprinomectin treatment at calving in pastured dairy cattle. *Veterinary Parasitology*, *105*(3), 191-206.
- Reichel, M. P., Vanhoff, K., & Baxter, B. (2005). Performance characteristics of an enzyme-linked immunosorbent assay performed in milk for the detection of liver fluke (*Fasciola hepatica*) infection in cattle. *Veterinary Parasitology*, *129*(1-2), 61-66.
- Roest, H. P., IJzermans, J. N., & van der Laan, L. J. (2021). Evaluation of RNA isolation methods for microRNA quantification in a range of clinical biofluids. *BMC Biotechnology*, *21*(1), 1-11. doi:10.1186/s12896-021-00706-6
- Salimi-Bejestani, M. R., Daniel, R. G., Felstead, S. M., Cripps, P. J., Mahmood, H., & Williams, D. J. (2005a). Prevalence of

- Fasciola hepatica in dairy herds in England and Wales measured with an ELISA applied to bulk-tank milk. *Veterinary Record*, 156(23), 729-731.
- Salimi-Bejestani, M. R., McGarry, J. W., Felstead, S., Ortiz, P., Akca, A., & Williams, D. J. (2005b). Development of an antibody-detection ELISA for Fasciola hepatica and its evaluation against a commercially available test. *Research in Veterinary Science*, 78(2), 177-181.
- Sanchez, J., & Dohoo, I. (2002). A bulk tank milk survey of Ostertagia ostertagi antibodies in dairy herds in Prince Edward Island and their relationship with herd management factors and milk yield. *The Canadian Veterinary Journal*, 43(6), 454.
- Sloan, S., Jenvey, C. J., Piedrafita, D., Preston, S., & Stear, M. J. (2021). Comparative evaluation of different molecular methods for DNA extraction from individual Teladorsagia circumcincta nematodes. *BMC Biotechnology*, 21(1), 1-13. doi:10.1186/s12896-021-00695-6
- Torgerson, P., & Claxton, J. (1999). Epidemiology and Control. In: Dalton, J. P. (Ed.). *Fasciolosis*. CABI Publishing, Wallingford, 113-149.
- Vercruysse, J., & Claerebout, E. (2001). Treatment vs non-treatment of helminth infections in cattle: defining the threshold. *Veterinary Parasitology*, 98(1-3), 195-214.
- Vorobyeva, N. V., Glagoleva, T. I., Mal, G. S., Zavalishina, S. Y., & Fayzullina, I. I. (2018). Influence of physical exercise on the activity of brain processes. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(6), 240-244.
- Yıldız, B., Kamiloğlu, N. N., Öziç, C., & Coşkun, Y. (2021). Effects of hypoxia on the mRNA expressions of TRAIL-mediated cell death related genes in hypoxia-tolerant rodent (Nannospalax nehringi) and some characteristics of these proteins. *Archives of Biological Sciences*, 73(1), 5-15.
- Zavalishina, S. Y. (2018a). Functional properties of anticoagulant and fibrinolytic activity of blood plasma in calves in the phase of milk nutrition. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(5), 659-664.
- Zavalishina, S. Y. (2018b). Physiological Dynamics of the Blood Coagulation System Activity in Calves During the Phase of Dairy Nutrition. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(5), 680-685.
- Zavalishina, S. Y. (2018c). Functional activity of the blood clotting system in calves during the phase of milk and vegetable nutrition. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(5), 720-725.
- Zavalishina, S. Y. (2018d). Anti-coagulant and fibrinolytic activity of blood plasma in healthy calves of dairy-vegetative nutrition. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9(5), 753-758.
- Zavalishina, S. Y. (2020a). Functional Activity of the Cardiorespiratory System and the General Level of Physical Capabilities Against the Background of Regular Physical Exertion. *Bioscience Biotechnology Research Communications*, 13(4), 2327-2331. doi:10.21786/bbrc/13.4/105
- Zavalishina, S. Y. (2020b). Functional Features of Hemostasis in Weakened Newborn Calves Treated with Aminosol. *Bioscience Biotechnology Research Communications*, 13(3), 1251-1256.
- Zavalishina, S. Y., Bakulina, E. D., Eremin, M. V., Kumantsova, E. S., Dorontsev, A. V., & Petina, E. S. (2021). Functional changes in the human body in the model of acute respiratory infection. *Journal of Biochemical Technology*, 12(1), 22-26.
- Zavalishina, S. Y., Shalupin, V. I., Rodionova, I. A., Kumantsova, E. S., Rysakova, O. G., Ryazantsev, A. A., & Sibgatulina, F. R. (2022). Influence of Regular Basketball Practice in Adolescence on the Functional Capacity of the Heart. *Journal of Biochemical Technology*, 13(1), 20-24.