

Amycolatopsis alba var. nov DVR D4, a bioactive actinomycete isolated from Indian marine environment

Venkata Ratna Ravi Kumar Dasari*, Sri Rami Reddy Donthireddy

Received: 09 September 2010 / Received in revised form: 2 November 2011, Accepted: 5 November 2011, Published online: 25 December 2011, © Sevas Educational Society 2008-2011

Abstract

The taxonomic position of a bioactive marine isolate, strain DVR D4 was established using a polyphasic approach. The organism merits species status in the genus *Amycolatopsis* according to the chemical and phenotypic data. Phylogenetic analysis of the strain based on its 16S rDNA sequence shows that there was 100% pairwise similarity and identity with no nucleotide gaps with the species *Amycolatopsis alba* strain DSM 44262. As the organism was distinguished with substantial differences in some of the phenotypic characteristics and other properties, it was proposed as a strain variety of *Amycolatopsis alba* and designated as *Amycolatopsis alba* var. nov DVR D4 (GenBank accession number of strain D4 is JN872327).

Key words: *Amycolatopsis alba*, polyphasic taxonomy, phylogenetic analysis, 16S rDNA sequence, bioactive, antagonistic activity.

Introduction

The genus *Amycolatopsis*, assigned to the family *Pseudonocardiaceae* (Warwick et al. 1994; Embley et al. 1988) was proposed by Lechevalier et al. (1986) to accommodate nocardioform actinomycetes having type IV cell composition, type PII phospholipid pattern and lacking mycolic acid. The members of the genus *Amycolatopsis* are Gram-positive, non-acid fast, non-motile actinomycetes that form branched vegetative hyphae that undergo fragmentation into rod-like and squarish elements. The members of this genus have high G+C (>55%) content in the DNA (Cwale et al. 2011).

Venkata Ratna Ravi Kumar Dasari*, Sri Rami Reddy Donthireddy

Centre for Biotechnology, College of Engineering, Andhra University, Visakhapatnam, Andhra Pradesh, INDIA-530 003

Tel: 00919440649700
Email: ravidvr@gmail.com

Only 10 novel species have been described for this genus until the last decade, But since 2000, many novel species have been described which were isolated from various terrestrial environments (Carlson et al. 2007; Groth et al. 2007; Lee et al. 2006; Tan et al. 2006; Saintpierre-Bonaccio et al. 2005; Kim et al. 2002; Huang et al. 2001; Goodfellow et al. 2001) and clinical material (Huang et al. 2004; Labeda et al. 2003). At the time of writing, the genus contains 48 species (Labeda et al. 2011; Albarracin et al. 2010; Atchareeya et al. 2010; Chen et al. 2010; Duangmal et al. 2010; Tamura et al. 2010; Tang et al. 2010; Liras & Demain 2009; Soon 2006).

Literature suggests that there have been some important novel bioactive compounds isolated from the members of the genus *Amycolatopsis* (Demain & Zhang 2005; Zhang et al. 2005; Trenin et al. 2001; Krishna et al. 1998; Malkova et al. 1991). In an effort to explore the relatively untapped potential members of this genus and investigate the potential application of their secondary metabolites, we attempted to isolate and identify strains having the capability of producing bioactive compounds from the marine sediments of coastal Visakhapatnam, India.

Materials and Methods

Isolation

Sediments from Bay of Bengal (NTPC area, Visakhapatnam, India) were collected by grab sampler. Strain DVR D4 was isolated on a starch casein agar (SCA- soluble starch, 10.0g; vitamin free casein, 0.3g; KNO₃, 2.0g; NaCl, 2.0g; K₂HPO₄, 2.0g; MgSO₄.7H₂O, 0.05g; CaCO₃, 0.02g; FeSO₄.7H₂O, 0.01g; agar, 20.0g; sterilized natural aged sea water, 1L; pH, 7.2; supplemented with rifampicin 2.5µg/ml and cycloheximide 75µg/ml to inhibit bacterial and fungal contamination, respectively) plate, which had been seeded with a sediment sample suspension and incubated at 28°C for 14 days (Ramesh & Narayanasamy 2009). The isolate was maintained on YEME (Yeast Extract Malt Extract) slants at 4°C and as a glycerol suspension (20%v/v) at -20°C (Williams & Cross 1971).

Taxonomy

The cultural and morphology properties were examined by using standard procedures (Holt et al. 1994) and procedures of Williams et




Figure 1: Neighbor-joining tree (Saitou and Nei, 1987) based on almost complete 16S rDNA sequences showing relationships between strain DVR D4 and representatives of the family *Pseudonocardiaceae* and related taxa.

Table 1: 16S rDNA similarity values between strain DVR D4 and the representatives of the genus *Amycolatopsis*

Rank	Name	Strain	Accession	Pair-wise Similarity	Identities	Diff/Total nucleotides
1	<i>Amycolatopsis alba</i>	DSM 44262	NR_024888.1	100%	1428/1428	0/1428
2	<i>Amycolatopsis coloradensis</i>	DSM 44225T	AJ421142.1	99%	1421/1431	4/1431
3	<i>Amycolatopsis umgeniensis</i>	UM16	DQ110876.1	99%	1412/1423	3/1423
4	<i>Amycolatopsis</i> sp. 232068	-	GU130127.1	99%	1417/1431	3/1431
5	<i>Amycolatopsis japonica</i>	HBUM173558	EU841704.1	99%	1415/1434	8/1434
6	<i>Amycolatopsis azurea</i>	IMSNU 20053T	AJ400709.1	99%	1414/1434	9/1434
7	<i>Amycolatopsis orientalis</i>	IMSNU 20057T	AJ293755.1	98%	1413/1435	11/1435
8	<i>Amycolatopsis regifaucium</i>	GY293	AY129769.1	98%	1409/1435	11/1435
9	<i>Amycolatopsis decaplanin</i>	DSM 44594	NR_025562.1	98%	1407/1433	8/1433

al. (1983) after growth on YEME Agar for 7-10 days at 28°C. Phenotypic tests were carried out following the procedures of Goodfellow et al. (1997) and Gordon et al. (1974). Chemotaxonomic characters were determined as described previously (Wu et al. 1989; Collins, 1985; Minnikin et al. 1984; Hasegawa et al. 1983; Lechevalier & Lechevalier 1980; Minnikin et al. 1980).

16S rDNA analysis

Genomic DNA was prepared by using standard methods (Maniatis et al. 1982; Marmur 1961). The G+C content of the DNA was determined using the Genetool software with PGEM as the control. The amplified product was sequenced using universal primers (5' to 3') fD2 (ccgaattcgtcgacaacAGAGTTTGATCATGGCTCAG) and rP1 (cccgggatccaagcttACGGTTACC-TTGTTACGACTT). The nucleotide sequence was obtained by processing DNA sequencing samples using ABI 3130 (4 capillary) electrophoresis instruments. The resultant 16S rDNA sequence was aligned using Gene Tool Lite version 1.0 software program against corresponding sequences retrieved from the Genbank database. The tree making algorithm used was Neighbor joining (Saitou & Nei, 1987) with Max Seq diff of 0.75. The phylogenetic tree was constructed (newick format) using Tree view software.

Fermentation

A full grown slant culture of the strain DVR D4 on YEME agar slant was transferred to Erlenmeyer flasks containing 50 ml seed medium (composition: soyabean meal, 10.0g; corn steep liquor, 10.0g; glucose, 5.0g; CaCO₃, 5.0g; sterilized natural aged sea water, 1.0L; pH 7.0) and incubated for 2 days at 28°C on a orbital shaker (220 rpm). A 5% of this inoculum was transferred into the optimized production medium (composition: D-glucose, 20.0g; malt extract, 40.0g; yeast extract, 4.0g; dipotassium hydrogen phosphate, 5.0g; sodium chloride, 2.5g; zinc sulphate, 0.04g; calcium carbonate, 0.4g; 1.0L sterile distilled water with pH 6.0) (Dasari et al. 2011a,b). The inoculated production flasks were incubated for 96 h at 28°C on a rotary shaker (220 rpm).

Antibacterial activity (cup-plate method)

The culture broth was centrifuged at 4000 rpm for 10 min, at 8°C and clear culture filtrate was separated for their antagonistic activity against the selected Gram-positive and Gram-negative bacteria by cup-plate method. Inoculated with test organisms, cups of 6 mm diameter were prepared in the nutrient agar plates and the cups are filled with the 50 µl of clear culture supernatant; and the diameter of inhibition zones were measured after incubation for 24 h at 37°C.

Table 2: Antibiotic resistance and inhibition similarities of *Amycolatopsis alba* var. nov DVR D4 and *Amycolatopsis alba*

Antibiotic	Strain D4	<i>Amycolatopsis alba</i> (Frederick and Raymond, 1993)
Cephalosporin (10 µg/ml)	+	N/A
Tetracycline (30 µg/ml)	+	+
Rifampicin (5 µg/ml)	+	+
Gentamycin (10 µg/ml)	+	-
Lincomycin (2 µg/ml)	+	-
Neomycin (30 µg/ml)	+	+
Penicillin (10 U)	-	-
Vancomycin (30 µg/ml)	-	-
Streptomycin (10 µg/ml)	-	-
Resistant, "+"; Inhibition, "-"		

Results and Discussion

An almost complete 16S rDNA sequence was determined for strain DVR D4 (1428 nucleotides). Comparison of this sequence with those of representation reference strains of the family *Pseudonocardaceae* shows that the organism belongs to the genus




Figure 2: Spore chain structure of *Amycolatopsis alba* var. nov DVR D4; phase contrast microscope X400

Amycolatopsis (Fig. 1). The GenBank accession number for the 16S rDNA sequence of strain D4 is JN872327. The 16S rDNA sequence similarity between the strain DVR D4 and its nearest neighbor, *Amycolatopsis alba*, is 100 %; this value corresponds to 0 differences out of 1428 nucleotide positions compared (Table 1). The chemotaxonomic data of the test strain are consistent with its assignment to the genus *Amycolatopsis* (Lechevalier et al. 1986). Strain DVR D4 contains *meso*-diaminopimelic acid as the wall




Figure 3: Scanning electron micrograph of *Amycolatopsis alba* var. nov DVR D4 grown on YEME medium for 7 days at 28°C; Magnification, X 9,000; Bar, 2.0µm.

diamino acid, arabinose and galactose as major wall sugars. It also contains phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylinositol diagnostic polar lipids but it lacks mycolic acids; the G+C content of the DNA is 58.9 mol%. This chemical profile serves to distinguish strain DVR D4 from members of all other wall chemotype IV taxa, except those classified in the genus *Amycolatopsis* (Holt et al. 1994; Embley 1992; Goodfellow & Lechevalier 1989; Henssen et al. 1987; Lechevalier et al. 1986). The classification of strain DVR D4 in the genus *Amycolatopsis* is also supported by phenotypic and morphological properties. The organism is aerobic, non-motile, Gram-positive, non-acid alcohol-

fast and produces branching substrate mycelium which fragments into cylindrical elements. Colonies have wrinkled surfaces and are covered with fine white hyphae; the hyphae are segmented into long strands of conidia that are arranged in a typical cobweb morphology characteristic of some nonstreptomycete actinomycetes (Fig. 2). The conidia are cylindrical and have a smooth surface (Fig. 3).

Strain DVR D4 was resistant to the following antibiotics ($\mu\text{g/ml}$): Cephalosporin (10), Tetracycline (30), Rifampicin (5), Gentamycin (10), Lincomycin (2) and Neomycin (30). It was inhibited by Penicillin (10 U), Vancomycin (30 $\mu\text{g/ml}$) and Streptomycin (10 $\mu\text{g/ml}$) (Table 2). The strain DVR D4 can be distinguished from the published *Amycolatopsis* species *Amycolatopsis alba* using a combination of phenotypic properties (Table 3). The chemical, genotypic and phenotypic data show that strain DVR D4 merits species status in the genus *Amycolatopsis*. Even though there was 100% pair-wise similarity and identity with no nucleotide gaps with

Table 4 Antibacterial activities of product produced by *Amycolatopsis alba* var. nov DVR D4

Test Organism	Zone of inhibition (mm)
<i>B. subtilis</i> NCIM 2439	18
<i>B. pumilus</i> NCIM 2327	16
<i>B. saccharolyticum</i> NCIM 2238	22
<i>S. aureus</i> NCIM 5021	20
<i>S. griseus</i> NCIM 2622	14
<i>E. coli</i> NCIM 2067	12
<i>A. formicans</i> NCIM 2319	20
<i>P. taetrolens</i> MTCC 1612	20

the species *Amycolatopsis alba* strain DSM 44262, there are substantial differences in some of the phenotypic characteristics (Tables 2 and 3). Therefore, it is proposed that this organism as a strain variety of *Amycolatopsis alba* and designated as *Amycolatopsis alba* var. nov DVR D4.

Table 3 Phenotypic characters of *Amycolatopsis alba* var. nov DVR D4

Character	Result	
	Strain D4	<i>Amycolatopsis alba</i> (Frederick and Raymond, 1993)
Gram staining	+	+
Spore staining	-	-
Motility	-	-
Color of aerial mycelium	White	White
Growth at 15°C	-	ND
Growth at 25°C	-	ND
Growth at 28°C	+	ND
Growth at 30°C	+	+
Growth at 37°C	-	ND
Growth at 42°C	-	ND
Growth at pH 5.0	+	ND
Growth at pH 6.0	+	ND
Growth at pH 6.5	+	ND
Growth at pH 7.0	+	ND
Growth at pH 7.5	+	ND
Growth at pH 8.0	+	ND
Growth at pH 9.0	+	ND
Growth at pH 10.0	+	ND
Growth on NaCl 2%	+	ND
Growth on NaCl 5%	-	+
Growth on NaCl 7%	-	ND
Growth on NaCl 10%	-	ND
Starch hydrolysis	-	ND
Casein hydrolysis	-	ND
Citrate utilization	-	+
Gelatin liquefaction	+	ND
H ₂ S production	-	ND
MR (Methyl Red)	-	ND
VP (Vogues Proskauer)	-	ND
Nitrate reduction	-	-
Indole	-	ND
Catalase	+	ND
Oxidase	-	ND
Urease	+	+
Acid production from		
Arabinose	-	+
Galactose	+	+
Glucose	+	ND
Mannitol	+	+
Raffinose	-	+
Salicin	-	+
Xylose	+	+
Sucrose	-	-
Rhamnose	-	-
Meso-inositol	-	+
Fructose	+	ND

+, positive or present; -, negative or absent; ND, not done

The antagonistic principles secreted by the *Amycolatopsis alba* var. nov. DVR D4 exhibited broad antagonistic spectrum against the tested bacterial species studied (Table 4). To the best of my knowledge from literature search, this was the first report of *Amycolatopsis alba* isolation from marine environment and also this was the first bioactive actinomycete strain of *Amycolatopsis alba* isolated in India.

Conclusion

The bioactive actinomycete was aerobic, non-motile, Gram-positive, non-acid filamentous, differentiated into substrate; abundant white aerial mycelium was produced and it also fragments into cylindrical elements. Cylindrical, smooth spores are formed with typical cobweb morphology. It grows on 2 % NaCl at 28°C and in pH range of 5.2 to 10.0. It was positive for urease, catalase and liquefaction of gelatin but negative for indole, oxidase, hydrolysis of starch and casein, utilization of citrate, H₂S production and nitrate reduction. Acid was produced from the following carbohydrates: galactose, glucose, mannitol, xylose and fructose. Acid was not produced from the arabinose, raffinose, salicin, sucrose, rhamnose and meso-inositol. The organism contains meso-diaminopimelic acid as the wall diamino acid, arabinose and galactose as major wall sugars. It also contains phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylinositol diagnostic polar lipids but it lacks mycolic acids; the G+C content of the DNA is 58.9 mol%.

Acknowledgements

Author Dasari is thankful to UGC, New Delhi, India for the financial assistance under UGC- Research Fellowships in Science for Meritorious Students.

References

- Albarracin VH, Alonso-Vega P, Trujillo ME, Amoroso MJ, Abate CM (2010) *Amycolatopsis tucumanensis* sp. nov., a copper-resistant actinobacterium isolated from polluted sediments. *Int J Syst Evol Microbiol* 60:397-401
- Atchareeya C, Pathom-Aree W, Yokota A, Chartchai K, Lumyong S (2010) *Amycolatopsis thailandensis* sp. nov., a poly (L-lactic acid)-degrading actinomycete isolated from soil. *Int J Syst Evol Microbiol* 61:839-843
- Carlsohn MR, Groth I, Tan GYA, Schutze B, Saluz HP, Munder T, Yang J, Wink J, Goodfellow M (2007) *Amycolatopsis saalfeldensis* sp. nov., a novel actinomycete isolated from a medieval alum slate mine. *Int J Syst Evol Microbiol* 57:1640-1646
- Chen J, Su JJ, Wei YZ, Li QP, Yu LY, Liu HY, Zhang YQ, Zhang YQ (2010) *Amycolatopsis xylanica* sp. nov., isolated from soil. *Int J Syst Evol Microbiol* 60:2124-2128
- Collins MD (1985) Isoprenoid quinone analysis in classification and identification. In: Goodfellow M, Minnikin DE (ed) *Chemical Methods in Bacterial Systematics*, Academic Press, London, pp 267-287
- Cwale Z, Igbinosa EO, Okoh AI (2011) Assessment of antibiotics production potentials in four actinomycetes isolated from aquatic environments of the Eastern Cape province of South Africa. *African J Pharm Pharmacol* 5:118-124
- Dasari VRRK, Nikku MY, Donthireddy SRR (2011a) Screening of antagonistic marine actinomycetes: optimization of process parameters for the production of novel antibiotic by isolate *Amycolatopsis alba* var. nov. DVR D4. *J Microbial Biochem Technol* (Article in Press)
- Dasari VRRK, Muthyala MKK, Nikku MY, Donthireddy SRR (2011b) Novel Pyridinium compound from marine actinomycete, *Amycolatopsis alba* var. nov. DVR D4 showing antimicrobial and cytotoxic activities *in vitro*. *Microbiol Res* (Article in Press)
- Demain AL, Zhang L (2005) Natural products and drug discovery. In: *Natural Products Therapeutic Medicine*, Humana Press, pp 3-32
- Duangmal K, Mingma R, Pathom-Aree W, Thamchaipenet A, Inahashi Y, Matsumoto A, Takahashi Y (2010) *Amycolatopsis samanae* sp. nov., isolated from roots of *Samanea saman* (Jacq.) Merr. *Int J Syst Evol Microbiol* 61:951-955
- Embley TM (1992) The family *Pseudonocardiaceae*. In: Balows A, Truper HG, Dworkin M, Harder W, Schleifer KH (ed) *The Prokaryotes*. Volume 1, Springer, Berlin, pp 996-1027
- Embley TM, Smida J, Stackebrandt E (1988) The phylogeny of mycolate-less wall chemotype IV Actinomycetes and description of *Pseudonocardiaceae* fam. nov. *Syst Appl Microbiol* 11: 44-52
- Frederick PM, Raymond CY (1993) *Amycolatopsis alba* sp. nov., isolated from soil. *Int J Syst Bacteriol* 43:715-720
- Goodfellow M, Lechevalier MP (1989) The genus *Nocardia*. In: Williams ST, Sharpe ME, Holt JG (ed) *Bergey's Manual of Systematic Bacteriology*, Williams & Wilkins, Baltimore, pp 2350-2361
- Goodfellow M, Brown A, Cai J, Chun J, Collins MD (1997) *Amycolatopsis japonicum* sp. nov., an actinomycete producing (S,S)-N,N'-ethylenediamine-disuccinic acid. *Syst Appl Microbiol* 20:78-84
- Goodfellow M, Kim SB, Minnikin DE, Whitehead D, Zhou ZH, Mattinson-Rose AD (2001) *Amycolatopsis sacchari* sp. nov., a moderately thermophilic actinomycete isolated from vegetable matter. *Int J Syst Evol Microbiol* 51:187-193
- Gordon RE, Barnett DA, Handerhan JE, Pang CHN (1974) *Nocardia coeliaca*, *Nocardia autotrophica* and the *nocardin* strain. *Int J Syst Bacteriol* 24:54-63
- Groth I, Tan GYA, Gonzales JM, Laiz L, Carlsohn MR, Schutze B, Wink J, Goodfellow M (2007) *Amycolatopsis nigrescens* sp. nov., an actinomycete isolated from a Roman catacomb. *Int J Syst Evol Microbiol* 57:513-519
- Hasegawa T, Takizawa M, Tanida S (1983) A rapid analysis for chemical grouping of aerobic actinomycetes. *J Gen Appl Microbiol* 29:319-322
- Henssen A, Kothe HW, Kroppenstedt RM (1987) Transfer of *Pseudonocardia azurae* and *Pseudonocardia fastidiosa* to the genus *Amycolatopsis*, with amended species description. *Int J Syst Bacteriol* 37:292-295
- Holt JG, Krieg NR, Sneath PHA, Staley JT, Williams ST (1994) *Bergey's manual of determinative bacteriology* (9th edn). The Williams and Wilkins Co., Baltimore, Maryland, USA
- Huang Y, Pas Ciak M, Liu ZH, Xie Q, Gamian A (2004) *Amycolatopsis palatopharyngis* sp. nov., a potentially pathogenic actinomycete isolated from a human clinical source. *Int J Syst Evol Microbiol* 54:359-363
- Huang Y, Weihong Q, Zhitang L, Zhiheng L, Michael G (2001) *Amycolatopsis rubida* sp. nov., a new *Amycolatopsis* species from soil. *Int J Syst Bacteriol Evol Microbiol* 51:1093-1097
- Kim B, Sahin N, Tan GYA, Zakrzewska-Czerwinska J, Goodfellow M (2002) *Amycolatopsis eurytherma* sp. nov., a thermophilic actinomycete isolated from soil. *Int J Syst Evol Microbiol* 52:889-894
- Krishna PSM, Venkateshwarlu G, Rao LY (1998) Studies on fermentative production of rifamycin using *Amycolatopsis mediterranei*. *J Microbiol Biotechnol* 14:689-691
- Labeda DP, Donahue JM, Williams NM, Sells SF, Henton MM (2003) *Amycolatopsis kentuckyensis* sp. nov., *Amycolatopsis*

- lexingtonensis* sp. nov. and *Amycolatopsis pretoriensis* sp. nov., isolated from equine placentas. *Int J Syst Evol Microbiol* 53:1601–1605
- Labeda DP, Goodfellow M, Chun J, Zhi XY, Li WJ (2011) Reassessment of the systemic of the suborder *Pseudonocardineae*: transfer of the genera within the family *Actinosynnemataceae* Ladeba and Kroppenstedt 2000 emend. Zhi et al. 2009 into an emended family *Pseudonocardineae* Embley et al. 1989 emend. Zhi et al. 2009. *Int J Syst Evol Microbiol* 61:1259–1264
- Lechevalier HA, Lechevalier MP (1980) The chemotaxonomy of actinomycetes. In: Dietz A, Thayer DW (ed) *Actinomycete Taxonomy* (Society for Industrial Microbiology Special Publication Number 6), Society of Industrial Microbiology, Arlington, pp 277–284
- Lechevalier MP, Prauser H, Labeda DP, Ruan JS (1986) Two genera of nocardioform Actinomycetes: *Amycolata* gen. nov. and *Amycolatopsis* gen. nov. *Int J Syst Bacteriol* 36:29–37
- Lee SD, Kinkel LL, Samac DA (2006) *Amycolatopsis minnesotensis* sp. nov., isolated from a prairie soil. *Int J Syst Evol Microbiol* 56:265–269
- Liras P, Demain AL (2009) Enzymology of beta-lactam compounds with cephem structure produced by actinomycete (Chapter 16). *Methods Enzymol* 458:401–429
- Malkova IV, Borisova OK, Pavlova MV, Zemlianitskaia EP, Serquuva TI (1991) *In vitro* activity of a new glycopeptides antibiotic eremomycin in relation to obligate anaerobic gram positive bacteria. *Antibiot Khimioter* 36:17–20
- Maniatis T, Fritsch EF, Sambrook J (1982) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York
- Marmur J (1961) A procedure for the isolation of deoxyribonucleic acid from microorganisms. *J Mol Biol* 3:208–218
- Minnikin DE, Hutchinson IG, Caldicott AB, Goodfellow M (1980) Thin-layer chromatography of methanolysates of mycolic acid-containing bacteria. *J Chromatogr* 188:221–233
- Minnikin DE, O'Donnell AG, Goodfellow M, Alderson G, Athalye M, Schaal A, Parlett JH (1984) An integrated procedure for the extraction of isoprenoid quinones and polar lipids. *J Microbiol Methods* 2:233–241
- Ramesh S, Narayanasamy M (2009) Screening of marine actinomycetes isolated from the Bay of Bengal, India for antimicrobial activity and industrial enzymes. *World J Microbiol Biotechnol* 25:2103–2111
- Saintpierre-Bonaccio D, Amir H, Pineau R, Tan GYA, Goodfellow M (2005) *Amycolatopsis plumensis* sp. nov., a novel bioactive actinomycete isolated from a New-Caledonian brown hypermagnesian ultramafic soil. *Int J Syst Evol Microbiol* 55:2057–2061
- Saitou N, Nei M (1987) The neighbor joining method: a new method for constructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Soon Dong Lee (2006) *Amycolatopsis jejuensis* sp. nov. and *Amycolatopsis halotolerans* sp. nov., novel actinomycetes isolated from a natural cave. *Int J Syst Evol Microbiol* 56:549–553
- Tamura T, Ishida Y, Otaguro M, Suzuki K (2010) *Amycolatopsis helveola* sp. nov. and *Amycolatopsis pigmentata* sp. nov., isolated from soil. *Int J Syst Evol Microbiol* 60:2629–2633
- Tan GYA, Robinson S, Lacey E, Goodfellow M (2006) *Amycolatopsis australiensis* sp. nov., an actinomycete isolated from arid soils. *Int J Syst Evol Microbiol* 56:2297–2301
- Tang SK, Wang Y, Guan TW, Lee JC, Kim CJ, Li WJ (2010) *Amycolatopsis halophila* sp. nov., a halophilic actinomycete isolated from a salt lake. *Int J Syst Evol Microbiol* 60:1073–1078
- Trenin AS, Fedorova GB, Laiko AV, Dudnik IuV (2001) Increase in eremomycin production by regeneration and UV-irradiation of *Amycolatopsis orientalis* subsp. *eremomycini* protoplasts. *Antibiot Khimioter* 46:6–11
- Warwick S, Bowen T, McVeigh H, Embley TM (1994) A phylogenetic analysis of the family *Pseudonocardiaceae* and the genera *Actinokineospora* and *Saccharothrix* with 16S rRNA sequences and a proposal to combine the genera *Amycolata* and *Pseudonocardia* in an emended genus *Pseudonocardia*. *Int J Syst Bacteriol* 44:293–299
- Williams ST, Cross T (1971) *Actinomycetes, methods in microbiology*. Volume 4, Academic press, New York
- Williams ST, Goodfellow M, Alderson G, Wellington EMH, Sneath PHA, Sackin MJ (1983) Numerical classification of *Streptomyces* and related genera. *J Gen Microbiol* 129:1743–1813
- Wu C, Lu X, Qin M, Wang Y, Ruan J (1989) Analysis of menaquinone compound in microbial cells by HPLC. *Microbiology* 16:176–178
- Zhang L, An R, Wang J, Sun N, Zhang S, Hu J, Kuai J (2005) Exploring novel bioactive compounds from marine microbes. *Curr Opin Microbiol* 8:276–281