

Evaluation of Renaoprotective Effect of *Costus Afer* Leaf Extract on Rats exposed to Cyclosporine: Antioxidant and Anti-inflammatory Pathways

Maha A. Althaiban

Received: 23 October 2018 / Received in revised form: 11 March 2019, Accepted: 17 March 2019, Published online: 10 April 2019
© Biochemical Technology Society 2014-2019
© Sevas Educational Society 2008

Abstract

Costus afer leaf (Cost) extract used as traditional herbal therapy for various diseases. Long term treated with cyclosporine (CsA) concomitant with development of renal toxicity. The current study aims to assess the nephroprotective mechanism effect of Cost in CsA nephrotoxic rats. Forty male rats were distributed into 4 groups; control (Cont), CsA; rats intraperitoneal (i.p) injected with CsA (twenty five mg/kg b.wt) for twenty one days, Cost 750 mg/kg b.wt +CsA, Cost 1500 mg/kg b.wt +CsA; rats received Cost orally for twenty one days, followed by i.p. injection with CsA. Biochemical samples were collected after 24 h from the last dose of CsA as well as renal tissue samples were collected for histopathological examination. Renal oxidative stress biomarkers (lipid peroxides (MDA) and catalase (CAT)) and the serum anti-inflammatory biomarkers (tumor necrosis factor interleukin-1beta (IL-1 β) and -alpha (TNF- α)) were determined. Kidney functions; serum levels of creatinine, uric acid and urea as well as serum ionic levels of sodium Na⁺ and potassium K⁺ were measured. The results of this study revealed that injection of CsA induced significant increase in renal MDA, serum anti-inflammatory cytokines (TNF- α and IL-1 β), serum kidney function parameters and the serum ionic K⁺ levels, with significant decrease in the renal CAT and serum ionic Na⁺ compared with control group. Renal tissues showing congestion, focal hemorrhage and interstitial nephritis, with coagulating necrosis of the renal tubules in the CsA group. Oral administration of Cost extract significantly ameliorated CsA-induced renal oxidative stress. It reduced CsA-induced elevation in serum anti-inflammatory cytokines and kidney function parameters, as well as the changes in ionic Na⁺ and K⁺ levels compared with CsA group. It also protected against CsA-induced histopathological changes. Therefore, Cost extract ameliorates nephrotoxicity caused by CsA through antioxidant and anti-inflammatory mechanisms.

Keywords: *Costus afer* leaf, Cyclosporine, renal toxicity, antioxidant, anti-inflammatory.

Maha A. Althaiban

Food and Nutrition Department, Faculty of Home Economics, King Abdulaziz University, Jeddah, Saudi Arabia.

Introduction

Kidney is the organ in the body that removes blood metabolic waste such as uric acid, urea and creatinine and excrete them by urine; the kidney also removes drugs and various toxic agents in the blood (Kunworarath et al., 2014). One of the most common kidney problems is nephrotoxicity, which is caused by toxic compounds that may be endogenous or exogenous such as chemotherapeutic agents, antiplastic agents (cyclosporine and cyclophosphamide), aminoglycosides and antibiotics, which are the most important exogenous toxic agents (Sembulingam and Sembulingam, 2010). Drugs can cause oxidative stress by producing free radicals that are mostly available as by-products or as metabolic by-products (Goyal et al., 2016). The damage to nucleic acids, membrane lipids and tissue proteins caused by free radicals when generated excessively at the cellular level (Lakshmi, Muvvala and Shashank, 2014). The drug CsA is the most widely used medicine in organ transplantation, which has immunosuppressive effect, currently CsA had been investigated for a wide range of autoimmune diseases in clinical trials (Capasso et al., 2008). Other adverse effects of CsA are hyperactivity, encephalopathy, neurotoxicity, hypertension and nephrotoxicity, in addition to the immunosuppressive effect (Gnieszka et al., 2010). The generation of reactive oxygen species is widely attributed to nephrotoxicity due to chronic CsA treatment (Bobadilla and Gamba, 2007).

Management of nephrotoxicity is still a challenge to the modern scientific community. Unfortunately, drugs have little to offer alleviation of kidney ailments. Thus given rise to research involved in identification of safe, inexpensive and available alternatives from natural resources (Kotnis et al., 2004). For several years, plants have been used as medicine to help maintain human health (Uy et al., 2019). And, medicinal plants have also been widely used as medicinal and aromatic plants since the ancient times (Niazi et al., 2019). Due to the presence of various bioactive compounds, medicinal plants have been proved to be a major source of drugs (Pereira, Barros and Ferreira, 2016). Bioactive compounds with antioxidant property can prevent and combat oxidative stress-related diseases (Anyasor et al., 2010). Recent studies revealed the Cost extract has significant

nephroprotective properties (Gnieszka et al., 2010; Kotnis et al., 2004).

Costus afer Ker Gawl of the Zingiberaceae family, now referred to as Costaceae, is a herbaceous, rhizomatous monocotyledon. The common names for Cost include ginger lily, bush cane or spiral ginger (Iwu, 2014). The Cost leaves and stem as medicinal plant parts have received a lot of attention as a result of its antioxidant activity, as well as in treating a vast number of disease conditions including malaria, rheumatoid arthritis, diarrhea, diabetes mellitus, stomach ache, cough and cold (Anaga et al., 2004). *CostusAfer* has proven to possess hepatoprotective, nephroprotective antimicrobial, anti-inflammatory and hypolipidemic effects (Anyasor et al., 2014). The Cost active compounds are flavonoids, tannins, alkaloids; phenols and saponins, which are powerful water-soluble antioxidants that prevent oxidizing damage to cells (Anyasor et al., 2010). Fruits and vegetables have been recognized as natural sources of various bioactive compounds (Messouadi et al., 2019). The purpose of the present work is to evaluate the potential protective role of Cost against kidney toxicity Induced by CsA.

Material and Methods

Drugs and chemicals and plant material

Cyclosporine A (CsA) soft capsules, provided by Novartis Pharmaceuticals, Australia. All chemical and kits with high grade obtained from Sigma-Aldrich (St. Louis, MO) Chemical Co. *Costus afer* (Cost) leaves was purchased from local market in Jeddah, Saudi Arabia.

Preparation of Costus afer leaf extract

Costus afer leaves dried powder (500 g) was macerated in 1 L of ethanol (80%) at room temperature and mixed for 48 hrs by magnetic stirrer at 100 rpm speed. The extract was concentrated at 40°C under vacuum using rotary evaporator, the condensed residue was further freeze-dried (Siddhuraju and Becker, 2003). The extract was stored in non-permeable glass containers at 4°C until used.

Experimental design

Forty male rats weighting 120-150 g purchased from animalhouse of King Fahd Medical Research Center, KAU. They were adhering under the rules of Canadian ethical approval from the local biomedical ethical committee of KAU with free water and standard diet. After one week of acclimatization rats were distributed into 4 groups (n=10 in each group). Group I (Cont); rats received distilled water for 2 weeks then intraperitoneal (i.p) injected with olive oil (vehicle) for 21 days. Group II (CsA); rats distilled water for 2 weeks then were i.p. injected with CsA at a dose (25 mg/kg b.wt) diluted in olive oil for a period of twenty one days according to Chandramohan and Parameswar (Chandramohan and Parameswari, 2013). Group III (Cost 750 mg/ kg b.wt +CsA) (Ezejiofor, Orish and Orisakwe, 2014). Group IV (Cost 1500 mg/

kg b.wt +CsA). Rats in groups III and IV received Cost orally for twenty one days, followed by i.p. injected with CsA.

Samples collection

Twenty four hours after the last injected dose of the CsA drug, rats were anesthetized then blood and renal samples were collected. The serum samples were separated and stored at -80 °C until used for biochemical analysis. The renal samples were either prepared for biochemical analysis or preserved in neutral buffer formaldehyde solution for histopathological studies.

Biochemical analysis

- **Renal oxidative stress biomarkers;** the lipid peroxidation level (MDA) and the activity of catalase (CAT) enzyme were estimated in homogenized renal using ELISA kits.
- **The anti-inflammatory cytokines;** these serum levels of tumor necrosis interleukin-1beta (IL-1 β) and - α (TNF- α) were assessed by ELISA kits.
- **The level of kidney functions;** blood urea nitrogen (BUN), uric acid (UA) and serum levels of creatinine (Cr), as well as serum levels of sodium (Na⁺) and potassium (K⁺) were measured using colorimetric kits.
- All kits used were obtained MyBioSource, USA followed the steps and instructions of kits.

Histopathological studies

A kidney tissue portion from each group was fixed in formaldehyde solution and prepared for examined under microscope after staining with hematoxylin and eosin to determine histopathological changes.

Statistical analysis

All results were analyzed by SPSS ver. 22, by using ANOVA, values are expressed as mean \pm SEM, P-value < 0.05 considered significance.

Results

Effect of Cost on renal oxidative stress in nephrotoxic rats by CsA

The levels of non-enzymatic MDA and enzymatic CAT antioxidant levels in the renal tissue in different groups are represented in Figure (1). The level of MDA was significantly (p < 0.001) increased with significantly (p < 0.001) decreased in CAT activity in CsA compared with the control group. Oral administration of Cost induced significant improvement in the antioxidant status of renal. The MDA level decreased significantly (p < 0.001) and the CAT enzyme activity increased significantly (p < 0.001) in the Cost administration (750 mg/kg) and the (1500 mg/kg) groups as compared with CsA group. Significant (p <

0.05) changes were observed in Cost (750 mg/kg)+ CsA and Cost (1500 mg/kg)+ CsA, which indicated the protective effect of Cost is a dose-dependent. .

Effect of Cost on serum anti-inflammatory cytokines in nephrotoxic rats by CsA

The serum levels of IL1- β and TNF- α in different groups are showed in Figure (2). The levels of TNF- α and IL1- β were significantly ($p < 0.001$) increased in CsA compared with the control group. Oral administration of Cost induced significant decreases in the serum anti-inflammatory cytokines status, there were significant ($p < 0.001$) decreases in TNF- α and IL1- β levels in both Cost groups as compared with CsA intoxicated group. Significant ($p < 0.05$) changes were observed in Cost (750 mg/kg) + CsA and Cost (1500 mg/kg)+ CsA.


Effect of Cost on renal functions in nephrotoxic rats by CsA

Table (1) represents the levels of serum renal functions (Cr, BUN and UA) in different groups of experiments. The serum of kidney functions level in the CsA group was significantly ($p <$

0.001) increased in CsA group compared to control group. Cost administration (750 and 1500 mg/kg) reduced the serum kidney function levels significantly ($p < 0.001$) as compared with CsA group, which may be indicator for renal toxicity induced by CsA. Significant ($p < 0.05$) changes were observed in Cost (750 mg/kg) + CsA and Cost (1500 mg/kg) + CsA, which indicated the protective effect of Cost is a dose-dependent.


Effect of Cost on ionic sodium and potassium in nephrotoxic rats by CsA

Figure (3) shows the levels of serum ionic Na⁺ and K⁺ in different experimental groups. The serum level of ionic Na⁺ was significantly ($p < 0.001$) decreased with significantly ($p < 0.001$) increased in the serum ionic K⁺ in CsA group compared with control group. Oral Cost administration significantly ($p < 0.001$) increased the ionic Na⁺ level with a significant ($p < 0.001$) increase in ionic K⁺ level in the Cost administration groups as compared with CsA intoxicated group. Significant ($p < 0.05$) changes were observed in Cost (750 mg/kg) + CsA and Cost (1500 mg/kg) + CsA.



In each group N = 10, values are expressed as mean \pm SEM, values. $p \leq 0.05$ considered significant; [@] versus Control, [&] versus CsA values. [#] between Cost (750 mg/kg) and Cost (1500 mg/kg) groups.

Figure 1: Effect of Cost extract on renal non-enzymatic (MDA) and enzymatic (CAT) levels against CsA-induced nephrotoxicity in rats




In each group N = 10, values are expressed as mean ± SEM, values. p ≤ 0.05 considered significant; @ versus Control, & versus CsA values. # between Cost (750 mg/kg) and Cost (1500 mg/kg) groups.

Figure 2: Effect of Cost extract on serum inflammatory TNF-α and IL1-β levels against CsA-induced nephrotoxicity in rats

Table 1: Effect of Cost extract on renal functions against CsA-induced nephrotoxicity in rats

Groups	Renal function Parameters		
	Cr (μmol/L)	BUN (mmol/L)	UA(umol/L)
Control	22.50 ± 1.04	7.86 ± 0.44	62.20 ± 2.62
CsA	50.48 ± 3.56 ^a	21.09 ± 1.48 ^a	122.59 ± 5.55 ^a
Cost (750 mg/kg) + CsA	33.52 ± 2.77 ^b	11.69 ± 1.15 ^b	76.95 ± 2.96 ^b
Cost (1500 mg/kg) + CsA	28.19 ± 1.61 ^{b,c}	8.50 ± 0.63 ^{b,c}	66.37 ± 2.69 ^{b,c}

In each group N = 10, values are expressed as mean ± SEM, values. p ≤ 0.05 considered significant; ^a versus Control, ^b versus CsA values. ^c between Cost (750 mg/kg) and Cost (1500mg/kg) groups.



In each group N = 10, values are expressed as mean ± SEM, values. p ≤ 0.05 considered significant; @ versus Control, & versus CsA values. # between Cost (750 mg/kg) and Cost (1500 mg/kg) groups.

Figure 3: Effect of Cost extract on serum ionic Na⁺ and potassium K⁺ levels against CsA-induced nephrotoxicity in rats

Histopathological results

Kidney of control group showing the normal histological structure. In the CsA group, renal tissue showing congestion of renal blood vessels, focal hemorrhage and interstitial nephritis, with coagulating necrosis of the renal tubules associated with

mononuclear cells infiltration. In the Cost (750 mg/kg) + CsA grouprenal tissue slight congestion of the glomerular tufts with slight vacuolation of some renal tubular were seen. Apparent normal appearance of renal tissue was revealed in the Cost (1500 mg/kg) + CsA group.




Figure 4: Effects of Costus Afer (Cost) leaves extract on the kidney tissue histopathological changes detected by H&E staining in CsA-

induced nephrotoxicity in rats (Power magnification x 66). Photo A represent control group, renal tissue showing the normal histological structure. Photo B represent CsA group, renal tissue showing congestion of renal blood vessels (small arrow), as well as focal hemorrhage (large arrow). Photo C represent CsA group, renal tissue showing focal interstitial nephritis (arrow), with necrosis (star). Photo D represent CsA group, renal tissue showing coagulating necrosis of the renal tubules associated with mononuclear cells infiltration (star).




Figure 5: Effects of *Costus Afer* (Cost) leaves extract on the kidney tissue histopathological changes detected by H&E staining in CsA-induced nephrotoxicity in rats (Power magnification x 66). Photo A represent Cost (750 mg/kg) +CsA group, renal tissue showing slight congestion of the glomerular tufts (arrows). Photo B represent Cost (750 mg/kg) +CsA group, renal tissue showing slightly necrobiotic changes and vacuolation of some renal tubular (arrows). Photo C represent Cost (750 mg/kg) +CsA group, renal tissue showing the normal appearance.

Discussion

The study was conducted to assess the protective effect and antioxidant activity of Cost against CsA induced nephrotoxicity. The use of CsA was found to be nephrotoxic in the first attempt after transplantation for immunosuppression (Lorber et al., 1987). The results showed that i.p. injection with CsA in a dose of 25 mg/kg b.wt. diluted in olive oil for a period of twenty one days, resulted in deterioration of renal function and the development of histopathological changes in the renal tissues. The results were in the same line with Gnieszka et al. (2010) who demonstrated that using CsA in a dose of 25 mg/kg causes a significant but unexpected nephrotoxicity that was not seen in initial animal experiments. Several studies have identified nephrotoxicity as the most common and clinically significant adverse effect of CsA (Gnieszka et al., 2010; Kotnis et al., 2004).

In the present study there were significant increases in renal MDA, anti-inflammatory cytokines (TNF- α and IL-1 β), kidney function parameters and the serum ionic K⁺ levels, with significant decreases in the renal CAT and serum ionic Na⁺ in intoxicated group compared with control group. The results obtained correspond to most of the experimental procedures reported (Agnieszka et al., 2010; Ciarcia, Damiano and Fiorito, 2012; Wongmekiat et al., 2011). This could be due to an increase in renal vascular resistance, a significant reduction in renal blood flow and activation of the sympathetic renal nervous system

which had been demonstrated as a concomitant stimulation of plasma renin activity (Elsayed, Bayomy and Azab, 2016). The exact cause for nephrotoxicity by CsA administration remains obscure, but several studies suggested deficiencies in magnesium, intracellular calcium, nitric oxide and oxidative stress (Parra et al., 1998). The generation of reactive oxygen species is widely attributed to nephrotoxicity due to chronic CsA treatment (Mohamadin et al., 2005).

The present study confirmed that, there has been a significant ($p < 0.05$) increases in the level of renal tissue MDA as well as in the level of TNF- α and IL-1 β pro-inflammatory cytokines compared with the extract pretreated groups. The results were consistent with other researchers who demonstrated that there is a significant increase in lipid peroxidation and in pro-inflammatory cytokines during CsA administration (Fong et al., 1990). Excessive MDA levels in kidney tissue and in pro-inflammatory cytokines had been taken into considered as an indicator of cellular harm because of excessive lipid peroxidation processes during antioxidant defense system malfunction (Marsoul et al., 2016).

In contrast, oral administration of Cost prevented this elevation in a dose dependent manner. The results were in agreement with Uboh et al., (2014), who reported that Cost has a remarkable inhibition of lipid peroxidation and radical scavenging due to its phytoconstituents such as flavonoids, alkaloids, saponins, phenols, terpenoids, tannins, and cardiac glycosides. Alkaloids

are known to have anti-inflammatory effect while, the other water-soluble antioxidants prevent cell damage from oxidizing and suggesting anti-inflammatory properties (Ukpai et al., 2012).

Antioxidant catalase is one of the major defenses against oxidative damage caused by ROS (Hussein, Ragab and El-Eshrawy, 2014). Intoxication with CsA resulted in a significant reduction in the activity of CAT enzyme compared to the untreated group. The reduction in CAT activity level may be due to NADPH production depletion during CsA administration (Anyasor et al., 2010). Oral administration of Cost with CsA led to significant elevation in renal tissue antioxidant activity compared with intoxicated group with CsA. The same effect were reported by Anyasor *et al.* (2014) who stated that the effect could be due to phytochemicals present in the Cost such as flavonoids, polyphenols, tannins and saponins which are known as plant antioxidant metabolites. Considering all types of natural antioxidants, polyphenols can be regarded as the major potent compound, because of being abundantly applied in the food industry, cosmetic, pharmaceutical and medicinal materials (Tabet et al., 2018).

This study also included several important biochemical parameters. These biological parameters are organ toxicity indicators. For example, serum urea, creatinine, uric acid and electrolytes (Na⁺ and K⁺ levels) which were used to evaluate kidney functions. In the current study, there were an increase in concentrations of serum uric acid, urea, creatinine and K⁺ level while there was a decrease in Na⁺ level detected in the CsA group compared with the normal group. Chronic Administration of CsA for 21 days resulted in deficiencies in renal functions characterized by high serum levels of uric acid, creatinine and urea, which could be used as a rough glomerular filtration index, indicating multiple kidney disorders (Burdmann, Andoh and Bennett, 2003). Pretreatment with Cost to CsA treated rats elevate the changes caused by CsA. The same results were reported by Uboh *et al.*, (2014) who stated that rats orally given Cost restored the levels of kidney function parameters (creatinine, urea and uric acid).

The present study demonstrated that administration of CsA lead to a significant decrease in potassium level and significant increase in serum sodium level in pretreated Cost groups compared to CsA group. The same results were seen in the study of Ezejiofor *et al.* (2014) who stated that rats orally given Cost significantly reduced creatinine and serum blood urea as well as sodium level and increased potassium level in gentamicin induced nephrotoxicity.

Histopathological studies had been carried out on all rats. The study results revealed a significant influence of the CsA administration on histopathological damage of CsA group. This results were in the same line with Gnieszka *et al.* (2010). On the other hand, there was an improvement in kidneys tissue of rat groups given Cost in combination with CsA.

In Conclusion, the results of the recent study demonstrate that Cost has a protective role that is likely mediated by its antioxidant

properties against CsA-induced nephrotoxicity. Further studies are needed to evaluate its potential utility of this extract in clinical conditions associated with renotoxicity.

References

- Agnieszka, K., Mariusz, M., Grażyna C., Krzysztof, C. and Elżbieta, K. (2010). Experimental models of cyclosporine nephrotoxicity. *Bull Vet InstPulawy* 54, 59-62.
- Anaga, A.O., Njoku, C.J., Ekejiuba, E.S., Esiaka, M.N and Asuzu, I.U.(2004). Investigations of the methanolic leaf extract of *Costusafer Ker* for pharmacological activities in vitro and in vivo. *Phytomedicine* , 11:242-8.
- Anyasor, G.N., Olusola, K., Oyelana, O.A and Akpofunre, B.E. (2010). Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costusafer Ker* Gawl. (Costaceae).*Afr. J. Biotechnol.*, 31:4880-4.
- Anyasor, G.N., OlusolaOgunwenmo, K., Oyelana, O.A and Akpofunre, B.E. (2010). Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costusafer Ker* Gawl. (Costaceae). *Afr. J. Biotechnol.* , 31:4880-4.
- Anyasor, G.N., OlusolaOgunwenmo, K., Oyelana, O.A. and Akpofunre, B.E.(2010). Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costusafer Ker* Gawl. (Costaceae).*Afr. J. Biotechnol.* , 31:4880-4.
- Anyasor, G.N., Onajobi, F., Osilesi, O., Adebawo, O. and Oboutor, E.M. (2014). Anti-inflammatory and antioxidant activities of *Costusafer Ker* Gawl. hexane leaf fraction in arthritic rat models. *J Ethnopharmacol* 2014; 155:543-51.
- Anyasor, G.N., Onajobi, F.D., Osilesi, O and Adebawo, O.O. (2014). Proximate composition, mineral content and in vitro antioxidant activity of leaf and stem of *Costusafer* (Ginger lily). *J. of Intercultural Ethnopharm.*, 3(3):128-34.
- Bobadilla, N. and Gamba, G. (2007). New insights into the pathophysiology of cyclosporine nephrotoxicity: a role of aldosterone. *Am J Renal Physiol*, 293, 2-9.
- Burdmann, E.A., Andoh, T. F. and Bennett, W.M. (2003) Cyclosporine nephrotoxicity. *SeminNephrol.*, 23(5):465-76.
- Capasso, G., Di Gennaro, C., Ragione, F., Manna, K., Ciarcia, R., Florio, S., Perna, A., Pollastro, R., Damiano, S., Mazzoni, O., Galletti, P. and Zappia V. (2008). In vivo effect of the natural antioxidant hydroxytyrosol on cyclosporine nephrotoxicity in rats. *Nephrol Dial Transplant*, 23, 1186-95.
- Chandramohan, Y. and Parameswari, C.S. (2013). Therapeutic efficacy of naringin on cyclosporine (A)- induced nephrotoxicity in rats: Involvement of hemeoxygenase-1. *Pharmacological Reports*, 65, 1336-44.
- Ciarcia, R., Damiano, S. and Fiorito, F. (2012). Hydrocortisone attenuates cyclosporin A-induced nephrotoxicity in rats. *J Cell Biochem*. 113:997-1004.
- Elsayed,A.S.I., Bayomy,M.F.F. and Azab, A.E. (2016). Effect of acute and chronic treatment of Cyclosporin A on liver and

- kidney functions in rats *Journal of Applied Pharmaceutical Science* 6 (03): 116-9.
- Ezejiolor, A.N., Orish, C.N and Orisakwe, O.E. (2014). Effect of aqueous leaves extract of *Costusafer Ker Gawl* (Zingiberaceae) on the liver and kidney of male albino Wister rat. *Anc. Sci. Life*, 33(1):4-9.
- Ezejiolor, A.N., Orish, C.N. and Orisakwe, O.E. (2014). *Costusafer Ker Gawl* leaves against gentamicin-induced nephrotoxicity in rats. *Kidney diseases*, 4 (8): 310-3.
- Fong, Y., Moldawer, L.L., Shires, G.T. and Lowry, S.F. (1990). The biologic characteristics of cytokines and their implication in surgical injury. *Surgery Gynecology and Obstetrics.*, 170(4): 363-78.
- Gnieszka, K., Mariusz M., Grażyna C., Krzysztof C. and Elżbieta K. (2010). Experimental models of cyclosporine a nephrotoxicity, *Bull Vet InstPulawy* 54: 59-62.
- Gnieszka, K., Mariusz M., Grażyna C., Krzysztof C. and Elżbieta K. (2010). Experimental models of cyclosporine a nephrotoxicity, *Bull Vet InstPulawy* 54: 59-62.
- Gnieszka, K., Mariusz, M., Grażyna, C., Krzysztof, C. and Elżbieta, K. (2010). Experimental models of cyclosporine a nephrotoxicity, *Bull Vet InstPulawy* 54: 59-62.
- Goyal, S.N., Mahajan, U.B., Chandrayan, G., Kumawat, V.S., Kamble, S., Patil, P., Agrawal, Y.O., Patil, C.R. and Ojha, S. (2016). Protective effect of oleanolic acid on oxidative injury and cellular abnormalities in doxorubicin induced cardiac toxicity in rats. *Am. J. of Translational Res.*, 8(1): 60-9.
- Hussein, S.A., Ragab, O.A. and El-Eshrawy, M.A. (2014). Protective effect of green Tea extract on cyclosporine a-induced nephrotoxicity in rats. *J of Biol. Sci.*, 14:248-57.
- Iwu, M.M. (2014). *Handbook of African Medicinal Plants*, CRC Press, 2nd edition, Boca Raton, New York, London.
- Kabouche, Z. 2019. Anti-hyperlipidemic, Anti-inflammatory and Antioxidant Activities of *Citrullus lanatus*. *World J Environ Biosci*, 8 (1), pp: 100-106.
- Kotnis, M.S., Patel, P., Menon, S.N and Sane, R.T. (2004). Renoprotective effect of *Hemidesmus indicus*, a herbal drug used in gentamicin-induced renal toxicity. *Nephrol.*, 9(3):142-52.
- Kunworarath, N., Muangnil, P., Itharat, A and Hiranyachattada, S. (2014). Acute and sub-chronic treatment of *Hibiscus sabdariffa* Linn. extract on renal function and lipid peroxidation in cisplatin-induced acute renal failure rats. *J. Physiol.*, 27(1): 5-12.
- Lakshmi, B.V.S., Muvvala, S and Shashank, P. (2014). Protective effect of *Vitisvinifera* in doxorubicin-induced oxidative stress in rats-A preliminary study. *Inter. J. of Adv. in Pharmacy Med. and Bioallied Sci.*, 2:7-14.
- Lorber, M.I., Van Buren, C.T., Flechner, S.M., Williams, C. and Kahan, B.D. (1987). Hepatobiliary and pancreatic complications of cyclosporine therapy in 466 renal transplant recipients. *Transplantation*, 43(1):35-40.
- Marsoul, R.D., Abbood, R.M. and Abbas, M.T. (2016). Effect of garlic oil on cyclosporine induced renal toxicity in rats. *Human J.*, 5(2): 209-221.
- Messaoudi, S.Tebibel, S. Beladjila, A.K. Khelifi Touhami, F. & Mohamadine, A.M., El-Beshbishy, H.A and El-Mahdy, M.A. (2005). Green tea extract attenuates cyclosporine A-induced oxidative stress in rats. *Pharmacological Research*. 51: 51-7.
- Niazi, M.Yari, F. Shakarami, A. 2019. A Review of Medicinal Herbs in the Lamiaceae Family Used to Treat Arterial Hypertension, *Entomol Appl Sci Lett*, 6 (1): 22-27.
- Parra, T., de Arriba, G., Arribas, I., Perez de Lema, G., Rodriguez-Puyol, D. and Rodriguez-Puyol, M. (1998). Cyclosporine A nephrotoxicity: role of thromboxane and reactive oxygen species. *J. Lab. Clin. Med.*, 131, 63-70.
- Pereira, C., Barros, L and Ferreira, I.C. (2016). Extraction, identification, fractionation and isolation of phenolic compounds in plants with hepatoprotective effects. *J. Sci. Food Agric.*, 96:1068-84.
- Sembulingam, K. and Sembulingam, P. (2010). *Essentials of Medicinal Physiology*, 5th Ed., New Delhi, India: Jaypee Brothers Medical Publishers (P) Ltd.
- Siddhuraju, P. and K. Becker, (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro climatic origins of drumstick tree (*Moringaoleifera* Lam). *Journal of Agricultural Food Chemistry*, 54: 2144-55.
- Tabet, A. Boukhari, A. & Nouidjem, Y. 2018. Phenolic content, HPLC analysis and Antioxidant activity extract from *Tamarix articulata*. *J Adv Pharm Edu Res*; 8(4):1-8.
- Uboh F.E, Saviour U, Eve M and Eyong E.U. (2014). Effect of *costusafer* Leaves' juice on nitrocellular thinner induced nephrotoxicity in rats. *Res J of Environ Toxicol .*, 8(1):37-45.
- Ukpai, C.F., Agbafor, K.N., Ndukwu, O.K., Agwu, A and Nwachukwu, S.N. (2012). Phytochemical composition of *costusafere* extract and its alleviation of carbon tetrachloride -induced hepatic oxidative stress and toxicity. *International Journal of Modern Botany.*, 2(5):120-6.
- Uy, I.A. Dapar, M.L.G. Aranas, T.A. Anthony R, R. Mindo, Cabrido, M.C.K. Torres, M.A. et al. (2019). Qualitative Assessment of the Antimicrobial, Antioxidant, Phytochemical Properties of the Ethanolic Extracts of the Roots of *Cocos nucifera* L. *Pharmacophore*, 10(2), 63-75.
- Wongmekiat, O., Gomonchareonsiri, S. and Thamprasert, K. (2011). Caffeic acid phenethyl ester protects against oxidative stress-related renal dysfunction in rats treated with cyclosporin A. *FundamClinPharmacol.*, 25:619-26.