

Effects of leucine-enkephalin on alpha-naphthyl acetate esterase and myeloperoxidase activity in haemolymph of the scallop *Chlamys farreri*

Dongwu Liu

Received: 23 September 2008 / Received in revised form: 6 October 2008, Accepted: 7 October 2008 Published online: 18 October 2008
© Sevas Educational Society 2008

Abstract

The study was designed to investigate the effects of leucine-enkephalin (L-ENK) on the activity of alpha-naphthyl acetate esterase (ANAE) and myeloperoxidase (MPO) in haemolymph of the scallop *Chlamys farreri*. Our results indicate that MPO activity with DAB and DEA substrate solution was induced when the concentration of L-ENK was 5 and 50 µg/ml, and the intracellular and extracellular MPO activity with two substrate solutions was highest when the concentration of L-ENK was 50 µg/ml. However, both the intracellular and extracellular MPO activity of 1 µg/ml group was significantly lower than that of control group. It was also found that the intracellular and extracellular ANAE activity was induced by L-ENK. The supernatant ANAE activity was highest when the concentration of L-ENK was 50 µg/ml, but the resultant HLS ANAE activity was highest when the concentration of L-ENK was 5 µg/ml. Overall, the data suggests an involvement of L-ENK in the regulation of the defense systems of the scallop *Chlamys farreri*. L-ENK may function as messengers in the bidirectional interaction between the neuroendocrine and the immune systems.

Keywords: *Chlamys farreri*; Leucine-enkephalin; Alpha-naphthyl acetate esterase; Myeloperoxidase; Haemolymph

Introduction

Phagocytosis is an important function of invertebrate immunocytes (Ottaviani et al. 1992). In bivalve species, phagocytic cells can be activated by foreign particles or organisms and release oxidative chemicals. This response is often referred to as an "oxidative burst". The initial metabolite of the "oxidative burst" is superoxide anion. Furthermore, superoxide anion can be dismutated

Dongwu Liu,

Analysis and Testing Center, Shandong University of Technology,
255049, Zibo, Shandong, P.R. China

Tel: 086-05332781987, Fax: 086-05332786781
E-mail: liudongwu@sdut.edu.cn

to hydrogen peroxide (H₂O₂), and be converted to other toxic reactive oxygen species (ROS). Hydrogen peroxide also interacts with myeloperoxidase (MPO) and halide to produce hypochlorous acid (HOCl), which is an even more potent microbicidal compound. In the "oxidative burst", all of metabolites play an important part in phagocyte-mediated killing of microorganisms (Adema et al. 1991; Volety and Chu 1995; Ordas et al. 2000).

It has been reported that myeloperoxidase (MPO) is present in some invertebrate phagocytes (Nakamura et al. 1985). The cellular immunity level is also related with Alpha-naphthyl acetate esterase (ANAE). ANAE is a lysosomal enzyme (Miyasaka et al. 1983; Mueller et al. 1975; Zicca et al. 1981), which has been considered to be a marker of T lymphocytes in peripheral blood and tissue sections (Beya et al. 1986; Miyasaka et al. 1983). It has been found that ANAE is present in the amebocytes of the fresh water snails, and the amebocyte enzyme histochemistry resembled that in human granuloma macrophages (McKerrow et al. 1985).

Mammalian-like opioid peptides have been found in invertebrates (Salzet and Stefano 1997; Breslin et al. 1993; Isaac et al. 1998; Stefano et al. 1998). It has been found that some opioid peptides function as endogenous messengers of the immune system, and participate in the regulation of the immune response (Hughes 1991; Stefano et al. 1995; 1990; 1993; 1991; Osman et al. 2003). Since the roles played by endogenous opioid peptides closely parallel those in vertebrates (Leung and Stefano 1987; Scharrer 1987; Stefano et al. 1989), the question arises whether the endogenous opioids affect the immune response of the scallop *Chlamys farreri*. The aims of the present study were to investigate the effects of leucine-enkephalin (L-ENK) on the activity of ANAE and MPO in the hemolymph of the scallop *Chlamys farreri*.

Material and methods

Experimental Animals

A batch of apparently healthy scallop *Chlamys farreri* was purchased from a commercial farm in Yantai, China. The scallops were acclimated in aerated seawater in 120 litre glass containers held at 21°C for at least three days prior to each experiment. Stocking densities were generally maintained at 20 molluscs per container.

Preparation of Haemocytes and Haemocyte Lysate Supernatant

A haemolymph sample about 10 ml was drawn from the posterior adductor muscle of 10 molluscs with a 25-gauge hypodermic needle. L-ENK (Sigma Co, USA) was added into the haemolymph at three different concentrations (1, 5, and 50 µg/ml), and the control group was not received any L-ENK (0 µg/ml). Following incubation at room temperature for 1 h, the haemolymph were centrifuged at 3000 rpm for 10 min. According to a protocol described by Sung et al (2004), the supernatants were collected in order to determine the extracellular enzymatic activity, and the cell pellets were used in preparing the haemocyte lysate supernatant (HLS) which was used to determine the intracellular enzymatic activity. The haemocyte lysate supernatant (HLS) was prepared according to procedures modified by Sung et al (1998). Briefly, cell pellets were resuspended in diluted water, and then homogenized with a sonicator equipped with a microtip and centrifuged at 6000 rpm for 10 min at 4°C.

Assay of alpha-naphthyl acetate esterase (ANAE) and myeloperoxidase (MPO) activity

The MPO activity in the haemocyte lysate supernatant (HLS) and supernatant was separately determined with 3, 3'-diaminobenzidine (DAB, Sigma Co, USA) and diethanolamine (DEA, Sigma Co, USA) substrate solution according to procedures described by Coles et al (1994) and Schlenk D et al (1991). The ANAE activity in the resultant HLS and supernatant was determined with alpha-naphthyl acetic acid (Sigma Co, USA) substrate solution.

Statistics

Data are expressed as mean ± SD, and statistical analysis was performed by One-way analysis of variance (ANOVA) followed with comparison test. $P < 0.05$ were considered significant and $P < 0.01$ were considered very significant. The protein concentration of the resultant HLS and supernatant was determined according to procedures described by Lowry et al (1951), using bovine serum albumin as a standard.




Figure 1: Extracellular and intracellular ANAE activity after haemolymph were added into different concentrations of L-ENK (n=6, 6 different measurements from the same extract). ^a $P < 0.01$ compared with control group, ^b $P < 0.01$ compared with 1 µg/ml group. The error bars represent the SD.

Results

The influence of L-ENK on ANAE activity with alpha-naphthyl acetic acid substrate solution

The results showed that the resultant HLS and supernatant ANAE activity of 1, 5 and 50 µg/ml group was very significantly higher than that of control group ($P < 0.01$) (Fig.1.A, B). However, there was no significant difference between ANAE activity of 1, 5 and 50 µg/ml group (Fig.1.A). The resultant HLS ANAE activity

was highest at the concentration of 5 µg/ml, which was 0.98 ± 0.02 (U) (Fig.1.B). Moreover, intracellular ANAE activity of 5 µg/ml group was very significantly higher than that of 1 µg/ml group ($P < 0.01$) (Fig.1.B).

The influence of L-ENK on MPO activity with DEA substrate solution

Using DEA as substrate solution, the supernatant MPO activity of 5 and 50 µg/ml group was not significantly higher than that of control group ($P \geq 0.05$), and the supernatant MPO activity of 1 µg/ml group was not significantly lower than that of control group ($P \geq 0.05$) (Fig.2.A). The resultant HLS MPO activity of 5 and 50 µg/ml group was very significantly higher than that of control group ($P < 0.01$), but the resultant HLS MPO activity of 1 µg/ml group was very sign-




Figure 2: Extracellular and intracellular MPO activity with DEA substrate solution after haemolymph were added into different concentrations of L-ENK (n=6, 6 different measurements from the same extract). ^a $P < 0.01$ compared with control group, ^b $P < 0.01$ compared with 1 µg/ml group; ^c $P < 0.01$ compared with 5 µg/ml group. The error bars represent the SD.

ificantly lower than that of control group ($P < 0.01$) (Fig.2.B). Both the resultant HLS and supernatant MPO activity was highest at the concentration of 50 µg/ml, which was 6.09 ± 1.72 (U) and 464.99 ± 53.69 (U), respectively (Fig.2.A, B). However, intracellular and extracellular MPO activity of 5 µg/ml group was very significantly higher than that of 1 µg/ml group ($P < 0.01$), and MPO activity of 50 µg/ml group was very significantly higher than that of 5 µg/ml group ($P < 0.01$) (Fig.2.A, B).

The influence of L-ENK on MPO activity with DAB substrate solution

We also examined intracellular and extracellular MPO activity using DAB as substrate solution. The results showed that both the resultant HLS and supernatant MPO activity of 5 and 50 µg/ml group was very significantly higher than that of control group ($P < 0.01$). However, both the resultant HLS and supernatant MPO activity of 1 µg/ml group was very significantly lower than that of




Figure 3: Extracellular and intracellular MPO activity with DAB substrate solution after haemolymph were added into different concentrations of L-ENK (n=6, 6 different measurements from the same extract). ^a $P < 0.01$ compared with control group, ^b $P < 0.01$ compared with 1 µg/ml group; ^c $P < 0.01$ compared with 5 µg/ml group. The error bars represent the SD.

control group (Fig.3.A, B). Both the resultant HLS and supernatant MPO activity was highest at the concentration of 50 µg/ml, which was 20.95±1.33 (U) and 475.32±17.90 (U), respectively (Fig.3.A, B). Moreover, intracellular and extracellular MPO activity of 5 µg/ml group was very significantly higher than that of 1 µg/ml group (P<0.01), and MPO activity of 50 µg/ml group was very significantly higher than that of 5 µg/ml group (P<0.01) (Fig.3.A, B).

Discussion

It has been found that the nervous and immune systems of invertebrates can exchange information through opioid peptides (Stefano et al. 1996; 1998; 1991). In our previous research, L-ENK participates in the regulation of nitric oxide synthase and catalase activity in the haemolymph of *Chlamys farreri* (Liu 2008; Liu and sun 2008). The present studies demonstrated that the MPO activity with DAB and DEA substrate solution was induced when the concentration of L-ENK was 5 and 50 µg/ml. However, the MPO activity was decreased when the concentration of L-ENK was 1 µg/ml. Both the intracellular and extracellular MPO activity with two substrate solutions was highest when the concentration of L-ENK was 50 µg/ml. The intracellular and extracellular ANAE activity was also induced by L-ENK. The intracellular ANAE activity was highest when the concentration of L-ENK was 5 µg/ml.

In phorbol myristate acetate-stimulated rat peritoneal macrophages, M-ENK increased H₂O₂ and NO production mainly through δ1 opioid receptor (Vujic et al. 2004). The haemocytes of invertebrate contain the µ3 subtype opioid receptor, which mediates the inhibition of cytokine-induced activation and chemotaxis by morphine and other opiates (Makman et al. 1996). In our previous experiments, we have found delta receptors and L-ENK on haemocytes of the scallop *Chlamys farreri* (Liu 2008). Thus we surmise that L-ENK may affect the activity of ANAE and MPO by binding with different opioid receptors on haemocytes. At 1 µg/ml L-ENK, L-ENK maybe binded with mu opioid receptors and MPO activity was inhibited. Although intracellular ANAE activity was highest when the concentration of L-ENK was 5 µg/ml, there is not significant difference with 50 µg/ml group. However, it is a complex network between the nervous and immune systems. The reason why different concentration of L-ENK has different function on the defense systems of the scallop *Chlamys farreri* needs to be further researched.

In summary, our results indicate that L-ENK participates in the regulation of the defense systems of the scallop *Chlamys farreri*. L-ENK may function as messengers in the bidirectional interaction between the neuroendocrine and the immune systems.

References

Adema CM, Van der Knaap WPW, Siminia T (1991) Molluscan haemocyte-mediated cytotoxicity: the role of reactive oxygen intermediates. *Rev Aquat Sci* 4:201-223

Beya MF, Miyasaka M, Dudler L et al (1986) Studies on the differentiation of T lymphocytes in sheep: 1. Two monoclonal antibodies that recognize all ovine T lymphocytes. *Immunol* 57:115-121

Breslin MB, Lindberg I, Benjannet S et al (1993) Differential processing of proenkephalin by prohormone convertases 1(3) and 2 and furin. *J Biol Chem* 268:27084-27093

Coles JA, Pipe RK (1994) Phenoloxidase activity in the haemolymph and hemocytes of the marine mussel *Mytilus edulis*. *Fish Shellfish Immunol* 4:337-352

Hughes TKJr, Smith EM, Barnett JA et al (1991) Lipopolysaccharide and opioids activate distinct populations of *Mytilus edulis* immunocytes. *Cell Tissue Res* 264:317-320

Isaac R, Schoofs L, Williams TA et al (1998) A novel peptide-processing activity of insect peptidyl-dipeptidase A (angiotensin I-converting enzyme): the hydrolysis of lysyl-arginine and arginyl-arginine from the C-terminus of an insect prohormone peptide. *Biochem J* 330:61-65

Leung MK, Stefano GB (1987) Comparative neurobiology of opioids in invertebrates with special attention to senescent alterations. *Prog Neurobiol* 28:131-159

Liu DW (2008) Effects of Leucine-enkephalin on catalase activity and glutathione level in haemolymph of the scallop *Chlamys farreri*. *J Pep Res and Therapeutics* 14:16-20

Liu DW, Sun HS (2008) Immunohistological detection of leucine-enkephalin and influence of leucine-enkephalin on the nitric oxide synthase activity of the scallop *Chlamys farreri*. *Mar Freshw Behav Physiol* 2:147-154

Lowry OH, Rosebroung NJ, Farr AL et al (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275

Makman MH, Dobrenis K, Downie S et al (1996) Presence of opiate alkaloid-selective mu3 receptors in cultured astrocytes and in brain and retina. *Adv Exp Med Biol* 402:23-28

McKerrow JH, Jeong KH, Beckstead JH (1985) Enzyme histochemical comparison of biomphalaria glabrata amebocytes with human granuloma macrophages. *J Leukoc Biol* 37:341-347

Miyasaka M, Heron I, Dudler L et al (1983) Studies on the differentiation of T lymphocytes in the sheep. I. Recognition of sheep T lymphocyte differentiation antigen by a monoclonal antibody T-80. *Immunol* 49:545-552

Mueller J, Brun DEL REG, Buerki H et al (1975) Nonspecific acid esterase activity: a criterion for differentiation of T and B lymphocytes in Mouse lymph nodes. *Eur J Immunol* 5:270-274

Nakamura M, Mori K, Inooka S et al (1985) In vitro production of hydrogen peroxide by the amoebocytes of the scallop, *Patinopecten yessoensis* (Jay). *Dev Comp Immunol* 9:407-417

Ordas MC, Novoa B, Figueras A (2000) Modulation of the chemiluminescence response of Mediterranean mussel (*Mytilus galloprovincialis*) haemocytes. *Fish Shellfish Immunol* 10:611-622

Osman AM, Gomma M, Saad AH (2003) A possible role for an enkephalinergic system in the internal defense mechanism of *Biomphalaria alexandrina* exposed to *Schistosoma mansoni*. *J Egypt Soc Parasitol* 33:841-861

Ottaviani E, Franchini A, Fontanili P (1992) The presence of immunoreactive vertebrate bioactive peptide substances in hemocytes of the freshwater snail *Viviparus ater* (Gastropoda, Prosobranchia). *Cell Mol Neurobiol* 12:455-462

Salzet M, Stefano GB (1997) Invertebrate proenkephalin: delta opioid binding sites in leech ganglia and immunocytes. *Brain Res* 768:224-232

Scharrer B (1987) Insects as models in neuroendocrine research. *Annu Rev Entomol* 32:1-16

Scharrer B (1987) Neurosecretion: beginnings and new directions in neuropeptide research. *Annu Rev Neurosci* 10:1-17

Schlenk D, Martinez PG, Livingstone DK (1991) Studies on myeloperoxidase activity in the common mussel, *Mytilus edulis* L. *Comp Biochem Physiol* 99:63-68

Stefano GB, Cadet P, Dokun A et al (1990) A neuroimmunoregulatory-like mechanism responding to stress in the marine bivalve *Mytilus edulis*. *Brain Behav Immun.* 4:323-329

- Stefano GB, Leung MK, Bilfinger TV et al (1995) Effect of prolonged exposure to morphine on responsiveness of human and invertebrate immunocytes to stimulatory molecules. *J Neuroimmunol* 63:175-181
- Stefano GB, Leung MK, Zhao X et al (1989) Evidence for the involvement of opioid neuropeptides in the adherence and migration of immunocompetent invertebrate hemocytes. *Proc Natl Acad Sci USA* 86:626-630
- Stefano GB, Salzet B, Fricchione GL (1998) Enkephalin and opioid peptide association in invertebrates and vertebrates: immune activation and pain. *Immunol Today* 19:265-268
- Stefano GB, Salzet-Raveillon B, Salzet M (1998) *Mytilus edulis* hemolymph contain prodynorphin. *Immunol Lett* 63:33-39
- Stefano GB, Scharrer B, Smith EM et al (1996) Opioid and opiate immunoregulatory processes. *Crit Rev Immunol* 16:109-144
- Stefano GB, Shipp MA, Scharrer B (1991) A possible immunoregulatory function for [Met]-enkephalin-Arg6-Phe7 involving human and invertebrate granulocytes. *J Neuroimmunol* 31:97-103
- Stefano GB, Smith EM, Cadet P et al (1993) HIV gp120 alteration of DAMA and IL-1 alpha induced chemotactic responses in human and invertebrate immunocytes. *J Neuroimmunol* 43:177-184
- Stefano GB, Smith EM, Hughes TK (1991) Opioid induction of immunoreactive interleukin-1 in *Mytilus edulis* and human immunocytes: an interleukin-1-like substance in invertebrate neural tissue. *J Neuroimmunol* 32:29-34
- Sung HH, Chang HJ, Her CH et al (1998) Phenoloxidase activity of hemocytes derived from *Penaeus monodon* and *Macrobrachium rosenbergii*. *J Invertebr Pathol* 71:26-33
- Sung HH, Huang YT, Hsiao LT (2004) Phenoloxidase activity of *Macrobrachium rosenbergii* after challenge with two pathogens: *Aeromonas veronii* and *Lactococcus garvieae*. *Fish Pathol* 39:1-8
- Volety AK, Chu FE (1995) Suppression of chemiluminescence of eastern oyster (*Crassostrea virginica*) haemocytes by the protozoan parasite *Perkinsus marinus*. *Dev Comp Immunol* 19:135-142
- Vujic V, Stanojevic S, Dimitrijevic M (2004) Methionine-enkephalin stimulates hydrogen peroxide and nitric oxide production in rat peritoneal macrophages: interaction of mu, delta and kappa opioid receptors. *Neuroimmunomodulation* 11:392-403
- Zicca A, Leprini A, Codoni A et al (1981) Ultrastructural localization of alpha naphthyl acid esterase in human TM lymphocytes. *Am J Pathol* 105:40-46