# Geographical Distribution Map and Epidemiological Pattern of Animal Bite in the North of Iran

# Seyed Akbar Hosseini, Mohamad Reza Vafaeenasab\*, Javad Rafinejad, Ali Almodaresi, Abbas Ali Dehghani Tafti, Masoud Mirzaei, Ahmad Ali Hanafi-Bojd

Received: 01 July 2019 / Received in revised form: 22 November 2019, Accepted: 27 November 2019, Published online: 27 December 2019 © Biochemical Technology Society 2014-2019

© Sevas Educational Society 2008

# Abstract

Introduction: Animal bite is a public health problem in Iran and in the world, and despite the great progress in preventing and treating these cases, it has had high economic, social, and psychological consequences. This study aimed to investigate the epidemiological and geographical status of animal bites in Mazandaran province in 2007-2016. Methods: This was a retrospective and longitudinal study conducted in 2018. The population of the study was all animal bitten people in the geographical area of Mazandaran province in 2007-2016. The data collection tool was a checklist based on the documentation available at the animal bite registration offices in disease prevention health centers. Frequency and

#### Seyed Akbar Hosseini

Department of Health in Disasters and Emergencies, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran, hosseini.ph@gmail.com

#### Mohamad Reza Vafaeenasab\*

Yazd Cardiovascular Research Centre, Shahid Sadoughi University of Medical Sciences, Yazd, Iran, rzvafaee@yahoo.com

#### Javad Rafinejad

Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, jrafinejad@tums.ac.ir

#### Ali Almodaresi

Engineering College, GIS & RS Department Yazd Branch, Islamic Azad University, Yazd, Iran, almodaresi@iauyazd.ac.ir

#### Abbas Ali Dehghani Tafti

Department of Health in Disasters and Emergencies, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran, aadtafti@ssu.ac.ir

#### Masoud Mirzaei

Department of Epidemiology & Biostatistics, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran, mmirzaei@ssu.ac.ir

#### Ahmad Ali Hanafi-Bojd

Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, aahanafibojd@tums.ac.ir percentage descriptive statistics and SPSS software were used for data analysis. Arc GIS software was also used to draw the geographical map. Results: A total of 32469 cases of the animal bite were reported in 2007-2016 in Mazandaran province. The prevalence of the disease was 45.2% in the urban area and 54.8% in the rural area. Most cases were in 2015 with 7621 patients and the least in 2013 with 3305 patients. Among the counties of Mazandaran province, Sari was the first with 4623 cases of the animal bite. Conclusion: Given the high incidence of animal bite in Mazandaran province, it is necessary to provide appropriate intervention programs for prevention and reduction of the animal bite, especially in summer and spring and rural population and emphasize on immunization of domestic dogs.

Key words: Animal Bite, Rabies, Geographical Distribution, ArcGIS, Mazandaran

## Introduction

One of the issues that threaten people's health is an animal bite. An animal bite comes from a scar from the tooth or claw of a pet or wild animal which causes the burden of disease and attenuation. (Bögel and Motschwiller, 1986) The animal bite has always been a major health threat in Iran and other countries around the world. Despite widespread advances in prevention and treatment, animal bites are still increasing and a lot of money is annually spent on vaccination and antibody serum. (Jaindl et al., 2012)

According to the World Health Organization (WHO) estimation, over ten million people in the world are treated for bites each year. (Rezaeinasab et al., 2007) Based on the statistics of disease control and prevention center, nearly 4.5 million people are bitten each year in the United States. (CDC. Dog Bites, 2013) In Iran, in 2013, there were about 150,000 cases of animal bites and 4 cases of rabies. (Goya, 2014) Animal bites are important threats to human health because some of the subsequent infections, such as rabies, are highly fatal so that the highest fatality rate among infectious diseases is rabies. (Willoughby et al., 2005) Rabies is a viral disease transmitted from warm-blooded animals to humans and causes death by creating acute and lethal encephalitis in humans and other mammals. (Madhusudana, 2005) According to the WHO reports, annually, 10 million people receive anti-rabies treatment to prevent the disease, and about 33,000 deaths are caused by rabies worldwide, most of which occur in Asia and Africa. Iran is one of the countries in the world with rabies in both wild and domestic animals. Rabies is still one of the health and economic

problems in Iran and almost all provinces are more or less involved with the disease. (Mazaheri et al., 2010)

According to the Pasteur Institute of Iran, in 2009, 131413 animal bites were reported from all over Iran treated with anti-rabies and 7 cases died from the disease. (Fayaz 2008) Given the increasing trend of stray dog populations and the growing statistics of rabies bites and dispersal in many provinces of the country, it is required to pay more attention to disease control and research in various aspects. (Zeynali et al., 1999)

Rabies is of great importance due to its fatality, the increasing incidence of animal bites in humans, damage to livestock, and economic losses. (Bokaee et al., 2009) In addition to the financial costs of preventing and treating animal bites, the psychological and social consequences of bites, and the remaining scars can also affect people's lives. (Kahn et al., 2003) Therefore, by examining the epidemiology of animal bites in a community and knowing how people are exposed, the number of animal bite cases in emergencies as well as the economic burden thereof can be reduced. (Rahmani et al., 2014; Manouchehrifar et al., 2013) Due to non-compliance with the dog keeping laws and rabies virus in warm-blooded animals in the area, which is mainly transmitted by biting, there is a high cost to vaccinate bitten people annually. Therefore, analyzing the available data can be a useful guide for planning health education and reducing the burden of this disease in the health system of the country. (Fayaz et al., 2011; Mazaheri et al., 2010)

Hence, in this study, the epidemiological and geographical status of bites was investigated. Given that most cases are found in the Caspian Sea, northeast, and southwestern regions of the country, (Jafari-Khounigh et al., 2015) this study was conducted in Mazandaran province.

# Methods

60

The present research was a retrospective-longitudinal and routine data-based study conducted in 2018. The population of the study was all animal bitten people in the geographical area of Mazandaran province in 2007-2016. According to disease control centers, an animal bitten person is defined as a person who has referred to health centers for animal bites and fear of rabies for vaccination and serum therapy. Therefore, this definition was considered as a criterion of the animal bite. Other inclusion criteria included residence in urban and rural areas of Mazandaran province and completeness of patient's information. The data collection tool was a checklist based on the documentation available at the animal bite registration offices in disease prevention health centers. The studied characteristics included age, sex, number of bites, type of residence, the month of bite, year of bite, and city. To conduct this study, the required data were extracted from animal bite registration offices at Mazandaran and Babol health centers.

The researchers have adhered to the Helsinki Ethical principles in all cases so that all patients' information was kept confidential. The information was coded without mentioning patients' personal information such as name, surname, etc. Frequency and percentage descriptive statistics and independent t-test and one-way ANOVA were used to analyze the data using SPSS software. ArcGIS software was also used to draw the geographical map.

# Results

Based on the results of animal bite cases, a total of 32469 cases were reported in Mazandaran province in 2007-2016. The prevalence of the disease was 45.2% in the urban area and 54.8% in the rural area. Most cases were in 2015 with 7621 patients and the least in 2013 with 3305 patients (Chart 1). Among the counties of Mazandaran province, Sari was the first with 4623 cases of animal bites (Table 1). No animal bites were recorded during the study period in the northern Savadkoh, Simorgh, and Kalardasht.

Table 1: Absolute frequency distribution of animal bite cases in Mazandaran province based on region, the season of disease, and residence during 2007-2016

Months									Residence						
February- March	January- February	December- January	November- December	October- November	September- October	August- September	July-August	June-July	May-June	April-May	March- April	Urban area	Rural area	Total of animal bites	Cities
150	225	226	227	319	320	375	374	285	343	317	317	1478	2000	3478	Amol
91	83	67	51	55	74	97	89	71	77	79	97	203	728	931	Babol
75	104	76	99	166	173	194	189	176	179	148	176	1080	676	1756	Babolsar
153	164	190	188	283	271	320	324	306	288	302	259	1780	1268	3048	Behshahr
110	179	188	177	205	288	312	286	248	295	220	254	1120	1642	2762	Tonekabon
45	70	77	57	79	92	119	101	99	95	112	81	328	699	1027	Jooybar
90	119	114	103	148	166	201	172	176	168	183	175	874	939	1813	Chaloos

66 86 93 87 124 121 116 148 149 122 119 148 920 454 1374 Ramsar   231 344 359 349 420 424 479 433 326 425 346 387 1793 2830 4623 Sari   39 51 38 55 40 60 72 78 82 61 50 69 277 418 695 savadkooh   22 26 15 24 24 58 51 51 59 73 56 78 294 243 537 Abbasabad   14 22 21 27 39 53 67 57 53 64 53 65 373 162 535 Freydoonkenar   163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86
231 344 359 349 420 424 479 433 326 425 346 387 1793 2830 4623 Sari   39 51 38 55 40 60 72 78 82 61 50 69 277 418 695 savadkooh   22 26 15 24 24 58 51 51 59 73 56 78 294 243 537 Abbasabad   14 22 21 27 39 53 67 57 53 64 53 65 373 162 535 Freydoonkenar   163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98
39 51 38 55 40 60 72 78 82 61 50 69 277 418 695 savadkooh   22 26 15 24 24 58 51 51 59 73 56 78 294 243 537 Abbasabad   14 22 21 27 39 53 67 57 53 64 53 65 373 162 535 Freydoonkenar   163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20
22 26 15 24 24 58 51 51 59 73 56 78 294 243 537 Abbasabad   14 22 21 27 39 53 67 57 53 64 53 65 373 162 535 Freydoonkenar   163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20 22 40 44 43 54 42 17 53 291 344 Miyandorood   68 97 114 109 151<
14 22 21 27 39 53 67 57 53 64 53 65 373 162 535 Freydoonkenar   163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20 22 40 44 43 54 42 17 53 291 344 Miyandorood   68 97 114 109 151 161 180 157 149 167 147 156 401 1255 1656 Neka
163 191 173 202 247 302 346 281 266 292 291 259 1518 1495 3013 Ghemshahr   66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20 22 40 44 43 54 42 17 53 291 344 Miyandorood   68 97 114 109 151 161 180 157 149 167 147 156 401 1255 1656 Neka
66 67 78 81 86 95 124 100 113 103 105 100 537 581 1118 Galoogah   58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20 22 40 44 43 54 42 17 53 291 344 Miyandorood   68 97 114 109 151 161 180 157 149 167 147 156 401 1255 1656 Neka
58 64 71 59 98 100 144 135 117 124 95 112 526 651 1177 Mahmoodabad   18 16 17 11 20 22 40 44 43 54 42 17 53 291 344 Miyandorood   68 97 114 109 151 161 180 157 149 167 147 156 401 1255 1656 Neka
18   16   17   11   20   22   40   44   43   54   42   17   53   291   344   Miyandorood     68   97   114   109   151   161   180   157   149   167   147   156   401   1255   1656   Neka
68 97 114 109 151 161 180 157 149 167 147 156 401 1255 1656 Neka
00 77 114 107 101 100 107 147 100 401 1255 1050 100 100 100 100 100 100 100 100
54   72   76   74   100   127   163   143   131   104   125   164   586   744   1330   Noor
61   63   74   75   95   130   135   139   124   131   115   109   535   717   1252   Noshahr
1574   2043   2067   2055   2699   3037   3535   3301   2973   3165   2905   3023   14676   17793   32469   Total

Based on the results of two independent t-tests and one-way ANOVA, there was a significant relationship between season and animal bites. However, there was no significant relationship between residence and animal bites (Table 2).

0.04 478.6 Spring 516.3 Summer 410.1 Fall 299.1 Winter Season

Table 2: The relationship between animal bites in Mazandaran province and season and type of residence during 2007-2016

Significance level	Mean bite	Туре	Variable	
0.41	936.4	Rural	Type of	
0.41	772.4	Urban	residence	



Chart 1: Frequency distribution of animal bites in Mazandaran province in 2007-2016



Figure 1: Distribution map of animal bites in Mazandaran province based on the studied years.



Figure 2: Distribution map of animal bites in cities of Mazandaran province during 2007-2016.

#### Discussion

According to the results, there was an average of 5004 animal bites per year in Mazandaran province. The results of Naghibi et al. in Mazandaran province also showed that 25869 animal bites were reported in Mazandaran province in 2008-2012 and the average incidence of bites in the five-year period was 203 cases per 100,000. (Naghibi et al., 2014) A comparison of the incidence of animal bites in the study period shows that the incidence rate is increasing each year. This result can be related to public awareness of the consequences of rabies and their referral to rabies prevention centers, improving the registration and reporting system for animal bites, tourism development, and the presence of families in rural, forest, and recreational areas.

The prevalence of the disease was 45.2% in the urban area and 54.8% in the rural area. The results of the study by Zohrevandi et al. (2014) entitled "Epidemiology of animal bites in Rasht city" also showed that 61.3% of animal bites were in rural areas. The study of Babaeeian-Moghaddam et al. (2015) in Quchan city showed that animal bites were more in rural areas. Similar studies conducted by Bahonar et al. in Ilam, Majidpour, et al. in Ardebil, as well as Amiri and Khosravi in Shahrood (2009), have also found that most of the animal bites were in rural areas. Given that in rural areas people are more likely to be in contact with cattle and dogs, it can be inferred that the incidence of bites is higher in these areas.

Most bites occurred in summer and after that in spring. In the study of Zohrevandi et al. (2014), most bites were in spring and summer. The results of Dadipour and colleagues in the North of Iran (Dadypour et al., 2009) and Charkazi et al. in Golestan province (Charkazi et al., 2013) were in line with the results of the present study. Most of the animal bites in the study of Babaeeian-Moghaddam (2015) were in spring and May-June. In the study of Bahonar et al. (2008) most bites were also in spring. (Bahonar et al., 2008) Majidpour's study showed that animal bites in summer were more common than other seasons. The cause of the increase in animal bites in summer can be attributed to the onset of the breeding season. More frequency, this is justified in two ways in spring and summer. First, the risk of being bitten in Mazandaran province increases due to holidays and an increased number of passengers. On the other hand, depending on the type of farmers' activity in spring and summer, the risk of being bitten is high, especially in this occupational group.

Generally, the findings of this study suggest the need to control animals, such as dogs and cats through appropriate training in teenagers and adolescents. Moreover, comprehensive educational content should be designed to teach how to treat dogs and provide targeted and principled training using educational theories. Furthermore, adopting proper laws and forcing dog owners to use collars can also reduce animal bites. Finally, it is suggested that by repeating the present study elsewhere in the country and determining the bite pattern of each area, the necessary conditions for the prevention and treatment of bitten patients will be provided.

# Conclusion

The results of the study showed that the number of animal bites increased during the years studied. These results could be due to increased awareness of the dangers of animal bites and increased public access to appropriate treatment, vaccine and anti-rabies serum, and partly due to increased bites. Therefore, it makes the process of vaccine supply difficult. On the other hand, according to the Millennium Development Goals and the Principles of Health for All, one of the concerns of health officials and planners is to reduce health-related problems, such as physical disability, illness and death, mental distress, social problems, and economic losses. Therefore, the results of this study can be used for planning and policy-making by the authorities. In this study, the incidence of animal bites in different seasons and months of year and locations were determined. According to the results of this study, it seems necessary to provide educational and health intervention programs to pet owners, especially dog owners, with emphasis on animal vaccination in order to prevent and reduce animal bite cases. Moreover, the authorities, in May-June and spring, should be prepared for more bitten people referral and provide the necessary vaccine and anti-rabies serum supplies, equipment, and staff.

# References

- Amiri, M., & Khosravi, A. (2009). Animal bites epidemiology in Shahroud city.
- Babaeeian-moghaddam, M., Hashemi-nazari, S. S., & Khodakarim, S. (2015). Epidemiological study on animal bite cases and its related injury in Quchan district in2013. Safety Promotion and Injury Prevention, 3(1), 9-14.
- Bahonar, A. R., Bokaie, S., Khodaveirdi, K. H., Nikbakht Boroujeni, G. H., & Rad, M. A. (2008). A Study of Rabies and the Frequency of Animal Bites in the Province of Ilam, 1994-2004. *Iranian Journal of Epidemiology*, 4(1), 47-51.
- Bögel, K., & Motschwiller, E. (1986). Incidence of rabies and postexposure treatment in developing countries. *Bulletin of the World Health Organization*, 64(6), 883.
- Bokaee S, Fayaz A, Poor Mehdi Broojeni M, Haghdoost AA. Zolfaghari MR, & Esfandiari B. (2009). Epidemiological survey of animal bite and rabies in the Caspian Sea littoral provinces during, 1996-2006. J Faculty veterinary Med, 5(1):5-14.
- CDC. Dog Bites. (2013). Available at: http://www.cdc.gov/homeandrecreationalsafety/dogbites/index.html.
- Charkazi, A., Behnampour, N., Fathi, M., Esmaeili, A., Shahnazi, H., & Heshmati, H. (2013). Epidemiology of animal bite in Aq Qala city, northen of Iran. *Journal of education and health* promotion, 2.
- Dadypour, M., Salahi, R., & Ghezelsofla, F. (2009). Epidemiological survey of animal bites in Kalaleh district, North of Iran (2003-05). *Journal of Gorgan University of Medical Sciences*, 11(1), 76-9.
- Fayaz A. (2008). Report to WHO collaborating center for reference and research on rabies.
- Fayaz, A., Fallahian, V., Simani, S., Eslamifar, A., Mohammadian, A., Hazrati, M., ... & Elm, E. (2011). Epidemiological

characteristics of persons exposed to rabies in Tehran referred to Pasteur Institute of Iran during the years of 1993-1994 and 2008-2009. *Research in Medicine*, *35*(3), 168-173.

- Goya M. (1393). Available at http://www.behdasht.gov.ir/?siteid=1&pageid=177&newsvie w=114492.
- Jafari-Khounigh, A., Nayerpour, A., Ahari, A. M., Mahmoodi, J., Rezadokht-Mamaghani, P., & Jeddi, M. B. (2015). Epidemiology of animal bites in Azarshahr town: A crosssectional study of key determinants. *Depiction of Health*, 5(4), 21-26.
- Jaindl, M., Grünauer, J., Platzer, P., Endler, G., Thallinger, C., Leitgeb, J., & Kovar, F. M. (2012). The management of bite wounds in children—a retrospective analysis at a level I trauma centre. *Injury*, 43(12), 2117-2121.
- Kahn, A., Bauche, P., Lamoureux, J., & Dog Bites Research Team. (2003). Child victims of dog bites treated in emergency departments: a prospective survey. *European journal of pediatrics*, 162(4), 254-258.
- Madhusudana, S. N. (2005). Rabies: An ancient disease that still prevails. *Indian Journal of Medical Research*, *122*(1), 4.
- Majidpour, A., Arshi, S., SADEGHI, H., Shamshirgaran, S. M., & Habibzadeh, S. (2004). Animal bites: epidemiological considerations in Ardabil province, 2000.
- Manouchehrifar, M., Khosravi, S., Khavanin, A., & Derakhshandeh, N. (2013). Report of 267 cases of scorpion bite referring to an emergency department during one year. *Emergency*, 1(1), 24.
- Mazaheri, V. H. N. K., Holakouie Naieni, K., Simani, S., Yunesian, M., Fayaz, A., Mostafavi, E., & Biglari, P. (2010). Geographical distribution of animal bite and rabies in the

Caspian Sea littoral provinces during 2002-2007. Journal of School of Public Health and Institute of Public Health Research, 8(3), 37-46.

- Naghibi, S. A., YazdaniCharati, J., & Shojaie, J. (2014). Epidemiological characteristic of animal-bite cases in Mazandaran, 2004-2011. *Journal of Mazandaran University* of Medical Sciences, 24(117), 218-224.
- Rahmani, F., Khojasteh, S. M. B., Bakhtavar, H. E., Rahmani, F., Nia, K. S., & Faridaalaee, G. (2014). Poisonous spiders: bites, symptoms, and treatment; an educational review. *Emergency*, 2(2), 54.
- Rezaeinasab, M., Rad, I., Bahonar, A. R., Rashidi, H., Fayaz, A., Simani, S., ... & Rad, M. A. (2007). The prevalence of rabies and animal bites during 1994 to 2003 in Kerman province, southeast of Iran. *Iranian Journal of Veterinary Research*, 8(4), 343-350.
- Willoughby Jr, R. E., Tieves, K. S., Hoffman, G. M., Ghanayem, N. S., Amlie-Lefond, C. M., Schwabe, M. J., ... & Rupprecht, C. E. (2005). Survival after treatment of rabies with induction of coma. *New England Journal of Medicine*, 352(24), 2508-2514.
- World Health Organization. (2008). Human and animal rabies. Available at: Http://www.who.int/rabies / en. Accessed Jan 20, 2008.
- Zeynali, M., Fayaz, A., & Nadim, A. (1999). Animal bites and ra bies: situation in Iran. *Arch Iran Med*, *2*(3), 120-4.
- Zohrevandi, B., Asadi, P., Kasmaie, V. M., Tajik, H., & Fatemi, M. S. (2014). Epidemiologic study of animal bites in Rasht County, Guilan Province, Iran's North, 2012. *Iranian journal* of emergency medicine, 1(1), 11-15.