Analgesic Utilization Pattern in Department of Surgery, Esic-Pgimsr & Model Hospital, Rajajinegar, Bangalore, India

Farhad Marghoubi *, Malini muraleedharan Nair, NS Kodandaram

Received: 22 November 2017 / Received in revised form: 18 May 2018, Accepted: 23 May 2018, Published online: 05 September 2018 © Biochemical Technology Society 2014-2018 © Sevas Educational Society 2008

Abstract

Objective: This study aims to analyze the utilization pattern of analgesics in the department of surgery, ESIC-PGIMSR & Model Hospital, Rajajinegar, Bangalore. Methodology: This was a prospective observational study, carried out over a period of 6 months, Patients admitted under the department of surgery was enrolled using inclusion and exclusion criteria by taking / obtaining their consent. The well designed / structured data collection forms were prepared and used. The data required for the study was collected from patient case sheet & by interacting with the care givers into the data collection form. Result: The study was conducted in the in-patient wards of the department of surgery at ESIC-PGIMSR & model hospital, Rajajinagar, Bengaluru. Majority of the patients was male. The study showed that majority of patients belonged to the age group (33-45, 27.2%). Mean age of population was 45 years old. For the pain management of post-operative patients in the department of surgery four analgesics i.e. paracetamol, tramadol, diclofenac and Mefenamic acid were prescribed. The result of the study shows that paracetamol and tramadol (NON-NSAID) were the most commonly prescribed analgesics, it may be due to its lesser side effects and their effectiveness when compared to others. Majority of analgesics were prescribed, using brand name, the prescription of medications in brand name could be change to nonexclusive name, additionally this assumes a significant role in rational use of medication (proper dispensing of drugs, cost, safety and efficacy). The most widely recognized route of organization was intravenous. Conclusion: The utilization of analgesics basically relies on the severity of pain. In mild pain, single analgesics are ordinarily utilized while at least analgesics are utilized in moderate or severe pain. Paracetamol (8.9%) and diclofenac (6.1%) alone and combination of paracetamol with tramadol (44.2%) are the most common practice in various surgical wards of our hospital. In this study the most common procedures were general surgical procedures like appendectomy (17.7%) and majority of patients were given subarachnoid anesthesia (74.1%), paracetamol (67.34%) was the most common analgesic which was prescribed for patients who were given subarachnoid anesthesia.

Keywords: analgesics, utilization pattern, NSAID, Department of Surgery.

Introduction

The world health organization (WHO) in 1977 has characterized drug usage as the showcasing, distribution, prescription and utilization of medications in a general public, with special accentuation on the resulting medical, social and economic results (WHO, 2003; Ahmed, Rahman and Khan, 2012). Pain is an unpleasant sensation happening in differing degrees of severity as a result of injury, disease, or emotional disorder (Kumarasingam, Revathy and Mukherjee, 2014). Poor pain control is dishonest, clinically unsound and monetarily inefficient (Bertolini et al., 2002). The ongoing activity of including pain as fifth vital sign in health care has underlined pain assessment is similarly imperative to that of temperature, pulse, blood pressure, and respiratory rate recording. "Pain is always subjective" (Mohammed, Beegum and Perumal, 2011; Elvir-Lazo and White, 2010).

Acute painful disorders are dealt with in instantly; on the other hand, severe post-operative pain and severe visceral pain are under analyzed and under treated. Effective postoperative pain control was accomplished by proficient utilization of health resources and patient's satisfactions (Swamy, Venkatesh and Nagaraj, 2010). Analgesics are characterized as the medications that mitigate pain without blocking nerve impulse conduction or particularly modifying sensory function (Mohammed, Beegum and Perumal, 2011). A specific analgesic dose that produces fruitful pain relief in one patient may create bearable adverse effects and inadequate pain control in

Farhad Marghoubi *, Malini muraleedharan Nair

Department of Pharmacy Practice, Acharya & B.M. Reddy College of Pharmacy, Bengaluru, India.

N S Kodandaram

Department of Surgery, ESIC-PGIMSR & Model Hospital, Bengaluru, India.

*Email: farhad.pharm2010@gmail.com

someone else. Expansive assortments of analgesics are accessible in the market which may lead to the issue of irrational prescription (Kamaldeen et al., 2012).

The effective relief of pain is of the most importance to whom treating patients undergoing surgery. Pain relief has important physiological benefits, observing of pain relief is progressively turning into significant postoperative quality measure. The aim for postoperative pain management is to diminish or take out pain and discomfort with at least of side effects. Traditionally the most commonly used postoperative analgesia is opioid based, and there are lots of supportive way to reduce opioid side effects (such as nausea and ileus) and relief the pain. The effectiveness of available drugs and drug's routes of administration is important in the postoperative regimen to the needs of the individual patient (Garimella and Cellini, 2013).

The most widely recognized strategy for postoperative pain relief is the traditional utilization of intramuscular opioid injections. Effective pain relief can be acquired with newer systems, for example epidural opioids and patient controlled analgesia (PCA). A few procedures, for example, pre-emptive analgesia have not been completely assessed and others, such as infiltrating local anesthetic into wounds, may not decrease the patient's analgesic needs. Patients ought to be advised before surgery because a clarification of what to expect, may decrease their analgesic requirements (Ogboli-Nwasor, Sa'adatu and Yusufu, 2012).

Analgesics including NSAIDs and paracetamols are commonly prescribed group of drugs in clinical practice for the management of pain and inflammation. Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) are commonly available over the counter. The normal adverse drug effects with this group of medication are the gastrointestinal tract (GIT) toxicity. Along these lines, periodic assessment of medication utilization patterns should be done to empower appropriate changes in the prescription so as to increase therapeutic advantages and reduce the adverse effects (Fehintola, and Ganiyu, 2008).

The investigation of successful use of analgesic in postoperative period is a huge constituent of medical audit which helps in observing, assessing and building required modifications in the recommending practices to accomplish a rational and cost effective medical care. The upsides of effective use of analgesic in postoperative period incorporate patient solace and in this way fulfillment, earlier mobilization, quicker recovery with less probability of the development of neuropathic pain, a diminished risk of deep vein thrombosis, fewer pulmonary and cardiac complications, and decreased expense of care (Ismail, Siddiqui and Rehman, 2018). Hence this study was intended to investigate the utilization pattern of analgesics in the department of surgery.

Materials and Methods

The study was conducted in ESIC-PGIMSR & Model Hospital, Rajajinagar. A study of 6 months was designed in which planning was done from November-December 2017 followed by data collection January-March 2018 & interpretation and thesis writing was done on April 2018. Data was collected by using a self-designed data collection form for surgery, which consists of details like age, sex, lab data, disease condition, history of the illness, co-morbidities, drug therapy and other relevant information. Patients admitted under the department of surgery was enrolled using inclusion and exclusion criteria by taking / obtaining their consent. The well designed / structured data collection joins were prepared and used. The data required for the study was collected from patient case sheet & by interacting with the care givers into the data collection form. All the hospitalized cases were sorted out excluding OPD cases and 147 cases were randomly selected for the study.

Result and Discusion

In This study we prospectively observed the utilization pattern of analgesics in the surgery department. A total of 147 patients were enrolled in the study, based on the inclusion and exclusion criteria. The study was conducted for a period of 6 months in the department of surgery in ESI hospital Rajaji Negar, Bangalore.

Majority of the sample population were belonged to the age group of 18-30. We were able to observe that male patients (104, 71%) were predominant than female patients (43, 29%). This was in similarly to a studies conducted by Roshini A et al., (2015) conducted a study at suraram, Telangana, India as result total of 218 patients (103, 47%) were female and (115, 53%) were male.

In our study the mean age of the sample population was found to be 48 years. This was in similarly to a study conducted by Sen, et al., (2015) the mean age of the patients were 49 years.

Analoggies	Number of patients (N=147)	
Analgesics	Number	Average/patient
NSAIDs	52	0.3
NON NSAIDs	225	1.5
TOTAL	277	1.88

Table 1: Average Number of Analgesics Per Prescription

During this study it was observed that all patients were prescribed with analgesics (NSAIDs/Non-NSAIDs) for post-operative pain management. Commonly used NSAIDs were Mefenamic Acid and Diclofenac and Non-NSAIDs were Paracetamol and Tramadol. Out of this 4 analgesics paracetamol was the most commonly prescribed analgesic (128, 46.2%) followed by tramadol (97, 35%), diclofenac (51, 18.4%), the least used analgesics was Mefenamic acid (1, 0.4%). In this study A total of 277 analgesics were prescribed to the 147 patients The average number of analgesic per prescription was 1.88. table 1

During this study the Most Common Route of Administration of analgesics (paracetamol, tramadol, diclofenac) Was found to Intravenous (IV) i.e. (233, 83%) and (47, 17%) patients were managed with oral therapy (Paracetamol, Tramadol and Diclofenac). This is similarly to the study written by Roshini A et al., 2015, showed that in (76.6%)167 of the patients, analgesics was prescribed intravenously.

In this study analgesics was prescribed 277 times, 241 times (87%) was prescribed with trade names and (36, 13%) was prescribed with generic names. in this study (120, 81.6%) of patients were on combination analgesic therapy. combination of tramadol+ paracetamol was predominantly used (65, 44.2%) followed by, Diclofenac+ paracetamol (37, 25.2%). the findings were consisted with a study conducted by Sen, S et al; (2015) in which they observed tramadol+paracetamol as the most frequent analgesic combination followed by diclofenac+ paracetamol.

Combination of Analgesics	Frequency (number)	Frequency(percentage)
Tramadol + PCM	65	44.2
Diclofenac + PCM	37	25.2
Tramadol + PCM+ Diclofenac (Triple Drug Combination)	18	12.2
Total	120	81.63

Table 2: Utilization Pattern of combination therapy

During this study Triple drug combination was only given tramadol+ paracetamol+ diclofenac (18, 12.2%). In this study we found out that most patients were admitted for undergoing appendectomy (26,17.7%) followed by incision and drainage (14,9.5%), mesh repair (10,6. 8%).one finding was similar to the study done by Sen, S et al. (2013) showed that the most common procedure done was general like appendectomy (14.32%).

In our study Most commonly prescribed adjuvant drugs along with analgesics were PPI's like pantoprazole (139, 21.5%) followed by antibiotics. This was in similarly to a study conducted by Choudhury, et al. 92017) showed that Pantoprazole was the most commonly prescribed medication (21.55%) among the gastro protective drugs. In this study Among antibiotics ceftriaxone was the most commonly prescribed (122, 18.8%) followed by metronidazole (70,10.8%).

Type of anesthesia	Distribution between 147 patients	
Type of unestitestit	Frequency (number)	Frequency (percentage)
Subarachnoid	109	74.1
General	27	18.4
Local	5	3.4
Laryngeal mask airway (LMA)	4	2.7
Ankle block	1	0.7
Sedation	1	0.7

Table 3: Distribution of Types of Anesthesia

In our study, (109, 74.1%) patients were given Subarachnoid anesthesia, (27, 18.4%) patients were given General anesthesia, (5, 3.4%) patients were given Local anesthesia, (4, 2.7%) patients were given Laryngeal mask airway anesthesia and only 1 patient was (0.7%) given Ankle Block and 1 patient was given Sedation (0.7%) anesthesia. Table3

Table 4: Frequency of antibiotics

Antibiotics name	Distribution of drugs	
	Frequency(number)	Frequency(percentage)
Ceftriaxone	122	82.9

metronidazole	70	47.61
Cefixime	36	24.48
clindamycin	18	12.24
ciprofloxacin	12	8.16
amikacin	8	5.44
Amoxicillin + clavulanate	4	2.7
meropenem	3	2.04
cotrimoxazole	2	1.36

Antibiotics and GI drugs were the drugs given adjuvant along with analgesics post operatively. The most common antibiotic prescribed was Ceftriaxone (122, 82.9%) of patients followed by Metronidazole (70, 47.61%) patients. Among GI drugs i.e. Pantoprazole (139, 94.5%) patients coming under PPIs was the mostly prescribed followed by Ranitidine under H2 blockers was prescribed (8, 5.44%) patients for gastric irritation. Ondansetron was only the anti-emetic prescribed (105, 71.4%) patients. Table 4,5

Drugs name CATEGORY	Distribution of drugs		
	CAILGORI	Frequency(number)	Frequency(percentage)
Pantoprazole	Proton Pump Inhibitor	139	94.5
Ondansetron	Antiemetic	105	71.4
Ranitidine	H2 blocker	8	5.44

Table 5: Frequency of GI drugs

In this study we found out that Paracetamol was the most common analgesic which was prescribed for patients who were given General anesthesia (21, 14.28%), Local anesthesia (4, 2.7%) and Sub arachnoid anesthesia (99, 67.34%). Paracetamol and Tramadol were prescribed in equal ratio (1, 0.7%) in Sedation and (1, 0.7%) Ankle Block anesthesia. Tramadol was the most prescribed in Laryngeal Mask Airway anesthesia (3, 2.04%).

Conclusion

For the pain management of post-operative patients in the department of surgery ward The result of the study shows that Paracetamol and Tramadol were the most commonly prescribed analgesics, it may be due to its lesser side effects and their effectiveness when compared to others. Standard treatment guideline will also help rational use of analgesics which improves a cost effective treatment plan with the introduction of safe and effective analgesics for post-operative pain relief.

Majority of analgesics were prescribed, using brand name, the prescription of medication in brand name must be change to generic name, likewise this plays a significant role in rational utilization of medication (proper dispensing of drugs, cost, safety and viability).

The utilization of analgesics predominantly relies on the severity of pain. In mild pain, single analgesics are normally utilized while at least two analgesics are utilized in moderate or severe pain. Paracetamol and Diclofenac alone and combination of Paracetamol+Tramadol are the most common practice in various surgical wards of our hospital.

Use of anti-emetics, PPIs or H2 blockers and antibiotics in our surgical practices to avoid any post-operative nausea and vomiting (PONV), the post-operative stress bleeds and prevented infections is seen to the great extent. The investigation demonstrates that antiulcer drugs were recommended along with analgesics to reduce the significant adverse effects of NSAIDs that is gastric complications. The investigation is valuable in diminishing the irrational prescription, which help to reduce the morbidity and health care burden in the general public.

References

- Ahmed, M., Ali, N., Rahman, Z. U., & Khan, M. (2012). A study on prescribing patterns in the management of arthritis in the department of orthopaedics. Der Pharmacia Lettre, 4(1), 5-27.
- Allam Roshini A, Siva Abhishek, K Ashwini, J Manasa, Mohd Tahir Siddiqui. Prescribing patterns of analgesics in the post-operative period of surgical interventions. Indo American Journal of Pharm Research.2015, 5(07):2555-2559.
- Bertolini, G., Minelli, C., Latronico, N., Cattaneo, A., Mura, G., Melotti, R. M., & Iapichino, G. (2002). Gruppo Italiano per la Valutazione degli Interventi in Terapia I. The use of analgesic drugs in postoperative patients: the neglected problem of pain

control in intensive care units. An observational, prospective, multicenter study in 128 Italian intensive care units. Eur J Clin Pharmacol, 58, 73-77.

- Choudhury, D., Gairola, B., Roy, D., & Sikidar, P. (2017). evaluation of analgesic activity of aqueous extract of aloe Vera, in albino Wistar rats. International Journal of Pharmaceutical Sciences and Research, 8(4), 1850-1857.
- Elvir-Lazo, O.L., White, P.F. The role of multimodal analgesia in pain management after ambulatory surgery. Curr Opin Anaesthesiol. 2010; 23:697–703
- Fehintola, F A, and A A Ganiyu. "Prescriptions involving analgesic drugs at a secondary health facility in ibadan, Nigeria." Annals of Ibadan postgraduate medicine vol. 6,2 (2008): 34-8.
- Garimella, V., & Cellini, C. (2013). Postoperative pain control. Clinics in colon and rectal surgery, 26(3), 191–196. doi:10.1055/s-0033-1351138
- Ismail, S., Siddiqui, A. S., & Rehman, A. (2018). Postoperative pain management practices and their effectiveness after major gynecological surgery: An observational study in a tertiary care hospital. Journal of anaesthesiology, clinical pharmacology, 34(4), 478.
- Kamaldeen AS, Omuya LM, Muhammad AS et al. JAPS, 2012; 2: 194-198.
- Kumarasingam, T., Revathy, S., & Mukherjee, D. (2014). Drug utilization pattern of analgesics among postoperative patients in a tertiary care hospital. Pharm Lett, 6(3), 40-6.
- Mohammed TCH, Beegum IM, Perumal P. Int.J. PharmTech Res.2011,3(3):1521-29.
- Ogboli-Nwasor, E., Sa'adatu, T. S., & Yusufu, L. M. (2012). Pattern of postoperative pain management among adult surgical patients in a low-resource setting. Journal of pain research, 5, 117.
- Sen, S., & Bathini, P. (2015). Auditing analgesic use in post-operative setting in a teaching hospital. Journal of Clinical and Diagnostic Research: JCDR, 9(4), FC01.
- Sen, S., Chakraborty, R., Rekha, B., Revathi, D., Ayyanna, S. C., Hemalatha, G., ... & Sridhar, C. (2013). Anti-inflammatory, analgesic, and antioxidant activities of Pisonia aculeata: folk medicinal use to scientific approach. Pharmaceutical biology, 51(4), 426-432.
- Swamy RM, Venkatesh G, Nagaraj HK. Biomedical Research 2010; 21 (4): 401-405.
- WHO (World Health Organization). Introduction to drug utilization research, World Health Organization: Geneva, 2003. Assessed on 25-07-2013.