

Amelioration of mercuric chloride induced oxidative stress by *Hygrophila auriculata* (K.Schum) Heine via modulating the oxidant - antioxidant imbalance in rat liver

Sridhar MP, Nandakumar N, Rengarajan T, Balasubramanian MP*

Received: 01 July 2012 / Received in revised form: 15 February 2013, Accepted: 18 March 2013, Published online: 11 July 2013,
© Sevas Educational Society 2008-2013

Abstract

Medicinal plants and their phytochemical constituents are considered as paramount entity to treat various human ailments. The present study was designed to investigate the effect of ethanolic extract of *Hygrophila auriculata* against mercuric chloride induced oxidative damage in rat liver. Mercuric chloride (HgCl_2) (1mg/kg b.w, i.p) was administered three times in a week for two weeks to induce mercury toxicity. Mercury damage to the liver was evidenced from an increase in the activities of marker enzymes such as ALT, AST, ALP and ACP in serum. The level of lipid peroxidation was increased in liver tissue with concomitant decrease in both the enzymic and non-enzymic antioxidants such as superoxide dismutase, catalase, glutathione peroxidase, reduced glutathione, vitamin C and vitamin E. The glutathione metabolizing enzymes such as glutathione reductase, glucose-6-phosphatase dehydrogenase and glutathione-S-transferase were also decreased. Interestingly post treatment with ethanolic extract of *Hygrophila auriculata* (100 mg/kg b.w, p.o) for 10 days after mercury induction significantly ameliorated the mercury induced oxidative damage which was evidenced by the reduction in the marker enzymes, lipid per-oxidation and increase in the status of antioxidants and enzymes of glutathione metabolism. Thus, this remarkable hepato-protective activity of ethanolic extract of the plant might be due to the antioxidant nature, which scavenges excessive free radicals that were involved in the acceleration of per-oxidation reaction and the potent membrane protective property which ultimately render protection to the liver against HgCl_2 induced oxidative damage.

Key words: *Hygrophila auriculata*; Mercuric chloride; Antioxidant; Lipid per-oxidation

Sridhar M.P, Nandakumar N, Rengarajan T, Balasubramanian M.P*

Department of Pharmacology and Environmental Toxicology,
Dr.AL Mudhaliar Post Graduate Institute of Basic Medical Science,
University of Madras, Chennai -600 113, Tamilnadu, India

*Tel: 0091 4424 547143, 0091 4424 547147
Email: balumppet@gmail.com

Introduction

Heavy metals such as aluminum, cadmium, lead, zinc and mercury affect many tissues like liver, kidney and brain etc (Huang et al. 1996). Due to industrialization and changes in the environment during the twentieth century, humans and animals are exposed to numerous chemical form of mercury. Since mercury is ubiquitous in the environment, it is nearly impossible for most humans to avoid exposure to mercury in any of its forms on a regular basis (Fitzgerald and Clarkson 1991). All forms of mercury cause toxic effects in a number of tissues and organs depending upon the level of exposure, the duration of exposure and the route of exposure (Clarkson 1972). Since mercury is being widely used in different fields such as medical, agricultural and industrial fields, its exposure cannot be avoided by humans (Sharma et al. 2002). Heavy metals are known to produce oxidative damage in the liver tissues by enhancing peroxidation of membrane lipids (Chaurasia and Kar 1997), a deleterious process solely carried out by free radicals (Halliwell and Gutteridge 1990). Many studies have investigated possible relationship between lipid peroxidation (LPO) and cellular damage in hepatic tissues under various pathological conditions (Comporti 1985). Lewis and Willis (1962) have suggested that peroxide formation may lead to oxidative destruction of thiol groups of amino acids and proteins.

Cell membranes are targets for oxidative damage produced by xenobiotics including heavy metals (Halliwell and Gutteridge 1990). Per-oxidative decomposition of membrane lipids is catastrophic for living system. Antioxidant enzymes such as catalase, superoxide dismutase, glutathione peroxidase, glutathione-S-transferase and glutathione reductase are scavenges free radicals and lipid peroxides and deoxygenate them (Dwivedi et al. 1984). Previous reports suggest that LPO is enhanced by disturbances such as depletion of cellular antioxidants (Tan et al. 1984).

Many antioxidant compounds, occurring naturally from plant sources, have been identified as free radical scavengers (Duh 1998). Recently, interest has increased considerably in finding naturally occurring antioxidants from feed or medicinal flora to replace synthetic antioxidants, which are being restricted, due to their adverse side effects such as carcinogenicity, hepatotoxicity and nephrotoxicity (Ito et al. 1983). Natural antioxidants can protect the human body from free radicals and retard the progress of many chronic diseases and also retard the lipid oxidative rancidity in food,

cosmetics and pharmaceutical materials (Kinsella et al. 1993). In search of plant as a source of natural antioxidants, some medicinal plants and fruits have been extensively studied for their antioxidant activity and radical scavenging in the past few decades (Velioglu et al.1998). Among the various natural antioxidants phenolic compounds are reported to be active, quenching oxygen derived free radicals by donating hydrogen atom or an electron to the free radical (Yuting et al.1990). Also, phenolic compounds of plant materials have been shown to neutralize free radicals in various *in vitro* model systems (Zhang et al.1996).

Hygrophila articulata (K. Schum.) Heine (Syn) is a wild herb commonly found in moisture places on the banks of tanks, ditches and paddy fields throughout India and is one of the sources of Ayurvedic drug 'kokilaksha'. Aerial parts of the plant are used ethnobotanically for the treatment of body pain, jaundice and malaria, while the seeds are used for treatment of impotence and thus as aphrodisiac (Jain 1991). Aerial parts of *H. auriculata* have been reported to contain lupeol, stigmasterol and butelin while the seeds of the plant are reported to contain mainly fatty acids (Quasim and Dutta 1967). The plant has been shown to possess hypoglycemic activity in human subjects (Fernando et al.1989), hepatoprotective activity against paracetamol and thioacetamide intoxicification in rats (Singh and Handa 1999) and CCl₄-induced liver dysfunctions (Shailajan et al. 2005), antitumour (Mazumdar et al. 1997), anabolic and adrogenic activities (Jayatilak et al. 1976). *H. auriculata* seeds have been reported to ameliorate the activities of antioxidant enzymes such as glutathione peroxidase and catalase in hepatocarcinoma (Ahmed et al. 2001). However, there is a paucity of information regarding antioxidant properties in experimental animals using the whole part of this plant. Therefore, the present study was undertaken to investigate the effect of ethanolic extract of whole plant *H. auriculata* against mercuric chloride (HgCl₂)-induced toxicity in wistar albino rats in order to validate the ethnobotanical and clinical claims of the plant.

Materials and methods

Animals

Healthy male Wistar albino rats weighing 120±30g purchased from Tamil Nadu Veterinary and Animal Sciences University, Chennai, were used for this study. Animals were housed in polypropylene cages and were provided certified rodent pellet diet and water *ad libitum*. They were maintained at 27°C to 37°C with 12 h light and dark cycle. All animal experiments were performed in accordance with the strict guidelines prescribed by the Institutional Animal Ethical Committee (IAEC No. 07/041/05) after getting necessary approval.

Chemicals

Mercuric chloride was purchased from Sigma Chemical Company, USA. 5-5'-dithiobis-2-nitrobenzoic acid (DTNB) and 1-chloro 2, 4-dinitrobenzene (CDNB) were purchased from SISCO Research Laboratories, Chennai, India. All the other chemicals used were of analytical grade and were purchased locally.

Collection of plant material

The plant is widely distributed throughout India, Srilanka, Burma, Malaysia and Nepal. The plants were collected from Red Hills, Tamil Nadu in the month of August 2004. The plant specimen was authenticated by Dr. S. Jayaraman, Plant Anatomy Research Center, Chennai, Tamilnadu, India. A voucher specimen has been deposited at the herbarium unit of the Department of Pharmacology and Environmental Toxicology, University of Madras, Taramani, Chennai.

Preparation of plant extract

The whole plant was shade dried and coarsely powdered. The powder was then extracted with ethanol (60-70°C) using Soxhlet extractor. The extract was dried under reduced pressure using rotary flash evaporator. The percentage yield of alcoholic extract was 12% w/w. The extract was stored in refrigerator for further studies.

Experimental design

Rats were divided into 4 groups with 6 animals in each group. The experimental design was as follows: Group I rats served as controls and were treated with normal saline (0.9% NaCl). Group II rats were administered with HgCl₂ (1 mg/kg b.w., i.p.) three doses per week for two weeks. Group III rats were administered with HgCl₂ dosage as in group II and then extract of *H. auriculata* (100 mg/kg b.w, p.o) for ten days. Group IV rats were treated with *H. auriculata* extract alone. At the end of experimental period, animals were subjected to mild ether anesthesia, blood was collected from retro orbital plexus and the serum was separated by centrifugation at 3000 rpm for 15 min at 4°C. Animals were sacrificed by cervical decapitation and the liver was excised, washed in ice cold saline and blotted to dryness. A 10% homogenate of the liver tissue was prepared in Tris-Hcl buffer (0.1 M; pH 7.4), centrifuged at 1000 rpm for 10 min at 4°C to remove the cell debris. The clear supernatant was used for further biochemical assays.

Biochemical assays

Aspartate and alanine transaminases (AST and ALT) were assayed according to the method of King 1965, acid and alkaline phosphatase enzymes in serum by the method of Bergmeyer, 1963 as described by Balasubramanian et al. 1983. Lipid peroxidation (LPO) was determined in the liver tissue as described by Devasagayam 1987, superoxide dismutase (SOD) according to Marklund and Marklund 1974 and catalase (CAT) according to the method of Sinha 1972. The activity of glutathione peroxidase (GPx) was estimated by the method of Rotruck et al. 1973. The hepatic reduced glutathione (GSH) content was estimated by the method of Moron et al. 1979. Vitamin C and Vitamin E were estimated by the method of Omaye et al. 1971 and Desai 1984 respectively. The activity of glutathione metabolizing enzymes such as glutathione reductase (GR), glucose -6-Phosphatase dehydrogenase (G6PD) and glutathione -S- transferase (GST) were estimated by the method of Staal et al. 1969, Beutler 1963, Habig et al. 1974 respectively.

Statistical analysis

The data obtained was subjected to One way ANOVA and Tukey's multiple comparison test was performed using SPSS statistical package (Version 7.5). Values are expressed as mean ± S.D. Value p< 0.05 was considered significant.

Result

Fig 1 shows the status of serum AST, ALT, ACP and ALP in control and experimental animals. The levels of the marker enzymes such as AST and ALT were significantly increased in serum with 27.8% and 39.5% respectively due to mercuric chloride induced hepatotoxicity when compared to group I control. In addition the ACP and ALP levels were also increased with 24.5% and 28.5% respectively in group II animals when compared with group I control animals. On the other hand, the increased activities of AST, ALT, ACP and ALP in serum of group II mercuric chloride induced hepatotoxicity were significantly decreased in group III *H. auriculata* post treated animals. However, there was no significant changes were

observed in group IV *H. auriculata* alone treated animals when compared to group I control animals.




Figure 1: The levels of transaminases and phosphatases enzymes in serum of control and HgCl₂ treated rats. Units are expressed as: AST, ALT = μ moles of pyruvate liberated/mg protein/min. ACP, ALP = μ moles of p-nitrophenol liberated/mg protein/min. Each value represents mean ± SD of six animals, a – Group II, III & IV compared with Group I, b – Group III compared with Group II *p<0.001; #p<0.01; @p<0.05; NS – No significant

The changes in the LPO level of liver tissue of control and experimental rats were illustrated in Fig. 2. LPO was significantly increased (around 2 fold) in HgCl₂ treated rats (group II) when compared with control rats (group I). Whereas treatment with *H. auriculata* group II animals showed a significant fall in the LPO level.




Figure 2: Effect of *H.auriculata* on lipid peroxidation in liver tissue of control and experimental rats during HgCl₂ induced toxicity. Each value represents mean ± SD of six animals, a – Group II, III & IV compared with Group I, b – Group III compared with Group II *p<0.001; #p<0.01; @p<0.05; NS – No significant.




Figure 3: Effect of *H.auriculata* on enzymic and non-enzymic antioxidants in the liver of control and experimental animals. Unit are expressed as: SOD=units /mg protein; CAT=μmoles of H₂O₂ consumed /mg protein/min; GPx=μ g of GSH utilized/mg protein/min; GSH= μg/mg protein/min; VIT-E and VIT-C=mg/g of wet tissue.

The changes in the level of enzymic and non-enzymic antioxidants such as SOD, CAT, GPx, GSH, Vit-E and Vit-C in the liver of control and experimental rats were given in Fig -3. A significant decrease (2 fold) in the activity of these radical scavengers were noticed after the administration of HgCl₂ in group II rats, when compared with group I rats.

On the contrary, post treatment with *H. auriculata* the activities of these enzymic antioxidants were significantly reversed to normal. The activities of non-enzymic antioxidants such as GSH, Vitamin C and Vitamin E were significantly decreased (2 fold) in group II rats treated with mercuric chloride, when compared with group I control rats, which were significantly reverted to normal level in group III *H. auriculata* post treated animals (Fig 4).




Figure 4: Each value represents mean ± SD of six animals, a – Group II, III & IV compared with Group I, b – Group III compared with Group II *p<0.001; #p<0.01; @p<0.05; NS – No significant.




Figure 5: Effect of *H.auriculata* on glutathione metabolizing enzymes in liver of control and HgCl₂ treated rats. Enzyme activities are expressed as : GR - nmoles of NADPH oxidized /min/mg protein ; G6PD - nmoles of NADPH formed /min/mg protein ; GST - nmoles of CDNB - GSH conjugate formed /min/mg protein. Each value represents mean ± SD of six animals, a – Group II, III & IV compared with Group I, b – Group III compared with Group II *p<0.001; #p<0.01; @p<0.05; NS – No significant

The levels of glutathione metabolizing enzymes such as GR, G6PD and GST were depicted in Fig 5. The levels of members of the glutathione metabolizing enzymes were significantly (p <0.05) decreased in group II toxicity bearing animals when compared to group I control animals, which were reverted to normal level on post treatment with *H. auriculata* and were comparable to group II animals.

Discussion

Redox disturbances are known to negatively impact the biological system through generation of ROS, which modify proteins, lipids and DNA (Choudiere, 1994). Liver being one of the target organs for mercury accumulation has witnessed the toxic insult of mercury by way of alterations in the activities of transaminases and phosphatases. Transaminases (ALT and AST) being an important class of enzymes linking carbohydrate and amino acid metabolism, have established a relationship between the intermediates of the citric acid cycle. These enzymes are regarded as markers of liver injury since; liver is the major site of metabolism (Liss et al. 1965). Phosphatases (ALP and ACP) are membrane bound and their alterations are likely to affect the membrane permeability and

produce derangement in transport of metabolites. Moreover, Plaa et al. 1986 have reported that these phosphatase enzymes act as an indicator of cholestatic changes. In this study, HgCl₂ administration to rats lead to marked elevation in the level of serum transaminases and phosphatases which is indicative of hepatocellular damage. This might be due to the possible release of these enzymes from the cytoplasm, into the blood circulation rapidly after rupture of the plasma membrane and cellular damage. Several researchers have reported that increased serum phosphatases and transaminases during mercury poisoning (Sharma et al. 2005) which clearly corroborates our study. The reason for the activities of marker enzymes during the toxicity might be due to the fact that oxidative damage in a cell or tissue occurs when the concentration of reactive oxygen species (O₂, H₂O₂, and OH) exceeds beyond the antioxidant capability of the cell (Winterbourn 1993). Treatment with the ethanolic extract of *H. auriculata* reduced the level of above mentioned markers in serum of HgCl₂ induced rats. This indicates that *H. auriculata* tends to prevent liver damage by maintaining in the integrity of plasma membrane, thereby suppress the leakage of enzymes through membranes, exhibiting hepatoprotective activity. It is reported that the products of LPO inactivate cell constituents by oxidation or cause oxidative stress through undergoing radical chain reaction which ultimately leading to loss of membrane integrity (Sorg et al. 1998). The stimulation of LPO observed as result of Hg (II) administration could be due to the formation of free radicals thought an exhaustion of antioxidants leading to oxidative stress (Sanders et al. 1996). It has been reported that Hg (II) can inactivate a number of enzymes by blocking the functional sites through binding to sulphhydryl groups, which are part of catalytic or binding domains (Fridovich, 1986). Thus it was suggested that in addition to depletion of intracellular thiol pools, the oxidant pathway may be a primary mechanism of induction of the response for Hg to induce oxygen free radicals or promote formation of lipid peroxides. Free radical scavenging enzymes like SOD and CAT protect the biological systems from oxidative stress. The SOD dismutates superoxide radicals (O₂⁻) into hydrogen peroxide (H₂O₂) and O₂ (Murray et al. 2003). CAT further detoxifies H₂O₂ into H₂O and O₂ (Bhattacharjee et al. 2006). GPx also functions in detoxifying H₂O₂ similar to catalase. Thus, SOD, CAT and GPx act mutually and constitute the enzymic antioxidative defense mechanism against reactive oxygen species (Mahboob et al. 2001). The decrease in the activities of these enzymes in the present study could be attributed to the excessive utilization of these enzymes in inactivating the free radicals generated during the metabolism of HgCl₂. This is further substantiated by an elevation in the levels of lipid peroxidation. Similar reports have shown an elevation in the status of LPO in liver during HgCl₂ exposure (Shanmugasundaram et al. 2006) and our results are in accordance with these reports. Restoration in the levels of lipid peroxidation after administration of *H. auriculata* could be related to its ability to scavenge reactive oxygen species, thus preventing further damage to membrane lipids. Our results are in line with previous studies by Chaudiere, (1994) in which they have shown that *H. auriculata* exhibits excellent antioxidant property. Therefore this property of *H. auriculata* might have resulted in the recouplement in the activities of the above antioxidant enzymes to normalcy. In the present study, we have observed that the depletion of the levels of non-enzymic antioxidants such as GSH, vitamin-C and vitamin-E due to the excessive liver damage and oxidative stress caused by HgCl₂. The Non-enzymic antioxidants like vitamin-C and E act synergistically to scavenge the free radicals formed in the biological system. GSH acts synergistically with vitamin-E in inhibiting oxidative stress and acts against lipid peroxidation (George 2003). Vitamin-C also scavenges and detoxifies free radicals in combination with vitamin-E and glutathione (Das 1994). It plays a vital role by regenerating the reduced form of vitamin-E and preventing the formation of

excessive free radicals (Sies 1991). The decreased levels of these enzyme activities observed during HgCl₂ administration might be due to the excessive utilization of these vitamins in scavenging the free radicals formed during the metabolism of HgCl₂. *H. auriculata* treatment effectively restored the depleted levels of these non-enzymic antioxidants caused by HgCl₂. *H. auriculata* has been reported to maintain the GSH homeostasis in the system as suggested by Vijayakumar et al. (2006) and this might be the reason for elevated glutathione levels observed during *H. auriculata* treatment. Increase in GSH levels in turn contributes to the recycling of other antioxidants such as vitamin-E and vitamin-C (Exner et al. 2000). It shows that *H. auriculata* maintains the levels of antioxidant vitamins C and E by maintaining GSH homeostasis, thereby protecting the cells from further oxidative stress. A significant decrease in the activities of GSH dependent enzymes, G6PD, GST and GR were observed in HgCl₂ treated rats, which may be due to the decreased expression of these antioxidants during hepatic damage. Further, the decreased levels of cellular GSH might have also caused a reduction in their activities as GSH is a vital co-factor for these enzymes and these observations are in accordance with the reports of Gstraunthaler et al. (1983) in which they have demonstrated that HgCl₂ induced hepatic injury was escorted by a substantial fall in hepatic GSH level, GR, G6PD, GPx and GST activity, which improved on administration of antioxidants. It is likely that post treatment of *H. auriculata* similarly maintains the activity of GR and GST in the liver by inhibiting lipid per-oxidation and maintaining GSH levels. This is an indicative of the potent antioxidant activity possessed by *H. auriculata*.

Conclusion

From the present study, it can be concluded that HgCl₂ induced toxicity involves oxidative stress in its pathophysiology. The ethanolic extract of *H. auriculata* possesses a potent free radical scavenging and antioxidant activities. The extract is capable of increasing/maintaining the levels of antioxidant molecules and antioxidant enzymes in the liver and also of protecting against oxidative damage to liver tissues. The preliminary chemical examination of ethanolic extract of *H. auriculata* extract has shown the presence of polyphenols, flavonoids, triterpenes and sterols which may be responsible for the antioxidant activity. Further studies are in progress in order to the isolation of more active compounds present in the extract.

Acknowledgement

The authors extremely grateful to Dr.R.Venkatakrishna Murali, M.D., Ph.D., Professor and Head, Department of Pharmacology and Environmental Toxicology, Dr.A.L.Mudhaliar Post Graduate Institute of Basic Medical Sciences, University of Madras, Taramani, Chennai – 600113 for providing the laboratory facilities.

References

- Ahmed S, Rahman A, Mathur M et al (2001) Antitumor promoting activity of *Asteracantha longifolia* against experimental hepato carcinogenesis in rats. Food and Chemical Technology 39: 19-28
- Balasubramanian MP, Dhandaythapani S, Nellaiyappan K et al (1983) Comparative Studies on Phosphomonoesterase in helminthes. Helminthologia 20 : 111-120
- Bergmeyer HO (1963) Methods of enzymatic analysis. Academic press, New York
- Beutler E (1983) Active transport of glutathione disulfide from erythrocytes. In: Larson A, Orrenius S, Holmgren A, Mannerwik B, editors. Functions of glutathione-Biochemical,

- physiological, toxicological and clinical aspects. New York: Raven Press pp. 65
- Bhattacharjee R, Sil PC (2006) The protein fraction of *Phyllanthus niruri* plays a protective role against acetaminophen induced hepatic disorder via its antioxidant properties. *Phytother Res* 20 : 595-601
- Chaudiere J (1994) Some chemical and biochemical constraints of oxidative stress in living cells. In: Rice-Evans, C.A., Burdon, R.H. (Eds.), *Free radical damage and its control*. Elsevier Science Amsterdam pp: 25-66
- Chaurasia SS and Kar A (1997) Protective effects of Vitamin E against lead-induced deterioration of membrane associated type -I iodothyronine 5' - monodeiodinase (5'D-I) activity in male mice. *Toxicol* 124: 203 - 209
- Clarkson TW (1972) The pharmacology of mercury compounds. *Ann Rev Pharmacol Toxicol* 12: 375 - 406
- Comporti M (1985) Biology of Disease: Lipid peroxidation and cellular damage in toxic liver injury. *Lab Invest* 53: 599- 623
- Das S (1994) Vitamin E in the genesis and prevention of cancer. A review *Acta Oncol* 33: 615-619
- Desai ID (1984) Vitamin E analysis methods for animal tissues. *Meth Enzymol* 105: 138-147
- Devasagayam TPA and Tarachand U (1987) Decreased lipid peroxidation in the rat kidney during gestation. *Biochem Biophys Res Commun* 145 : 134-138
- Duh PD (1998) Antioxidant activity of burdock (*Arctium lappa* Linn): its scavenging effect on free radical and active oxygen. *J American Oil Chem Society* 75: 455 - 461
- Dwivedi RS, Kaur G, Srivastava RC and Krishnamurthy CR (1984) Lipid peroxidation in tin intoxicated partially hepatectomized rats. *Bull Environ Contam. Toxicol* 33: 200 – 209
- Exner R, Wessner B, Manhart N et al (2000). Therapeutic potential of glutathione. *Wien Klin Wochenschr* 112: 610-616.
- Fernando MR, Wickramasinghe SMDN, Thabrew EH (1989) A preliminary investigation of the possible hypoglycemic activity of *Asteracantha longifolia*. *J Ethno Pharmacol* 27: 7-14
- Fitzgerald WF and Clarkson TW (1991) Mercury and monomethylmercury: Present and future concerns. *Environ Health Perspectives* 96: 159-166
- Fridovich I (1986) Superoxide dismutases. *Adv Enzymol* 58: 61-97
- George J (2003) Ascorbic acid concentrations in dimethylnitrosamine-induced hepatic fibrosis in rats. *Clin Chim Acta* 335: 39-47
- Gstraunthaler G, Pfaller W and Kotanko P (1983) Glutathione depletion and in vitro lipid peroxidation in mercury and maleate induced acute renal failure. *Biochem Pathol* 32: 2969-2972.
- Habig WH, Pabste MJ and Jakboy WB (1974) Glutathione-S-transferase: the first enzymatic step in mercapturic acid formation. *J Biol Chem* 249 : 7130-7139
- Halliwell B and Gutteridge JMC (1990) Free radical and catalytic metal ions in human disease: an overview. *Meth Enzymol* 186: 1-85.
- Huang YL, Cheng SL and Lin TH (1996) Lipid peroxidation in rats administered with mercuric chloride. *Biol Trace Elem Res* 52: 193 - 206
- Ito N, Fukushima S, Hasegawa A et al (1983) Carcinogenicity of butylated hydroxyanisole in F344 rats. *J National Cancer Institute* 70: 343 - 347
- Jayatilak PG, Pardanani DS, Murty BD et al (1976) Effect of an indigenous drug (speman) on accessory reproductive functions of mice. *Ind J Exp Biol* 14: 170-173
- King J (1965) The transferases-alanine and aspartate transaminases. In: Van D (ed.), *Practical Clinical Enzymology* Nostrand Company Limited London pp: 191-208
- Kinsella JE, Frankel E, German B et al (1993) Possible mechanism for the protective role of the antioxidant in wine and plant foods. *Food Tech* 47: 85- 89
- Lewis SE and Wills ED (1962). Destruction of sulphhydryl groups of proteins and amino acids by peroxides of unsaturated fatty acids. *Biochem Pharmacol* 11: 901- 912
- Liss G, Greenberg RA and Tamburro CH (1985). Use of serum bile acids in the identification of vinyl chloride hepatotoxicity. *Am J Med* 78: 68-76
- Mahboob M, Shireen KF, Atkinson A et al (2001) Lipid peroxidation and antioxidant enzyme activity in different organs of mice exposed to low level of mercury. *J Environ Sci Health Part B* 36: 687-697
- Marklund SL, Marklund G (1974). Involvement of superoxide anion radical in the auto oxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 47: 469-474
- Mazumdar UK, Gupta M, Maiti S et al (1997) Antitumor activity of *Hygrophila spinosa* on Ehrlich ascites carcinoma and sarcoma 180 induced in mice. *Ind J Exp Biol* 35: 473-477
- Moron MS, Diferece JW and Mannerwik KB (1979) Levels of glutathione, glutathione reductase and glutathione-S-transferase activities in rat lung and liver. *Biochim Biophys Acta* 582 : 67-78
- Murray RK, Granner DK, Mayes PA et al (2003) *Harper's Illustrated Biochemistry*, 26th. The McGraw-Hill Companies Inc
- Omaye ST, Turnbull JD, Saulberlich HE (1971) Selected methods for the determination of ascorbic acid in animal cell, tissues and fluids. *Meth Enzymol* 62: 1-11
- Plaa GL, Hewitt WR (1986) Detection and evaluation of chemically induced liver injury. In: Hayes, A.W (Eds). *Principles and Methods of Toxicology* New York: Raven Press pp: 401 - 441
- Quasim C, Dutta NL (1967) Chemical investigation of *Asteracantha longifolia* Nees. *J Ind Chem Soc* 44: 82-83
- Rotruck JT, Pope AL, Ganther AL (1973) Selenium: Biochemical role as a component of glutathione peroxidase purification and assay. *Science* 179: 588-590
- Sanders BM, Goering PL, Jenkins K (1996) The role of general and metal- specific cellular responses in protection and repair of metal- induced damage: Stressproteins and metallothioneins. In: Chang, L.W (Ed). *Toxicology of Metals*. CRC Press, Boca Raton, Florida, USA pp: 165-187
- Shailajan S, Chandra N, Sane RT et al (2005) Effect of *Asteracantha longifolia* Nees. against CCl4 induced liver dysfunction in rat. *Ind J Exp Biol* 43: 68-75
- Shanmugasundaram P, Venkataraman S (2006) Hepato protective and antioxidant effects of *Hygrophila auriculata* (K. Schum) Heine Acanthaceae root extract. *J Ethno Pharmacol* 104: 124-128.
- Sharma MK, Madhu Kumar and Ashok Kumar (2002) *Ocimum sanctum* aqueous leaf extract provides protection against Mercury induced toxicity in Swiss albino mice. *Ind J Exp Biol* 40: 1079- 1082
- Sharma MK, Madhu Kumar and Ashok Kumar (2005) Protection against mercury -induced renal damage in Swiss albino mice by *Ocimum sanctum*. *Environ Toxicol Pharmacol* 19 :161- 167
- Singh A and Handa SS (1999) Hepatoprotective activity of thioacetolens and *Hygrophila auriculata* against paracetamol and thioacetamide intoxication in rats. *J Ethno Pharmacol* 49: 119-126
- Sinha AK (1972) Colorimetric assay of catalase. *Anal Biochem* 47: 389-394
- Sorg O, Schilter B, Honegger P et al (1998) Increased vulnerability of neurons and glial cells to low concentration of methylmercury in a prooxidant situation. *Acta Neuropathol* 96: 621-627
- Staal GEJ, Visser J and Veeger C (1969) Purification and properties of glutathione reductase of human erythrocytes. *Biochem Biophys Acta* 185: 39-48

- Tan KH, Mayer DJ, Belin L and Ketterer B (1984) Inhibition of microsomal lipid peroxidation by glutathione transferase B and A. A role of endogenous phospholipids. *Biochem J* 320 : 243-254
- Velioglu YS, Mazza G, Gao L, Oomah BD (1998) Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J Agri Food Chem* 46: 4113-4117
- Vijayakumar M, Govindarajan R, Rao GMM et al (2006) Action of *Hydrophila auriculata* against streptozotocin - induced oxidative stress. *J Ethno Pharmacol* 104: 356 - 361
- Winterbourn CC (1993) Superoxide as an intracellular sink. *Free Radic Biol Med* 14: 85- 90
- Yuting C, Rongliang Z, Zhongjian J, Young J (1990) Flavonoids as superoxide scavenger and antioxidants. *Free Radical Biol Med* 9: 19-21
- Zhang D, Yasuda T, Yu Y et al (1996) Ginseng extract scavenges hydroxyl radical and protects unsaturated fatty acids from decomposition caused by iron mediated lipid peroxidation. *Free Radical Biol Med* 2: 145- 150