# Cold-active detergent-stable extracellular α-amylase from *Bacillus cereus* GA6: Biochemical characteristics and its perspectives in laundry detergent formulation

### Roohi<sup>\*</sup>, Mohammed Kuddus, Saima

Received: 02 November 2012/Received in revised form: 29 November 2012, Accepted: 17 March, 2013, Published online: 30 June 2013, © Sevas Educational Society 2008-2013

#### Abstract

A psychro-halo-tolerant *Bacillus cereus* GA6 was isolated from soil of Gangotri glacier, Western Himalaya that produced maximum cold-active  $\alpha$ -amylase at 20±1 °C after 96 h of incubation in alkaline medium by using glycerol and ammonium acetate as a substrate. The enzyme was purified to homogeneity (55 kDa) with 53.33% yield and 175.92 U/mg specific activity; and was stable at low temperature (4-37 °C) and in alkaline medium (pH 7-11). The purified enzyme showed maximum activity at 22±1 °C (pH 9) along with K<sub>m</sub> and V<sub>max</sub> value of 0.27 mg/ml and 2600 U/ml, respectively. The activity was strongly inhibited by Fe<sup>2+</sup>, Zn<sup>2+</sup>, H<sub>2</sub>O<sub>2</sub> and CuSO<sub>4</sub> while enhanced considerably by Ca<sup>2+</sup>. The hydrolysates of soluble starch by the enzyme were mainly glucose, maltose and maltotriose. The enzyme showed exceptional resistance ability against chemical denaturants (Urea, SDS and EDTA) and significant compatibility with commercial laundry detergents thus advocating its suitability for detergent formulation.

**Keywords:** Cold-active enzyme; α-amylase; *Bacillus cereus*; biocatalyst; biodetergent.

#### Introduction

Amylases (EC 3.2.1.1) are of ubiquitous occurrence and are produced by plants, animals, and microbes but microbial sources are

#### Roohi<sup>\*</sup>

Protein Research Laboratory, Department of Biotechnology and Bioengineering, Integral University, Lucknow-226026, India

Tel.: 00919450358872 \*Email: roohi0607@gmail.com

#### **Mohammed Kuddus**

Department of Biochemistry, University of Hail, Hail, Kingdom of Saudi Arabia

#### Saima

Protein Research Laboratory, Department of Biotechnology and Bioengineering, Integral University, Lucknow-226026, India

the most preferred one for large scale production to meet industrial demands (Pandey et al. 2000). Out of different factors affecting microbial environment, temperature is one of the most important as it influences most biochemical reactions. Thereby psychrotolerantmicrobes have developed various adaptive strategies to live in harsh environmental conditions (Margesin et al. 2002). Psychrotrophs are known to synthesize enzymes with low activation energies and high activities at low temperatures (Morita et al. 1997) conferring considerable progress toward energy savings. These enzymes are of commercial interest because they might be used in applications at low temperature and there is scientific interest in the relationship between protein structure and thermal stability of enzymes.

Amylases constitute one of the most important groups of industrial enzymes and account for nearly 25% of the total sale of enzymes (Burhan et al. 2003). Alpha amylases (EC 3.2.1.1) belongs to the enzyme class of hydrolases which randomly cleaves the 1,4-α-Dglucosidic linkages between the adjacent glucose units in linear amylose chain of starch. For commercial purpose of enzymes, detergent industries are the primary consumers in terms of both volume and value. Detergent enzymes account for about 30% of the total worldwide enzyme production and represent one of the largest and most successful applications of modern industrial biotechnology (Rao et al. 1998). The use of enzymes in detergent formulation not only enhances the detergent ability to remove tough stains but also makes the detergent environmentally safe. Out of the vast pool of enzymes exploited in detergent industries worldwide, there is always a need of new enzymes with novel properties that can further enhance the wash performance of currently used enzyme-based detergents (Gupta et al. 1999). Amylases are used in the formulation of enzymatic detergent, and 90% of all liquid detergents contain these enzymes (Hmidet et al. 2009). These enzymes are used in detergents for laundry and automatic dishwashing to degrade the residues of starchy foods. Most of the reported cold-active enzymes for detergent application were originated from Antarctic ecosystem and were mainly proteases (Gerday et al. 1997; Russel et al. 1997).

Comparatively little information is available on cold-active  $\alpha$ amylases from microorganisms isolated from glacier regions, which may be an ideal habitat for cold-adapted microorganisms. The present study includes production optimization, purification and biochemical characterization of cold-active  $\alpha$ -amylase produced by novel psychro-tolerant *Bacillus cereus* strain GA6 isolated from soil of Gangotri glacier, Western Himalaya, India and its application in detergent formulation for cold washing.

#### **Materials and Methods**

#### Isolation of cold-active $\alpha$ -amylase producing bacteria

In order to isolate psychro-tolerant bacteria, twenty five soil samples were collected from vicinity of Gangotri glacier, Western Himalaya, India during winter season of 2011. The glacier is situated in between  $30^{\circ}44'-30^{\circ}56$ 'N and  $79^{\circ}04'-79^{\circ}15$ 'E having temperature 2-5 °C in summer and subzero in winter. The samples were serially diluted in sterile cold saline solution and subjected to isolation of  $\alpha$ -amylase producing microbes at low temperature (Mishra and Behera 2008). Production of  $\alpha$ -amylase was carried out by using the modified method of Abe et al. (1988). Protein concentration was determined by the method of Lowry et al. (1951) using BSA as standard.

#### Alpha-amylase assay

Amylolytic activity with starch (Himedia) as a substrate was assayed by the method of Swain et al. (2006). This amylase assay was based on reduction in blue color intensity resulting from enzyme hydrolysis of starch. The reaction mixture consisted of 0.2 ml enzyme (cell free supernatant), 0.25 ml of 0.1% soluble starch solution and 0.5 ml of phosphate buffer (0.1M, pH 6.0) incubated at 50°C for 10 min. The reaction was stopped by adding 0.25 ml of 0.1N HCl and color was developed by adding 0.25 ml Gram's iodine solution. The optical density of blue color solution was determined by using spectrophotometer at 690 nm. The activity was expressed in units. One unit of enzyme activity is defined as the quantity of enzyme that causes 0.01% reduction of blue color intensity of starch-iodine solution at 50°C in one min per ml.

#### Identification of potential isolates

The potential isolate was identified and characterized by using Bergeys Manual of Systematic Bacteriology (Garrity 1985). The identity of the isolate was further confirmed by 16S rRNA analysis as described in our previous article (Roohi et al. 2011).

## Optimization of media and fermentation conditions for enzyme production

The effect of incubation period on enzyme production and cell growth was determined by inoculating production media with bacterial isolate (1%) and incubated at 20±2 °C for different time intervals (24-168 h) at 120 rpm. Alpha-amylase production was estimated after every 24 h for 8 days using the method of Swain et al. (2006). Growth kinetics was obtained by measuring the cell density at 660nm. To determine optimum temperature and pH for fermentation, the inoculated media was incubated at different temperatures (4-50 °C) and pH (pH 5-12) in a rotary shaker (120 rpm) and also in static condition. The effect of additional supplements (1%) in production media such as glucose, sucrose, maltose, lactose and glycerol as a carbon source; and glycine, casein, yeast extract, ammonium acetate and ammonium sulphate as a nitrogen source were evaluated in terms of enzyme yield. To assess the impact of metals, media was supplemented with maximum tolerance level of different metal ions and incubated under optimized conditions for 48 h. The heavy metals used were  $Ca^{2+}$ ,  $Cu^{2+}$ ,  $Zn^{2+}$ ,  $Fe^{2+}$ ,  $Mg^{2+}$  and  $Hg^{2+}$ . Amylase activity was measured as per standard protocol (Swain et al. 2006).

#### *Extraction and purification of cold-active* $\alpha$ *-amylase*

The isolated strain Bacillus cereus GA6 was grown in optimized conditions and cells were removed by centrifugation at 10000g for 10 min at 4°C. Cell-free supernatant was subjected to ammonium sulfate precipitation (10-80% w/v) and the precipitate was resuspended in 0.1M phosphate buffer (pH 6) that was dialyzed against the same buffer for 24 h. The amount of salt required for percentage saturation was calculated by the method given by Deutscher (1990). The partially purified protein was subjected to ion exchange chromatography on a DEAE cellulose column (Bangalore Genei, India) that was pre-equilibrated with phosphate buffer (0.1M, pH 6). The protein was eluted (flow rate 60 ml/h) with a linear gradient of NaCl (0.1 to 1M) in the same buffer. A total of 40 fractions were collected and assayed for protein and enzyme activity. All subsequent steps were carried out at 4°C. Fractions having highest amylase activity was desalted and concentrated by dialysis and lyophilization, respectively.

#### Characterization of purified $\alpha$ -amylase

#### Determination of optimum pH and pH stability

The effect of pH on the activity of  $\alpha$ -amylase was measured by using the following buffers; 0.2M acetate (pH 4-5), 0.1M phosphate (pH 6-8), 0.05M borax-NaOH buffer (pH 9-10) and 0.05M sodium hydrogen phosphate buffer (pH 11-12) (Aneja 2003). Reaction mixtures were incubated at 20°C for 30 min and the respective activity was measured. To determine pH stability, the enzyme was pre-incubated at different pH (4-12) for 1 h but without substrate and then estimated the residual activity as per standard assay procedure.

#### Determination of optimum temperature and thermal stability

The effect of temperature on the enzyme activity was determined by performing the previously described standard assay procedure within a temperature range of 4-45 °C by incubating the reaction mixture for 1 h. Thermo-stability of the enzyme was examined by pre-incubating the enzyme for 3 h without substrate within a temperature range of 4-45 °C and residual enzyme activity was determined.

#### Effect of metal ions and inhibitors

The effect of metal ions and inhibitors on the activity of purified  $\alpha$ amylase was investigated by pre-incubating the enzyme with different metal ions (Ca<sup>2+</sup>, Zn<sup>2+</sup>, Hg<sup>2+</sup>, Co<sup>2+</sup>, Ba<sup>2+</sup>, and Cu<sup>2+</sup>; 5mM) and inhibitors (EDTA, H<sub>2</sub>O<sub>2</sub> and Urea, 1-10mM; CuSO<sub>4</sub> and SDS, 0.1-1%) for 30 min at 20±2 °C.

## Effect of substrate concentration on $\alpha$ -amylase activity and their kinetic studies

To determine  $K_m$  and  $V_{max}$  for enzyme-substrate catalyzed reaction; 0.1, 0.2, 0.5, 0.7, 1.0, 1.2 and 1.4 mg/ml soluble starch in 0.1M phosphate buffer (pH 6.0) was mixed with purified  $\alpha$ -amylase (5662 units) and the mixture was incubated at 22°C for 30 min and the reaction was stopped immediately by heating at 100°C for 2 min.  $K_m$  and  $V_{max}$  values were obtained from Lineweaver-Burk plot (1/[V] ml/Units versus 1/[S] ml/mg) and expressed as the mean of the three different experiments.

#### Thin-Layer Chromatography (TLC) analysis

The hydrolysis products of starch were identified by TLC on silica gel plates using the method of Zhang and Zeng (2008). Soluble starch (1%) were hydrolyzed by incubating with amylase in 0.1M phosphate buffer, pH 6 at 20°C for 6 h. Aliquots (5 ml) of the reaction mixtures were chromatographed on a silica gel with chloroform–acetic acid–ddw (18:21:3, v/v), and the products were detected by spraying the gel with aniline-diphenylamine-phosphate followed by baking at 120°C for 30 min. Standards used were Glucose (S1), maltose (S2), maltotriose (S3) and maltotetraose (S4).

#### SDS-PAGE and zymogram analysis

The molecular weight of purified amylase was determined by SDS-PAGE. Electrophoresis was performed by the method of Laemmli (1970). The gel was stained with Coomassie Brilliant Blue R-250 and the relative molecular mass of the protein was calculated using standard protein markers (Sigma), run simultaneously. SDS-PAGEzymogram was performed for identity confirmation by the proposed method of Lin et al. (1998).

#### Evaluation of enzyme for use in detergent formulation

#### Compatibility and stability analysis

Application of cold-active purified  $\alpha$ -amylase as a detergent additive was studied by using the method of Kuddus and Ramteke (2011). The enzyme compatibility with commercially available laundry detergents was studied. The used detergents were Ariel and Tide (Procter and Gamble, India), Surf Excel and Wheel (Hindustan Lever Ltd., India) and Ghari (Rohit Surfactants Private Limited). The detergent solution (1% w/v) was boiled for 15 min to inactivate the enzymes that could be part of their formulation (enzyme assay was performed to check the activity). The cold-active amylase (10 mg/ml) was then incubated with detergent solution for different time intervals (0.5-3 h) at 22±1 °C and the residual activity was determined in comparison to control (without any detergent).

#### Wash performance analysis

Wash performance analysis of purified enzyme was studied on white cotton cloth pieces (8x8 cm) stained separately with food gravy and mixture of baby food and chocolate and then left for overnight. The following sets were prepared for cold-washing:

a) Flask containing distilled water (100 ml) + stained cloth (with food gravy and mixture of baby food and chocolate, independently). b) Flask containing distilled water (98 ml) + stained cloth + 2 ml Tide detergent (1% w/v).

c) Flask containing distilled water (96 ml) + stained cloth + 2 ml Tide detergent (1% w/v) + 2 ml purified  $\alpha$ -amylase (4774 units/ml). The flasks were incubated for 30 min (22°C) in shaking condition (100 rev/min) and rinsed with cold water then air dried. The wash performance was analyzed by measuring relative reflectance of cloth pieces using digital reflectance meter (Aimil Ltd., New Delhi) to test enzyme efficiency for stain removal. Untreated stained cloth pieces were taken as control (Kuddus and Ramteke 2011; Beg and Gupta 2003; Adinarayana et al. 2003).

#### Statistical analysis

All the experiments were carried out in triplicates and the standard error in results was within 5%. Appropriate controls were maintained in all the experiments.

#### Results

#### Psychrotrophic α-amylase producing bacteria

Total thirty prominent bacterial colonies were screened from soil of Gangotri glacier, producing cold-active extracellular  $\alpha$ -amylase. Depending upon larger hydrolysis zone on starch agar media at 15±2 °C, six isolates were subjected to enzyme production in amylase producing broth media. On the basis of maximum enzyme production (3000 units) one potential isolate, designated as GA6, was selected for further studies.

#### Identification of potential isolate

The isolated cold-active  $\alpha$ -amylase producing bacterium was identified on the basis of 16S rRNA gene alignment with BLAST search from the GenBank database at NCBI (MIDILABS Inc., USA) that showed 99% homology with *Bacillus cereus*. The phylogenetic tree (Fig. 1) was constructed using software MEGA 4.1 (Tamura et al. 2007). Final sequence of GA6 was submitted to GenBank having an accession no. HQ832575. The strain was Grampositive, rod shape, aerobic, non-motile, spore forming and catalase positive with opaque colonies and could grow up to 5 mm diameter in size. The organism grew well in the pH range of 7-11 and temperature range of 4-37 °C but did not grow above 40°C. The isolate could tolerate up to 8% NaCl concentration.



Fig. 1. Phylogenetic tree showing the homology of strain GA6 with *Bacillus cereus* 

#### *Optimization of* $\alpha$ *-amylase production*

Alpha amylase production by strain GA6 began after 48 h and reached maximum at 96 h (4364 units) in logarithmic phase at  $15\pm2$  °C (Fig. 2). However, the cell growth was maximal at 120 h of incubation which indicate that the production of enzyme was independent to cell growth. Optimum temperature was found to be 20°C over 96 h of incubation (Fig. 3). However, there was continuous decline in enzyme production with increase in incubation temperature and it was totally inhibited at 50°C. The pH of the culture media strongly affects many enzymatic reactions and transport of compounds across the cell membrane and from Fig. 3 it was clear that amylase production was maximum at pH 10 (4732 units).

Enzyme production leads a four-fold boost in shaking condition (4662 units) at 120 rpm as compared to unperturbed condition (1204 units) at 20°C after 48 h incubation, established the necessity of

aeration for growth of the strain. Enhanced production of amylase was achieved with glycerol (4744 units), ammonium acetate (4746 units) and  $Ca^{2+}$  (4780 units) while lactose, glycine and  $Fe^{2+}$  worsened the production (Table 1).

#### Purification and characterization of $\alpha$ -amylase

Cold-active  $\alpha$ -amylase was purified to homogeneity by precipitating with ammonium sulphate (50%) and using a single step ion-exchange chromatography on a DEAE-cellulose (Fig. 4). Enzyme



Fig. 2. Effect of incubation period on growth and enzyme production by GA6 at  $15\pm1^{\circ}C$ 



Fig. 3. Effect of pH and temperature on enzyme production by GA6 after 96 h of incubation

Table 1: Effect of carbon, nitrogen and metal ions on  $\alpha\text{-amylase}$  production by GA6

S. No.	Additional supplements (1%)	$\alpha$ -amylase activity (units)
	Control	3225
	Carbon source	
1	Glucose	1476
2	Sucrose	1952
3	Maltose	1952
4	Lactose	1238
5	Glycerol	4744
	Nitrogen source	
1	Casein	1746
2	Glycine	317.4
3	Yeast extract	1905
4	Ammonium acetate	4746
5	Ammonium sulphate	476.1
	Metal ions	
1	FeCl <sub>3</sub>	462
2	CuSO <sub>4</sub>	625
3	CaCl <sub>2</sub>	4560
4	MgCl <sub>2</sub>	3345
5	ZnSO <sub>4</sub>	1250
6	HgCl <sub>2</sub>	1250

was eluted from the column as unbound fractions with 0.7M NaCl gradient. The  $\alpha$ -amylase exhibited a specific activity of 175.92

U/mg, corresponding to a purification factor of 7.9-fold and a total yield of 53.33% (Table 2).

#### Optimum pH and pH stability

The effect of pH on  $\alpha$ -amylase activity is shown in Fig. 5. The activity increased almost linearly from pH 5 to 9 and it was optimum at pH 9. However, 89 and 90% of the total enzyme activity was manifested at pH 8 and 10, respectively. At pH 7 and 11 the activity was 66 and 54%, respectively. The pH stability result suggests that enzyme was relatively stable over a pH range of 8 to 11 and retained about 80, 82 and 65% residual activity at pH 8, 10 and 11, respectively.



Fig. 4:. The elution profile of  $\alpha$ -amylase from *B. cereus* GA6 by using DEAE cellulose column

C . .

... n

Table 2. Purification of amylase from <i>B. cereus</i> GA6							
Purification steps	Total activity (Units)	Total protein (mgL <sup>-1</sup> )	Specific Activity (Umg <sup>-1</sup> )	Purification (factor)	Yield (%)		
Crude	5010	225	22.26	-	100		
enzyme							
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> precip <sup>n</sup> (50%) (dialyzed)	3894	64	60.84	2.73	77.72		
DEAE- cellulose pool	2674	15.2	175.92	7.903	53.33		

(lyophilized)



Fig. 5. Influence of pH on the activity and stability of purified  $\alpha$ -amylase from *B. cereus* GA6

#### Temperature optima and thermal stability

The activity and stability of the lyophilized enzyme preparation assayed at temperatures ranging from 4-45 °C at pH 9 are shown in Fig. 6. Enzyme activity increased with temperature and was found optimum at 22°C. Activity decreased sharply above 37°C and becomes almost zero at 45°C. Enzyme showed stability between 437 °C. The enzyme retained 56, 78 and 80% stability at 4°C, 15°C and 28°C, respectively of that at 22°C.

#### Stability of enzyme with metal ions and inhibitors

This cold-active  $\alpha$ -amylase showed 185% catalytic activation in presence of 5mM Ca<sup>2+</sup> (Table 3). Strongest inhibitory effect was observed in the presence of Fe<sup>2+</sup> and Zn<sup>2+</sup> where activity becomes zero. Cu<sup>2+</sup>, Ba<sup>2+</sup>, Co<sup>2+</sup>, and Hg<sup>2+</sup> also inhibited the enzyme activity and only 38.95, 58.82, 76.46 and 36.46 % activity remained, respectively. Enzyme showed remarkable resistance against EDTA, SDS and Urea as more than 50% activity still remained when incubated for 30 min at 20°C while CuSO<sub>4</sub> (1%) and H<sub>2</sub>O<sub>2</sub> (10mM) obtained as a potent inhibitors.



Fig. 6. Effect of temperature on the activity and stability of purified  $\alpha$ amylase from *B. cereus* GA6

Table 3. Effect of metal ions and inhibitors on the activity of purified amylase

Inhibitor	Concentration	% Relative activity	
Control	-	100	
$Ca^{2+}(CaCl_2)$	5mM	185	
$Ba^{2+}$ (BaCl <sub>2</sub> )	5mM	58.82	
$Co^{2+}(CoCl_2)$	5mM	76.46	
$Hg^{2+}(HgCl_2)$	5mM	36.46	
$Cu^{2+}$ (CuSO <sub>4</sub> )	5mM	38.95	
$Zn^{2+}$ (ZnCl <sub>2</sub> )	5mM	0	
$Fe^{2+}$ (FeSO <sub>4</sub> )	5mM	0	
$H_2O_2$	1mM	25	
	10mM	0	
EDTA	1mM	70	
	10mM	45	
CuSO <sub>4</sub>	0.1%	10	
	1%	0	
Urea	1M	60	
	10M	35	
SDS	0.1%	50	
	1%	25	

Effect of substrate concentration on  $\alpha$ -amylase activity and their kinetic studies

When activity of purified cold-active enzyme was measured with different concentration of starch (0.1 to 1.4 mg/ml) in optimized conditions, it was observed that amylolytic activity increased with substrate concentration linearly and reached to maximum at 1 mg/ml and then becomes constant (figure not shown). A Lineweaver-Burk plot (Fig. 7) indicates that this enzyme has apparent  $K_m$  and  $V_{max}$  values of 0.27 mg/ml and 2600 Units/ml, respectively for the hydrolysis of soluble starch.

#### TLC analysis

TLC was used to analyze the hydrolysis patterns of soluble starch digested by the purified amylase (Fig. 8). The main hydrolysis products of soluble starch were mainly glucose, maltose, maltotriose and a little of maltotetraose also. This hydrolysis patterns revealed that amylase from *B. cereus* GA6 is a typical alpha-amylase as it hydrolyze the  $\alpha$ -(1,4)-glycosidic linkage only.



Fig. 7. Lineweaver-Burk plot for the hydrolysis of starch by purified  $\alpha$ -amylase from *B. cereus* GA6



**Fig. 8.** TLC analysis of hydrolysis products of soluble starch digested by *B. cereus* GA6  $\alpha$ -amylase. (Lane 1: product generated after 0.5 h; lane 2: product generated after 2 h; lane 3: product generated after 6 h; Std: standard marker, S1. Glucose, S2. Maltose, S3. Maltotriose, S4. Maltotetarose)

#### Molecular weight

For determination of molecular weight, enzyme preparations and known molecular weight markers were subjected to SDS–PAGE. The purified protein showed a single band equal to a molecular mass of about 55 kDa on SDS-PAGE confirming that enzyme is composed of single polypeptide chain and activity staining gel confirming high enzyme purity (Fig. 9).

#### Detergent Compatibility and stability analysis

Cold-active  $\alpha$ -amylase purified from *B. cereus* GA6 showed excellent stability and compatibility with a wide range of locally available commercial detergents (1% w/v) which is a mandatory requirement for cold-washing. It was most compatible with 'Tide' detergent retaining 92.27% activity after 0.5 hour incubation at 22°C and it retained 86, 78 and 72% activity after 1.0, 2.0 and 3.0 hour incubation, respectively (Table 4). Next to Tide, the enzyme was compatible with 'Ghari' detergent exhibiting 89% activity after 0.5 hour incubation, and retained 81.8, 74.9 and 70.7% activity after 1.0, 2.0 and 3.0 hours of incubation, respectively at 22°C.



Fig. 9. SDS-PAGE and activity staining analysis of the purified  $\alpha$ -amylase from B.cereus GA6. (M: Marker; Lane 1: Crude protein; Lane 2: Purified lyophilized α-amylase; Lane 3: activity staining of purified enzyme).

Remaining activity of the enzyme was more than 64% with rest of the tested detergents even after 3.0 hours of incubation.

Table 4. Compatibility of amylase from B. cereus GA6 with commercial detergents at 20±1 °C

Dotorgonto	Relative residual amylase activity (%)				
Detergents	0.5 hour	1.0 hour	2.0 hour	3.0 hour	
Surf Excel	88.27	80.20	70.5	65	
Wheel	85.55	79	71.38	66.22	
Tide	92.27	86.11	78	72	
Ariel	80.91	77.50	70	64.50	
Ghari	89.07	81.80	74.92	70.70	

Wash performance analysis at low temperature

The wash performance on fabrics is dependent on efficiency of amylases for the removal of stains. Result suggests that this coldactive alkaline amylase exhibited high efficiency for the removal of stains when used in combination with 'Tide' detergent at 22±2 °C. The wash performance analysis of mixture of baby food and chocolate stains on cotton fabric showed an increase in reflectance from 56 to 78% when washed with detergent and enzyme (combined) as compared to detergent only (Fig. 10). Similarly increase in reflectance was from 52 to 82% (Fig. 10) for removal of food gravy stain from white cloth, when washed with detergent supplemented with enzyme (Fig. 11). Therefore, it may be recommended that the supplementation of the cold-active alkaline  $\alpha$ amylase isolated from Bacillus cereus GA6 could significantly perk up the cleansing of the starchy stains resulting in complete stains removal at low temperature.



Fig. 10. Removal of food gravy and chocolate stain at 20°C by cold-active αamylase (reflectance analysis) (W: Water, D: Detergent, E:Enzyme)

#### Discussion

In industrial process, bacterial  $\alpha$ -amylases are produced mainly from cultures of Bacillus subtilis var. amyloliquefaciens (Uhlig, 1998; Goyal et al., 2005). Alpha-amylases from Bacillus stearotermophilus and Bacillus licheniformis are well characterized and are profoundly used in the starch-processing industry. Since, alkali-tolerance characteristic and stability toward chelators and detergents are important features for use of amylolytic enzymes in detergents and starch-processing, amylases from psychro-tolerant bacteria are of special interest as a source of novel cold-adapted enzymes (Leveque et al. 2000; Saxena et al. 2007).



Fig. 11. Wash performance of B. cereus GA6 α-amylase in combination with Tide detergent at 20°C. (A) Cloth stained with food gravy; (B) Gravy-stained cloth washed with water only; (C) Gravy-stained cloth washed with enzyme only; (D) Gravy-stained cloth washed with detergent only; (E) Gravy-stained cloth washed with detergent and enzyme

Maximum  $\alpha$ -amylase was produced in its logarithmic phase from *B*. cereus GA6 at 15°C at 96 h of incubation. Similar observations were also reported for Bacillus sp. (Wijbenga et al. 1991) where maximum amylase production occurred into its late logarithmic growth phase and continued to secrete well into the stationary phase. Gupta et al. (2008) also reported same time duration for highest aamylase production but in case of Aspergillus niger. The optimal temperature for amylase production and growth of the Bacillus cereus GA6 were found to be at 20°C. Similar findings were also reported for cold-adapted Pseudoalteromonas arctica GS230 (Lu et al. 2010) producing maximum amylase at 20 °C. However, Flavobacterium balustinum A201 (Morita et al. 1997) and Bacillus sp. A-001 (Lealem and Gashe 1994) showed its highest enzyme production at 30° and 35°C, respectively. On pH optimum analyses, B. cereus GA6 produced a peak in alkaline side i.e. at pH 10.0. Comparable to the result, optimum pH was 9.0, 8.0 and 7.5 for Actinomycete strain AE-19 (Poornima et al. 2008); Micrococcus antarcticus (Fan et al. 2009) and Bacillus sp. (Lealem and Gashe 1994), respectively. The pH tolerant  $\alpha$ -amylase from microbes are commercially important for detergent industry and can be successfully used as an additive to remove starch based stains. Result showing the necessity of Ca<sup>2+</sup> for production of amylase was in agreement with Vishwanathan and Surlikar (2001) where increased production of Aspergillus flavus a-amylase occurs in presence of 5mM Ca<sup>2+</sup>.

It has been reported that the alkaline amylase of Bacillus sp. strain IMD370 showed maximum activity at pH 10 (McTigue et al. 1995). Optimum pH was 8.7 as observed by Igarashi et al. (1998) and Kobayashi et al. (1992) for Bacillus sp. KSM-1378 and Natronococcus sp. AH-36, respectively. However B. cereus GA6 aamylase showed maximum activity at pH 9.0. The enzyme was highly active at pH range of 7-10 for 1 h at 20°C. The enzyme was 80% stable between pH ranges 8-11 for 1 h. These findings were highly similar with the literature result (Chakraborty et al. 2009) where  $\alpha$ -amylase of marine Streptomyces sp. D1, was stable in buffer systems having pH 7-11. These results suggest that the activity of the enzyme is higher in alkaline pH and are pH stable at low temperature, making this enzyme attractive for the detergent industry. This can be further commercially utilized for hidedehairing process, where dehairing is carried out at pH values between 8.0 and 10.0. Optimal temperature for GA6-amylase was found to be at 22°C which decreased sharply above 37°C. The temperature profile of Antarctic bacterium, Alteromonas haloplanctis (Feller et al. 1994) proposed totally contradictory results where the bacterium faced temperatures close to that of their environment (-2 to 4°C) but the apparent optimal activity temperature was between 30-40°C. Ambiguity was also noticed for A. psychrolactophilus (Smith and Zahnley 2005) showing optimum activity between 40-50°C. B. cereus GA6 a-amylase showed good temperature stability from 4-37°C, giving maximum activity below 25°C and thereby highly recommended for the category of true psychro-tolerant enzymes (Morita 1975). This cold-active amylase is of considerable interest as it can efficiently hydrolyze starch at ambient to low temperatures, thereby reduces costs in producing alcohol and to use in many other low-temperature applications where industrial fermentation flourishes.

Most  $\alpha$ -amylases vary in their response to the chelators, chemical denaturants and metal ions. Almost twice activation was noticed in activity of *B. cereus* GA6 alkaline  $\alpha$ -amylase with 5mM Ca<sup>2+</sup> while Fe<sup>2+</sup> and Zn<sup>2+</sup> were proved to be potent inhibitors. Positive effects of calcium were also reported by Arikan (2007) where 130% increment in enzyme activity was noticed in the presence of 5mM Ca<sup>2+</sup>. Same results for calcium were also reported by Sindhu et al. (2011), Brawn and Kelly (1993) and Dong et al. (1997). An inhibitory effect of Zn<sup>2+</sup> on the activity of amylases was also observed by Arikan (2007), and Mamo and Gessesse (1999). B. cereus GA6 α-amylase has also strong inhibition by 100% with  $CuSO_4$  (1%) and  $H_2O_2$ (10mM). In order to have applications in detergent industries, amylase must be stable to various detergent ingredients, such as denaturants and chelators. The amylase from B. cereus GA6 exhibited 50% activity when preincubated with 0.1% SDS and 25% activity with 1% SDS at 20°C for 30 min and also showed considerable resistance against EDTA, SDS and Urea. Several authors (Arikan 2007; Kikani and Singh 2011; Park et al. 2010) reported the existence of inhibitor resistant amylase (SDS, urea, EDTA) but were actually thermophilic or thermostable in nature. To the best of my knowledge this is the first report where cold-active B. cereus GA6 alkaline-amylase found to be a chelator resistant at low temperature range so are suitable in liquefaction of starch, in detergent and textile industries and in other industrial applications in fairly cold environments. This resistance, which is essential requirements, suggests that the use of enzyme as effective additive in detergents. 1 mg/ml soluble starch was found optimum for B. cereus GA6 amylase and it also showed less K<sub>m</sub> (0.27 mg/ml) and high V<sub>max</sub> (2500 Units/ml), suggesting strong affinity between enzyme and substrate. Our results are confirmed with the results in the literature where  $\alpha$ -amylase of halophilic *Bacillus cereus* MS6 showed maximum enzyme activity in 1 to 1.5% of starch concentration (Al-ZaZaee et al. 2011) and a K<sub>m</sub> of 0.88 mg/ml was obtained for a-amylase purified from Streptococcus bovis JB1 with

soluble potato starch (Freer 1993). The results of K<sub>m</sub> were in accordance with Li et al. (2011) and Alexsandro et al. (2011) where observed K<sub>m</sub> was 0.27 and 0.37 mg/ml, respectively. In other reports, Bano et al. (2011) also observed the Km and Vmax value of 2.68 mg/ml and 1773 Units/ml, respectively for α-amylase purified from Bacillus subtilis. The hydrolysis pattern of B. cereus GA6 amylase confirmed its type alpha as obtained by TLC analysis. These findings were highly similar with the literature results (Zhang and Zeng 2008; Li et al. 2011; Michelin et al. 2010) where glucose, maltose and maltotriose were found as final hydrolysates of soluble starch by purified cold-adapted amylase of Nocardiopsis sp. 7326, Rhizopus oryzae and Paecilomyces variotii, respectively. Molecular mass of 55 kDa having single polypeptide chain of B. cereus GA6  $\alpha$ -amylase was confirmed by the Zymogram analysis. Similar findings were also reported for Bacillus licheniformis (Haq et al. 2010) and Pseudomonas sp. 7193 (Zhang and Zeng 2008).

Compatibility and stability of *B. cereus* GA6 alkaline  $\alpha$ -amylase with commercial detergents at 20°C was excellent and retaining more than 92% activity after 0.5 h of incubation with 'Tide'. A similar compatibility result was obtained for *Bacillus* sp. SMIA-2 amylase retaining 75% of its activity after 20 min. of incubation with ,Tide, but at 50°C (Carvalho et al. 2008). Therefore, it may be suggested that the cold-active amylase from *B. cereus* GA6 is appropriate as an additive for commercial detergents for low temperature washing.

#### Conclusions

Cold-adapted *B. cereus* GA6 isolated from Gangotri glacier showed promising production of Ca<sup>2+</sup>-dependent, alkaline  $\alpha$ -amylase that was able to withstand a temperature up to 4°C for 24 h but having temperature optima of 22°C. Enzyme resistance against chemical denaturants and detergents were interesting properties as very few reports of such effects exist in the literature for psychro-tolerant amylase therefore suggests the potential of this enzyme in detergent industry for laundry and automatic dishwashing at low temperatures and to keep the environment safe and clean. The organism can also find applications in environmental bioremediation at cold regions.

#### Acknowledgements

Financial assistance from Council of Science and Technology, Uttar Pradesh, India (Project No. CST/3221) is gratefully acknowledged.

#### References

- Abe J, Bergman FW, Obata K, Hikuri S (1988) Production of raw starch digesting amylase by *Aspergillus* K-27. Appl Microbiol Biotech 27:447-450
- Adinarayana K, Ellaiah P, Prasad DS (2003) Purification and partial characterization of thermostable serine alkaline protease from a newly isolated *Bacillus subtilis* PE-11. AAPS Pharm Sci Tech 4(4):56-67
- Alexsandro SG, Silva RN, Lottermann MT, Morales AC, Ivares A, Pepe de Moraes LM, Torres FAG, de Freitas SM, Ulhoa CJ (2011) Biochemical and structural characterization of amy1: An alpha-amylase from Cryptococcus flavus expressed in Saccharomyces cerevisiae. Enz Res 157294-157301
- Al-ZaZaee MMA, Neelgund S, Gurumurthy DM, Rajeshwara AN (2011) Identification, characterization of novel halophilic *Bacillus Cereus* Ms6: a source for extra cellular α-amylase. Adv Enviro Biol 5(5):992-999
- Aneja KR (2003) Experiments in Microbiology, Plant Pathology and Biotechnology. New Age International, New Delhi, India

- Arikan B (2007) Highly thermostable, thermophilic, alkaline, SDS and chelator resistant amylase from a thermophilic *Bacillus* sp. isolate A3-15. Biores Technol 99(8):3071-3076
- Bano S, Qader SA, Aman A, Syed MN, Azhar A (2011) Purification and characterization of novel α-amylase from Bacillus subtilis KIBGE HAS. AAPS Pharm Sci Technol 12(1):255-261
- Beg QK, Gupta R (2003) Purification and characterization of an oxidation stable, thiol dependent serine alkaline protease from *Bacillus mojavensis*. Enz Microb Technol 32:294-304.
- Brawn SH, Kelly RM (1993) Characterization of amylolytic enzymes having both (alpha)-1,4 and (alpha)-1,6 hydrolytic activity from the thermophilic archea *Pyrococcus furiosus* and *Thermococcus litoralis*. Appl Environ Microbiol 59(8):2614– 2621
- Burhan AU, Nisa C, Gokhan C, Omer A, Ashabil, Osman G (2003) Enzymatic properties of a novel thermostable, thermophilic, alkaline and chelator resistant amylase from an alkaliphilic *Bacillus* sp. isolate ANT-6. Process Biochem 38:1397–1403
- Carvalho RVD, Correa TLR, Matos SJC, Mansur LRC, Martins MLL (2008) Properties of an amylase from thermophilic *Bacillus* sp. Braz J Microbiol 39:102-107
- Chakraborty S, Khopade A, Kokare C, Mahadik K, Chopade B (2009) Isolation and characterization of novel α-amylase from marine *Streptomyces* sp. D1. J Mol Catalyst B Enzym.58:17-23
- Deutscher MP (1990) Guide to protein purification: Methods in enzymology. Academic Press, New York
- Dong G, Vieille C, Savchenko A, Zeikus JG (1997) Cloning, sequencing and expression of the gene encoding extracellular alpha-amylase from *Pyrococcus furiosus* and biochemical characterization of the recombinant enzyme. Appl Environ Microbiol 63(9):3569-3576
- Fan HX, Liu Y, Liu ZP (2009) Optimization of fermentation conditions for cold-adapted amylase production by *Micrococcus antarcticus* and its enzymatic properties. Huan Jing Ke Xue 30(8):2473-2478
- Feller G, Narinx E, Arpigny J, Zekhnini Z, Swings J, Gerday C (1994) Temperature dependence of growth, enzyme secretion and activity of psychrophilic antarctic bacteria. Appl Microbiol Biotech 41:477–479
- Freer SN (1993) Purification and characterization of the extracellular alpha-amylase from streptococcus bovis JBL. Appl Environ microbial 59: 1398-1402
- Garrity GM (1985) Bergey's manual of systematic bacteriology book review. Williams and Wilkins, New York
- Gerday C, Aittaleb M, Arpigny JL, Baise E, Chessa JP (1997) Psychrophilic enzymes: a thermodynamic challenge. Biochem Biophys Acta 1342:119-131
- Goyal N, Gupta JK, Soni SK (2005) A novel raw starch digesting thermostable α-amylase from *Bacillus* sp. I-3 and its use in the direct hydrolysis of raw potato starch. Enz Microbiol Technol 37:723-734
- Gupta A, Gupta VK, Modi DR, Yadava LP (2008) Production and characterization of α-amylase from *Aspergillus niger*. Biotechnol 7:551-556
- Gupta R, Gupta K, Saxena RK, Khan S (1999) Bleach-stable alkaline protease from *Bacillus* sp. Biotechnol Lett 21:135-138
- Haq I, Ali S, Javed MM, Hameed U, Saleem A, Adnan F, Qadeer MA (2010) Production of alpha amylase from a randomly induced mutant strain of *Bacillus amyloliquefaciens* and its application as a desizer in textile industry. Pak J Bot 42(1):473-484
- Hmidet N, Ali N, Haddar A, Kanoun S, Alya S, Nasri M (2009) Alkaline proteases and thermostable α-amylase co-produced by *Bacillus licheniformis* NH1: Characterization and potential application as detergent additive. Biochem Eng J 47:71-79

- Igarashi K, Hatada Y, Hagihara H, Saeki K, Takaiwa M, Uemura T, Ara K, Ozaki K, Kawai S, Kobayashi T, Ito S (1998) Enzymatic properties of a novel liquefying alpha-amylase from an alkaliphilic *Bacillus* isolate and entire nucleotide and amino acid sequences. Appl Environ Microbiol 64:3282–3289
- Kikani BA, Singh SP (2011) Single step purification and characterization of a thermostable and calcium independent αamylase from *Bacillus amyloliquifaciens* TSWK1-1 isolated from Tulsi Shyam hot spring reservoir, Gujarat (India). Int J Biol Macromol 48(4):676-81
- Kobayashi T, Kanai H, Hayashi T, Akiba T, Akaboshi R, Horikoshi K (1992) Haloalkaliphilic maltotriose forming α-amylase from the archaebacterium *Natronococcus* sp. strain Ah-36. J Bacteriol 174:3439–3444
- Kuddus M, Ramteke PW (2011) Production optimization of an extracellular cold-active alkaline protease from *Stenotrophomonas maltophilia* MTCC 7528 and its application in detergent industry. Afr J Microbiol Res 5(7):809-816
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 227(5259):680–685
- Lealem F, Gashe BA (1994) Amylase production by a gram-positive bacterium isolated from fermenting tef (*Eragrostis tef*). J Appl Bacterio. 77:341-352
- Leveque E, Janacek S, Haye B, Belarbi A (2000) Thermophilic archeal amylolytic enzymes. Enz Microb Technol 26:3-14
- Li S, Zuo Z, Niu D, Singh S, Permaul K, Prior BA, Shi G, Wang Z (2011) Gene cloning, heterologous expression and characterization of a high maltose-producing α-amylase of *Rhizopus oryzae*. Appl Biochem Biotechnol 164(5):581-592
- Lin LL, Chyau CC, Hsu WH (1998) Production and properties of a raw starch-degrading amylase from the thermophilic and alkaliphilic *Bacillus* sp. TS-23. Biotechnol Appl Biochem 28(1):61–68
- Lowry OH, Rosebrough NJ, Farr AL, Randal RJ (1951) Protein measurement with the Folin phenol Reagent. J Biol Chem 193:265-275
- Lu M, Wang S, Fang Y, Li H, Liu S, Liu H (2010) Cloning, expression, purification, and characterization of cold-adapted αamylase from *Pseudoalteromonas arctica* GS230. Prot J 29(8):591-597
- Mamo G, Gessesse A (1999) Purification and characterization of two raw-starch-digesting thermostable α-amylase from a thermophilic *Bacillus*. Enz Microb Technol 25:433-438
- Margesin R, Feller G, Gerday C, Russell N (2002) Cold-adapted microorganisms: adaptation strategies and biotechnological potential. Wiley, New York
- McTigue M A, Kelly CT, Doyle EM, Fogarty WM (1995) The alkaline amylase of the alkalophilic *Bacillus* sp IMD 370. Enz Microb Technol 17:570–573
- Michelin M, Silva TM, Benassi VM, Peixoto-Nogueira SC, Moraes LA, Leao JM, Jorge JA, Terenzi HF, Polizeli ML (2010) Purification and characterization of a thermostable α-amylase produced by the fungus *Paecilomyces variotii*. Carbohydr Res 345(16):2348-53
- Mishra S, Behera N (2008) Amylase activity of a starch degrading bacteria isolated from soil receiving kitchen wastes. Afr J Biotech 7(18):3326-3331
- Morita RY (1975) Psychrophilic bacteria. Bacteriol Rev 39:144–167
- Morita Y, Nakamura T, Hasan Q, Murakami Y, Yokoyama K, Tamiya E (1997) Cold-active enzymes from cold adapted bacteria. JAOCS 74(4):441-444
- Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D, Mohan R (2000) Advances in microbial amylases. Biotechnol Appl Biochem 31:135-152

- Park KM, Jun SY, Choi KH, Park KH, Park CS, Cha J (2010) Characterization of an exo-acting intracellular alpha-amylase from the hyperthermophilic bacterium *Thermotoga neapolitana*. Appl Microbiol Biotechnol 86(2):555-66
- Poornima R, Sahu MK, Sivakumar K, Pushpavalli V (2008) Optimization of α-amylase production by *Actinomycete* strain AE-19 isolated from shrimp pond. Trends Appl Sci Res 3:45-52
- Rao MB, Tanksale AM, Ghatge MS, Deshpande VV (1998) Molecular and Biotechnological aspects of Microbial proteases. Microbiol Mol Biol Rev 62:597-635
- Roohi, Kuddus M, Ahmad IZ, Arif JM (2011) Production of coldactive extracellular α-amylase by newly isolated *Microbacterium foliorum* GA2 from Gangotri glacier, Western Himalaya, India. Asian J Biotech 3(5):449-459
- Russell NJ (1997) Psychrophilic bacteria: molecular adaptations of membrane lipids. Comp Biochem Physiol 118:489-493
- Saxena KR, Dutt K, Agarwal L, Nayyar P (2007) A highly thermostable alkaline amylase from a *Bacillus* sp. PN5. Biores Technol 98:260-265
- Sindhu R, Suprabha GN, Shashidhar S (2011) Purification and characterization of  $\alpha$ -amylase from *Penicillium janthinellum* (NCIM 4960) and its application in detergent industry. Biotechnol Bioinf Bioeng 1(1):25-32
- Smith MR, Zahnley JC (2005) Production of amylase by Arthrobacter psychrolactophilus. J Ind Microbiol Biotechnol 32:277-283
- Swain MR, Kar S, Padmaja G, Ray CR (2006) Partial characterization and optimization of production of extracellular α-amylase from *Bacillus subtilis* isolated from culturable cowdung microflora. Pol J Microbiol 55:289-296
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. Mol Biol Evol 24:1596-1599
- Uhlig H (1998) Industrial enzymes and their applications. Dissertation, University of California
- Vishwanathan P, Surlikar NR (2001) Production of α-amylase with *Aspergillus flavus* on Amaranthus grains by solid-state fermentation. J Basic Microbiol 41:57-64
- Wijbenga DJ, Beldman G, Veen A, Binnema DL (1991) Production of native-starch-degrading enzymes by a *Bacillus firmusflentus* strain. Appl Microbiol Biotech 35:180-184
- Zhang JW, Zeng RY (2008) Purification and characterization of a cold-adapted α-amylase produced by *Nocardiopsis* sp. 7326 isolated from Prydz Bay, Antarctic. Mar Biotech 10:75-82