

Secreted protease and activity of an endoproteinase for preliminary characterization of clinical *Candida* isolates

Francisca I Okungbowa, Rukhsana Chowdhury, Kalidas Pal

Received: 23 September 2009 / Received in revised form: 29 September 2009, Accepted: 15 July 2010, Published online: 25 July 2010
© Sevas Educational Society 2008-2010

Abstract

Nine clinical *Candida* strains (*Candida dubliniensis* (one), *Candida albicans* (six), and *C. tropicalis* (two)) were characterized by production of secreted protease, activity of endoproteinase A and with the HiCandida identification kit. Protease secretion was in the range of 1.3-1.8 cm while PrA activity was between 0.19 and 0.44 µg/min/mg protein. Results are indicative of the possible use of endoproteinase activity and SAP production for characterizing *Candida* strains.

Keywords: Biochemical activity, *Candida*, characterization, endoproteinase, secreted protease.

Introduction

Candida species are the most commonly isolated human opportunistic pathogens, especially in immunosuppressed individuals such as people infected with the Human Immunodeficiency Virus (HIV), those using in-dwelling catheters and those with prolonged chemotherapy (Santos *et al.*, 2002; Ahmad *et al.*, 2004; Kuhn *et al.*, 2004).

This study was undertaken with the aim of determining the possible use of secreted protease and activity of an endoproteinase for further characterizing nine clinical *Candida* isolates and to document additional data on *Candida* diversity.

Francisca I Okungbowa

Department of Plant Biology and Biotechnology (formerly Botany),
University of Benin, PMB 1154, Benin City, Nigeria

*Tel: 007098805057, Email: fiokun2002@yahoo.com

Rukhsana Chowdhury, Kalidas Pal

Infectious Diseases Division, Indian Institute of Chemical Biology,
Kolkata 700032, India.

MATERIALS AND METHODS

***Candida* strains:** Nine *Candida* strains isolated from genital samples (high vaginal swabs and urethra swabs) were obtained from the University of Benin Teaching Hospital (UBTH), Benin City, Nigeria. The *Candida* isolates were grown on Yeast Peptone Dextrose, YPD Agar (0.5% yeast extract, 1% peptone, 2% glucose and 2% agar powder, BD France) with added chloramphenicol (50 µg/ml) at 37°C for 48 hours.

The strains arbitrarily numbered 1-9 were identified as *C. dubliniensis* (isolate 1), *C. albicans* (isolates 2, 3, 5, 7, 8 and 9), *C. tropicalis* (isolates 4, 6). Pure cultures were maintained on YPD slants and stored at 0-4°C for future use.

***Saccharomyces cerevisiae*:** The strain used as control was a diploid standard laboratory strain, a gift from Dr. Anil Ghosh of the Department of Biotechnology, Indian Institute of Chemical Biology, Kolkata. Secreted protease and protease activity were determined according to Santos *et al* (2002) and Lowry *et al* (1951) respectively.

RESULTS

Secreted protease (SAP) was detected in all the nine *Candida* isolates. The diameters of halo (clear zone) surrounding inoculum were in the range of 1.3-1.8 cm (Fig 1). The production of SAP




Figure 1: Effect of diameter of halo on nine *Candida* isolates

was not detected in the *S. cerevisiae* strain. Activity of protease A (PrA) was observed in all nine *Candida* isolates, ranging from 0.19-0.35 $\mu\text{g}/\text{min}/\text{mg}$ protein for 15 minutes, and 0.2-0.44 for 30 minutes incubation (Fig 2). Enzyme activity was generally higher at 30°C. There were significant differences in PrA activities of isolates 1 and 2 and other isolates (at 15min incubation) and between those of 3 and 9 and others (at 30min incubation). Enzyme activity was significantly high for isolate 8 (30min incubation). Also, there was significant difference between values obtained for 15 minutes and 30 minutes incubation only for isolates 2, 8 and 9. With the HiCandida kit, six *C. albicans*, two *C. tropicalis* and one *C. dubliniensis* strains were identified. However, further separations were observed using SAP and PrA characterization, producing 2, 3 subgroups (*C. albicans*) and 0, 2 subgroups (*C. tropicalis*), respectively (Table 1).

169




Figure 2: Effect of enzyme activity on nine *Candida* isolates

Discussion and conclusions

Differences in proteolytic activity against albumin by SAP in *Candida* strains, have been used earlier to differentiate them into species and biotypes (Krzeminska – Jaskowiak *et al.*, 1994; Vidotto *et al.*, 2004). This is the first report of the use of an endoproteinase

Table 1: Speciation of nine *Candida* isolates by HiCandida Kit, SAP and PrA

Candida species	Number of same species identified by			
	Hi-Candida	SAP	PrA (15 min)	PrA (30 min)
<i>C. albicans</i>	6	5+1	(1) + (2) + (3)	(2) + (2) + (2)
<i>C. tropicalis</i>	2	2	(1) + (1)	(1) + (1)
<i>C. dubliniensis</i>	1	1	1	1

for characterizing *Candida* spp. There is the possibility of using the combination of SAP and PrA to characterize *Candida* species.

Acknowledgment

Authors are grateful to Third World Academy of Science (TWAS), Italy, and Council of Scientific and Industrial Research (CSIR), India, for fellowship to F. I. Okungbowa.

References

Ahmad S, Khan Z, Mokaddas E, Khan ZU (2004) Isolation and molecular identification of *Candida dubliniensis* from non-

human immuno-deficiency virus-infected patients in Kuwait. J Med Microbiol 53: 633-637

Krzeminska – Jaskowiak E, Pietkiewicz K, Ozdowska B (1994) Utilization of enzymatic activity in strains of *Candida* isolated from the vagina for typing and evaluation of pathogenicity. Med Dosw Mikrobiol 46: 225 – 231

Kuhn DM, Mukherjee PK, Clark TA, Pujol C *et al* (2004) *Candida parapsilopsis* characterization in an outbreak setting. Emerg Infect Dis 10: 1074 – 1081

Lowry OH, Rosebrough N.J, Farr AL, Randall RJ (1951) Protein Measurement with the Folin Phenol Reagent. J Biol Chem 193: 269-275

Santos AA, Sa EAC, Gaziri LCJ, Felli I (2002) Treatment of serum with supernatants from cultures of *Candida albicans* reduces its serum dependent phagocytosis. Braz J Microbiol 33: 79 – 83

Vidotto V, Ponton J, Aoki S *et al* (2004) Differences in extracellular enzymatic activity between *Candida dubliniensis* and *Candida albicans* isolates. Rev Iberoam Micol 21: 70 – 74.