

Polyphenolic Compounds of Plant of *Lepidium Ruderale* Linn. and Their Biological Activity

Shertayeva Nailya, Sabiraliyeva Zhanat, Taubayeva Raushan, Taithibekov Asker

Received: 18 August 2018 / Received in revised form: 18 December 2018, Accepted: 19 December 2018, Published online: 21 December 2018

© Biochemical Technology Society 2014-2018

© Sevas Educational Society 2008

Abstract

This article presents the results of the study of the plant *Lepidium ruderales* L. A method was developed for the complete extraction of polyphenolic substances from the plant, using the method of high-performance liquid chromatography flavonoids were identified and determined in the plant: kaempferol, quercetin; using mass spectrometry method, isovitexin belonging to the class C of flavonoids was identified; the chelating activity of the plant *Lepidium ruderales* L. was studied and it was revealed that the plant has a high biological activity; the content of organic acids was determined by capillary electrophoresis.

Keywords: Plants *Lepidium Ruderale* L., Flavanoids, Mass Spectrometry, High-Performance Liquid Chromatography Method, Astragalin, Luteolin, Kaempferol, Quercetin, Biological Activity, Capillary Electrophoresis Method, Organic Acids.

Shertayeva Nailya

Candidate of Chemical Sciences, Associate Professor, Department of chemistry and methods of teaching chemistry, Faculty of Chemistry and Biology, University of Taraz State Pedagogical University, Taraz City, Kazakhstan

Sabiraliyeva Zhanat

Candidate of Technical Sciences, Department of chemistry and methods of teaching chemistry, Faculty of Chemistry and Biology, University of Taraz State Pedagogical University, Taraz City, Kazakhstan

Taubayeva Raushan

PhD, Candidate of Agricultural Sciences, Department of chemistry and methods of teaching chemistry, Faculty of Chemistry and Biology, University of Taraz State Pedagogical University, Taraz City, Kazakhstan

Taithibekov Asker

Associate Professor, Department of chemistry and methods of teaching chemistry, Faculty of Chemistry and Biology, University of Taraz State Pedagogical University, Taraz City, Kazakhstan Country.

*Email: galiya008@mail.ru

Introduction

Lepidium ruderales L., which is known as Ban Helem in the Himalayas, is used as food and medicinal plant. It is used for treatment of cough, asthma, hemorrhoid, skin diseases, and rheumatism by the tribal inhabitants of the central Himalaya (Baoliang et al., 2003).

Recently, plant materials or their extracts as substitute and supportive medicine have been used increasingly, and several antimicrobial agents have been discovered and identified (Qusti et al., 2018). In the current study, *Lepidium ruderales* L. was tested for various biological activities like antifungal, antibacterial and antiviral agent. Benzyl glucosinolates, the principled elements of *Lepidium ruderales* L., showed anti-amebic activity. Quercetin and its derivatives were discovered in the plant *Lepidium ruderales* L. Quercetin – a plant flavanol, belonging to the group of catechins, the most abundant element in fruits and vegetables, has a higher antioxidant potential than vitamins C and E and a great biological activity (Jyota et al., 2003; Abu Zaker Khaled, 2001). In Kazakhstan, 20 species of peppergrasses are found (Baitenov, 2001).

Medicinal plants have been anticipated to be one of the most valuable resources in therapeutic practices for human diseases (Sargia et al., 2018). The use of herbal medicine represents a long history of human body (Sayed Ahmad et al., 2018). Historically, herbal medicines and their derivatives have demonstrated an important alternative for the treatment of illnesses (Nadri et al., 2018). Therefore, recently, there have been a lot of studies conducted on examining the contents of various plants. El Abidine Ababsa et al., (2018) in a study examined the chemical characterization and biological study of the species *Senecio Cineraria*.

The purpose of this study: Studying the polyphenolic compounds of the plant *Lepidium ruderales* L. and determining their biological activity.

Research object: the plant *Lepidium ruderales* L. (weedy *Lepidium*), collected in the Pavlodar region, in the nature

reservation territory of the Bayan-aul in the Republic of Kazakhstan.

In the course of this study, an optimal method was developed for extracting polyphenolic compounds from the overground part of *Lepidium ruderales* L., wherein 3 different methods for extracting substances were utilized: hot water, ethyl alcohol and methanol.

Table 1: The total content of polyphenolic compounds in different extract methods

Plant	Metanol	Etanol	Water
	The total amount of phenolic compounds, mg GAE / 100 g	The total amount of phenolic compounds, mg GAE / 100 g	The total amount of phenolic compounds, mg GAE / 100 g
<i>Lepidium ruderales</i> L.	71,4	51,9	56,3

As can be seen from table 1, the most complete extraction is achieved by using extraction with methanol.

Figure 1 shows the absorption spectra of the obtained extracts from *Lepidium ruderales* L.




Fig. 1- Absorption spectra of different extracts: 1- spectrum of methanol extract, 2- spectrum of ethanol extract, 3 - spectrum of water extract of a plant

Polyphenolic compounds were separated using high-efficiency liquid chromatography. For the separation efficiency, the corresponding column Zorbax SB C18 (3.5 μ m) 3x150 mm was chosen, the mobile phase: methanol - acetic acid solution 0.01% (25: 75); speed of the mobile phase: 1.2 cm³ / min; column temperature: 250 0C; detection under UV, λ = 254 nm, sample volume is 20 mm³. The chromatogram of flavonoids illustrated in Figure 2, the retention time indicated in Table 3.




Fig. 2- Chromatogram of flavonoids of the plant *Lepidium ruderales* L

Table 2: Quantitative content of flavonoids

Plant	Astragalin, g/kg	Luteolin, g/kg	Quercetin, g/kg	Kaempferol, g/kg
<i>Lepidium ruderales</i>	1,03	1,32	3,04	0,63
Total content	6,02			

Table 3: Identification of flavonoids *Lepidium ruderales* L.

Substance name	Retention time	Formula
Quercetin	3,005	<chem>O=C1C(=C(O)C(=C(O)C1=O)C2=CC(=C(O)C=C2)O</chem>
Kaempferol	3,763	<chem>O=C1C(=C(O)C(=C(O)C1=O)C2=CC=CC=C2</chem>
Astragalin	8,081	<chem>O=C1C(=C(O)C(=C(O)C1=O)C2=CC=CC=C2[C@@H]3O[C@H](CO)[C@@H](O)[C@H]3O</chem>
Luteolin	9,306	<chem>O=C1C(=C(O)C(=C(O)C1=O)C2=CC(=C(O)C=C2)O</chem>

To identify substances not identified by HPLC, mass spectrometry was used. For the determination of flavonoids by mass spectrometry, the method (Sheel Sharma, Nidhi Agarwal, 2011) was used. The results are shown in Figure 3.




Fig. 3 - Isovitexin mass spectra

Conditions for mass spectrometry analysis: ionization - negative, cell temperature - 1200C, source temperature - 5000C, column temperature - 400C, sample injected volume - 5 ml, analysis time -10 min (table 4).

It was found that the methanolic extract of the plant has the greatest chelating activity - 77.7%, and aqueous extract has a little less - 61.5%.

Table 4: Identification of flavanoids *Lepidium ruderales* L.

Substance name	retention time	formula
Isovitexin	4,07	

A study of the chelating properties of weedy *Lepidium* was conducted and it was found that different extracts from the plant exhibit different chelating ability, the data are shown in Table 5.

Capillary electrophoresis was used to determine the organic acids and 4 organic acids were detected in the extract, which were separated on a 55 cm long capillary with an inner diameter of 0.25 µm, the results of the study are shown in Table 6.

Table 6- Organic acid content

Substance	Contents, g/kg
Tartaric acid	0,4
Malic acid	0,7
Citric acid	1,2
Succinic acid	0,9
Total content of organic acids- 3,2	

Table 5: Chelating activity of various extracts of the overground part of weedy *Lepidium*

Plant	Extract	Concentration, µg/mg	Chelating activity (%)
<i>Lepidium ruderales</i> L.	Methanol	50	33,2
		100	45,6
		250	62,3
		500	77,7
	Ethanol	50	15,2
		100	25,9
		250	33,9
		500	51,5
	Water	50	23,1
		100	33,4
		250	46,7
		500	61,5

Findings: in the plant *Lepidium ruderales* L. during the study:

- a method was developed for the complete extraction of polyphenolic substances from a plant, and in this way it was determined that, extraction with methanol is 1.5–2 times more efficient than extraction with ethyl alcohol;
- using the high-performance liquid chromatography method in the plant, 6.02 g / kg of flavonoids were identified and determined including: astragaline - 1.03 g / kg; luteolin-1.32 g / kg; kaempferol-0.63 g / kg and quercetin-3.04 g / kg;
- by using mass spectrometry, Isovitexin belonging to the class C of flavonoids was identified;
- the chelating activity of the plant *Lepidium ruderales* L. was studied, and the plant was found to have a high biological activity of 77.7% in methanolic extract and 61.5% in an aqueous extract;
- by using capillary electrophoresis the organic acid content of 3.2 g / kg was determine: tartaric acid - 0.4 g / kg; malic

acid — 0.7 g / kg; citric acid - 1.2g / kg; succinic acid - 0.9 g / kg.

References

- Abu Zaker Khaled, Zhuravlev N.S. Quantitative determination of flavonoids in the leaves of some species of the genus *Rume L.* // National Pharmaceutical Academy of Ukraine, 2001. –C. 67-72.
- B. Sargia, B. Singh, N. Gupta, LK Gahlot, T. Gulati, Y. Hasija. MED-PDB: An online database of medicinal plants. *J Adv Pharm Edu Res* 2018;7(4):204-207.
- Baitenov M.S. *Flora of Kazakhstan*. Almaty, 2001.- T. 4 - 320 p.
- Baoliang Cui, Bolin Zheng, Kan He and Qun Yi Zheng. Imidazole Alkaloids from *Lepidium meyenii*.// *Jornal of Natural Products*, vol.66, No 8,2003.- p.1101-1103
- El Abidine Ababsal, Z. Kara Ali,W. Abidli,N. Akkal,S. & Jyota Agarwal and D.L. Verma. Quercetin Glycosides from Antioxidative Active aqueous Ethanolic Extract of *Lepidium ruderales L.*// *Academia Arena*, 3(3) 2011.- p.25-33
- Medjroubi,K. 2018. Chemical Characterization and Biological Study of the Species *Senecio Cineraria*. *World Journal of Environmental Biosciences*. 7 (3), 112-121.
- Nadri, S. Mahmoudvand, H. Mahmoudvand, H. Maryam Rashnoo,M. & Khaksarian,M. Chemical composition, antinociceptive and acute toxicity of *Pistacia atlantica* fruit extract. *Entomol Appl Sci Lett*, 2018, 5 (3): 8-12.
- Qusti, S.Y. Alseeni, M.N. Alharbi, R.A. Balgoon,M. Jambi, E.J. Alotaibi, S.A. (2018).Antibacterial activity of selected plants species extract cited in the holy Quran against clinical isolates. *Pharmacophore*, 8(5), 18-28.
- Sayed Ahmad,M. Shawky,A. Othman Ghobashy,M. & Ahmed Felifel, R.H. Effect of Some medicinal plants on life cycle of Citrus Brown Mites (*Eutetranychusorientalis*). *International Journal of Pharmaceutical Research & Allied Sciences*, 2018, 7(4):13-17.
- Sheel Sharma, Nidhi Agarwal. Nourishing and healing prowess of garden cress (*Lepidium sativum* Linn.), *Indian journal of natural products and resources*, Vol. 2 (3), September 2011, pp. 292-297.