

# Improvement of anaerobic production of hydrogen in the dark by genetic mutation strains of *Synechocystis* sp. strain PCC 6803

Takashi Yamamoto, Adipa Chongsuksantikul, Kazuhiro Asami, Kazuhisa Ohtaguchi\*

Received: 24 June 2012 /Received in revised form: 10 August 2012, Accepted: 10 August 2012, Published online: 11 June 2013  
© Sevas Educational Society 2008-2013

## Abstract

The anaerobic H<sub>2</sub> production in the dark by bidirectional hydrogenase of unicellular cyanobacterium *Synechocystis* sp. strain PCC 6803 has been studied. D-glucose addition to cell suspension has been known to enhance H<sub>2</sub> production, and the two mutants, glucose tolerant strain and its mutant lacking L-lactate dehydrogenase (LDH) gene, were introduced to improve H<sub>2</sub> production with D-glucose. Higher D-glucose uptake rate of glucose tolerant strain resulted in higher H<sub>2</sub> production rate comparing with the wild type strain. The LDH mutant showed higher initial H<sub>2</sub> production rate and H<sub>2</sub> yield from D-glucose than the other strains because the NADH which should be consumed originally by LDH can be redistributed to the H<sub>2</sub> production reaction.

**Keywords:** H<sub>2</sub> production, Bidirectional hydrogenase, *Synechocystis* sp. strain PCC 6803, D-glucose, Metabolic modification

## Introduction

Molecular hydrogen (H<sub>2</sub>) is a potential source of a non-polluting fuel. Cyanobacterial H<sub>2</sub> production is one of the candidates of sustainable H<sub>2</sub> production method. The unicellular cyanobacterium *Synechocystis* sp. strain PCC 6803 is a favored microorganism for the molecular biotechnological studies on the cyanobacterial H<sub>2</sub> production. The bidirectional hydrogenase of *Synechocystis* sp. strain PCC 6803 works as an enzyme forming H<sub>2</sub> and NAD(P)<sup>+</sup> from protons and NAD(P)H (Appel and Schulz 1996; Schmitz *et al.* 1995). This reaction is thought to play an important role for proper redox poisoning (Appel *et al.* 2000).

Some reports propose that more supply of NAD(P)H to hydrogenase improves H<sub>2</sub> production (Yamamoto *et al.*, 2012; Courmac *et al.*,

2004). Due to the oxygen sensitivity of hydrogenase photosynthesis inhibits its activity, H<sub>2</sub> production is performed under the dark and anaerobic condition. Under the condition NAD(P)H is produced by catabolism of intracellular glucan and cell constituting materials accumulated during photosynthetic growth. Supply of NAD(P)H is a key factor to improve H<sub>2</sub> production. D-glucose addition can also provide NAD(P)H source to cells even under the dark condition because *Synechocystis* sp. strain PCC 6803 has characteristic in that it is capable to consume D-glucose (Smith 1983). Catabolism of D-glucose to CO<sub>2</sub>, organic acids or other metabolites donates electrons to NAD(P)<sup>+</sup> to form NAD(P)H and improves H<sub>2</sub> production.

In the present report two genetic mutation strains of *Synechocystis* sp. strain PCC 6803 (hereafter PCC strain) were introduced to improve H<sub>2</sub> production. There are some advantages of using *Synechocystis* sp. strain PCC 6803 as a model microorganism, for instance genome DNA sequence was completely annotated on this strain (Kaneko *et al.* 1996) allowing us to construct mutant. Glucose tolerant strain of *Synechocystis* sp. strain PCC 6803 (hereafter GT strain) has been developed to grow on heterotrophic (Williams 1998). It seems that this strain has a high glucokinase activity which is key enzyme to utilize D-glucose (Kahlon *et al.* 2006). L-lactate dehydrogenase knock out mutant (hereafter *Δldh* strain) was constructed from GT strain. L-lactic acid is produced as one of the final products from D-glucose under the dark and anaerobic condition. The reaction from pyruvate to L-lactic acid which catalyzed by L-lactate dehydrogenase competes NADH consumption with hydrogenase, which results in decreasing of H<sub>2</sub> yield from D-glucose. Similar research has been studied by D-lactate dehydrogenase knock out mutant of *Synechococcus* sp. strain PCC 7002 (McNeely *et al.* 2010). This redirection of metabolic pathway succeeded to increase H<sub>2</sub> production by 5 folds.

## Methods

### Construction of *Δldh* mutant

The deletion of L-lactate dehydrogenase gene (slr 1556) was constructed by a PCR amplification of an *ldh* gene fragment using primer *ldh* forward (5'-GCCTATGATCGTCAATTTTCC-3') and *ldh* reverse (5'-TTCAGCAATATTTGCCAGTGTC-3') designed by us from the information of the chromosomal DNA of wild-type

Takashi YAMAMOTO\*, Adipa CHONGSUKSANTIKUL,  
Kazuhiro ASAMI, Kazuhisa OHTAGUCHI

Department of Chemical Engineering, Tokyo Institute of Technology, 12-1, Ookayama 2, Meguro-Ku, Tokyo 152-8552, Japan

\*Tel.: 0081 3 5734 2113; Fax: 0081 3 5734 2113.  
E-mail: paradox4@vesta.ocn.ne.jp

*Synechocystis* sp. strain PCC 6803 (Kaneko *et al.* 1996). PCR-amplified *ldh* fragment was ligated into *Sma*I restriction site of plasmid pUC19. The kanamycin cassette from plasmid pUC4K was inserted into the *Bgl*II restriction site at *ldh* gene fragment. Transformant colony was selected from BG-11 agar plate containing kanamycin and continuously re-streaked to new agar plates. Kanamycin resistant cell were grown at increasing concentrations to 120  $\mu\text{g mL}^{-1}$  and finally transferred into liquid BG-11 medium. The full segregation was confirmed by PCR (Figure S1) using the mentioned primer. Segregated cells maintained in 5 mL BG-11 medium containing 60  $\mu\text{g mL}^{-1}$  of kanamycin and 100  $\mu\text{L}$  of culture transfer to flesh medium every two weeks.

#### Cell preparation

*Synechocystis* sp. strain PCC 6803 was photosynthetically grown at 34 °C in a 100 mL bubble column (clear Pyrex glass) containing 80 mL HEPES buffer (pH 7.7) aqueous solution of BG-11 medium. The aqueous solution was aerated at 80 mL  $\text{min}^{-1}$  by the air containing 6%  $\text{CO}_2$  that was filtered through 0.45  $\mu\text{m}$  filter and fed from the bottom of the bubble column. The bubble column was placed in a 34 °C water bath. One side of the bubble column was illuminated by fluorescent lamps at 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . After the 24 h cultivation cells were harvested by centrifugation at 25 °C and 3000 rpm for 10 min. The cell pellets were washed with HEPES buffer solutions and centrifuged again.

#### $\text{H}_2$ production

After the above cell preparation, the settled cells were suspended in de-ionized water containing 50  $\text{mmol L}^{-1}$  HEPES buffer (pH 7.7) in a test tube. Initial cell mass concentration for  $\text{H}_2$  production was adjusted to 2.0  $\text{g L}^{-1}$ . Total volume of cell suspension was adjusted to 10 mL. Finally cell suspensions in the test tubes were sparged with  $\text{N}_2$  gas for a few minutes to remove  $\text{O}_2$  gas and sealed with butyl rubber caps. Test tubes were placed into a water bath at 34 °C under a dark condition and agitated at 120 rpm.

#### Measurement of $\text{H}_2$ production, cell mass concentration, D-glucose concentration in cell suspension and hydrogenase activity

At the measurement time, 300  $\mu\text{L}$  of gas sample was withdrawn from the test tube with a gasket syringe and 400  $\mu\text{L}$  of cell suspension was also withdrawn from the test tube with a syringe. After that, the test tube was sparged with  $\text{N}_2$  gas again.

An amount of  $\text{H}_2$  in gas sample was determined with a gas chromatograph and a thermal conductivity detector (GC-320, GL science Inc.; column, Molecular sieve 13X; carrier gas, nitrogen gas; column temperature, 315 K; injector temperature, 333 K; detector temperature, 353 K). Total  $\text{H}_2$  production per unit suspension volume of a run ( $y_{\text{H}_2}$ ) was calculated from a result.

Cell mass concentration in terms of dry cell weight (DCW) per unit suspension volume was monitored by measuring an optical density at 730 nm ( $\text{OD}_{730}$ ) utilizing a spectrophotometer. One unit absorbance of cell suspension was equivalent to 0.369  $\text{g-DCW L}^{-1}$ .

The amount of D-glucose contained in a reaction mixture ( $c_G$ ) was determined with a glucose tester (Wako Pure Chemicals, Japan) as mentioned elsewhere (Yamamoto, 2012).

The activity of hydrogenase in intact cells of *Synechocystis* sp. strain PCC 6803 was assayed under methyl viologen by the modified method of Gutthann *et al.* (2007). First, 4 mg of cells harvested from photobioreactor were suspended in a 2 mL of de-ionized water

containing 50  $\text{mmol L}^{-1}$  HEPES, 5  $\text{mmol L}^{-1}$  methyl viologen and 10  $\text{mmol L}^{-1}$  sodium hydrosulfite in 15 mL glass tubes. Then the cell suspensions in the test tubes were sparged with  $\text{N}_2$  gas for a few minutes to remove dissolved and gaseous  $\text{O}_2$  and sealed with butyl rubber caps. The test tubes were placed into a water bath at 34 °C under a dark condition and agitated at 120 rpm. Produced  $\text{H}_2$  in the gas phase of the test tubes were measured three times every 30 min and hydrogenase activity was calculated. Reproducibility was confirmed by 2 – 4 runs for each result.

## Results and discussion

#### Growth

PCC, GT and *Aldh* strain were autotrophically grown in BG-11 medium. No differences of specific growth rate in growth phase and final cell mass concentration in stationary phase of each strain were observed (data not shown). During cell preparation each strain did not secrete metabolites with detectable level indicating that L-lactate dehydrogenase did not work on growth phase, and *Aldh* strain has a growth characteristic equal to wild type.

#### Hydrogenase activity

Specific hydrogenase activities of each strain were measured at the beginning of  $\text{H}_2$  production phase. Their values were around 4.8  $\text{U g}^{-1}$ . Hydrogenase seems not to be affected by knock out of L-lactate dehydrogenase gene.

#### $\text{H}_2$ production

Figure 1 shows  $\text{H}_2$  production by each strain in a HEPES buffer solution or buffer solutions containing 5.6, 28 or 56  $\text{mmol L}^{-1}$  of D-glucose at 96 h. Cells completely consumed extracellular D-glucose in 48 h in the runs of each strain with 5.6  $\text{mmol L}^{-1}$  of D-glucose. On the other hands the greater part of D-glucose remained in cell suspension at 96 h in the runs with 28 or 56  $\text{mmol L}^{-1}$  of D-glucose. D-glucose improved  $\text{H}_2$  production for each strain. Optimum D-glucose concentrations were 5.6 or 28  $\text{mmol L}^{-1}$  of D-glucose. Excess D-glucose concentration inhibited  $\text{H}_2$  production. It seems that cells may use energy to drive sugar out of cells at high D-glucose concentration (Baebprasert *et al.*, 2010). Comparing with PCC strain, GT strain supplemented with D-glucose produced more  $\text{H}_2$ . The amount of  $\text{H}_2$  production in the runs with 5.6, 28 or 56  $\text{mmol L}^{-1}$  of D-glucose increased 13, 15 and 30%, respectively. On the other hands the results by *Aldh* strain not only with D-glucose but without D-glucose showed increasing of  $\text{H}_2$  production.




Figure 1: Hydrogen production per unit volume of cell suspension by PCC, GT and *Aldh* strain in buffer solutions containing 0, 5.6, 28 and 56  $\text{mmol L}^{-1}$  of D-glucose at 96 h.

Especially initial  $\text{H}_2$  production rates of *Aldh* strain were improved in the each run (Figure 2). This implies the mutant cells contained

more reducing materials for H<sub>2</sub> production than the other two strains at the beginning of H<sub>2</sub> production phase because less D-glucose was catabolized at that time.

#### L-lactic acid production

Figure 3 shows L-lactic acid production at 96 h. Before 72 h, *Aldh* strain did not produce L-lactic acid as detectable level (data not shown), which ensures that metabolic modification was successfully performed. Decreasing of L-lactic acid production seems to contribute to the increase of H<sub>2</sub> production by *Aldh* strain because more NADH would be redistributed to hydrogenase. Little amount of L-lactic acid was observed in the run with *Aldh* strain at 96 h. This might be because some mutant cells might turn back to wild type during cell preparation and H<sub>2</sub> production phase without kanamycin. Comparing with PCC and *Aldh* strain, H<sub>2</sub> production did not reach to the level, which is estimated that inability of 1 mol L-lactic acid production theoretically results in the raise of 1 mol H<sub>2</sub>. It seems that other NADH consumption pathway also competed with H<sub>2</sub> production.

#### D-glucose uptake rate

The improvement of H<sub>2</sub> production by GT strain resulted from increasing of D-glucose uptake rate. Figure 4 showed D-glucose uptake rate during 0-24 h in each run. Comparing with PCC strain,




Figure 2: Initial hydrogen production rate per unit volume of cell suspension by PCC, GT and *Aldh* strain in buffer solutions containing 0, 5.6, 28 and 56 mmol L<sup>-1</sup> of D-glucose.




Figure 3: Concentration of L-lactic acid produced by PCC, GT and *Aldh* strain in buffer solutions containing 0 or 56 mmol L<sup>-1</sup> of D-glucose at 96 h.

GT strain showed high D-glucose uptake rate. It is deduced that GT strain has high glucokinase activity and can utilize more D-glucose than PCC strain. On the other hand *Aldh* strain showed lower D-glucose uptake rate than PCC strain although this mutant was derived from GT strain. This might be because metabolic

modification affects D-glucose utilization characteristic. L-lactate dehydrogenase mainly works in cytoplasmic matrix; however,




Figure 4: Uptake rate of extracellular D-glucose per unit volume of cell suspension by PCC, GT and *Aldh* strain in buffer solutions containing 5.6, 28 and 56 mmol L<sup>-1</sup> of D-glucose. The uptake rates were calculated using data at 0 and 24 h for  $c_{G0} = 5.6$  mmol L<sup>-1</sup>, and at 0 and 96 h both for  $c_{G0} = 28$  and 56 mmol L<sup>-1</sup>.

hydrogenase exists on thylakoid membrane (Appel *et al.*, 2000). It is deduced that NADHs produced through glycolysis in cytoplasmic matrix of *Aldh* strain were locally accumulated in and inhibited D-glucose utilization.

#### H<sub>2</sub> yield from D-glucose

H<sub>2</sub> yield from extracellular D-glucose consumption ( $Y_{H2/G}$ ) is an important factor to consider a bioprocess. Extracellular D-glucose is utilized for producing cell constituents, intracellular glucan and secreted metabolites, then these processes produce or consume NAD(P)H. Knock out of L-lactate dehydrogenase in *Aldh* strain improved  $Y_{H2/G}$  as expected (Figure 5). It implies that more NADH derived from D-glucose consumed to produce H<sub>2</sub> in *Aldh* strain comparing with the other strains. Metabolic modification successfully increased the efficiency of D-glucose utilization. In the run with 5.6 mmol L<sup>-1</sup> of D-glucose, GT strain showed the highest yield among three strains. This is because H<sub>2</sub> produced after depletion of D-glucose was reflected on the data.




Figure 5: H<sub>2</sub> yields, calculated from extracellular D-glucose consumption during 0 and 96 h, of PCC, GT and *Aldh* strain in buffer solutions containing 5.6, 28 and 56 mmol L<sup>-1</sup> of D-glucose.

## Conclusion

D-glucose addition to a buffer solution in H<sub>2</sub> production phase improved H<sub>2</sub> production due to more supply of NAD(P)H to hydrogenase. D-glucose uptake rate and H<sub>2</sub> yield from D-glucose are important factors for H<sub>2</sub> production in extracellular D-glucose supply condition. In the present report, GT strain showed high D-glucose uptake rate. This contributed to the increase in H<sub>2</sub> production. On the other hand, *Aldh* strain was successfully constructed from GT strain and showed higher  $Y_{H2/G}$  values than

other strains. Furthermore initial H<sub>2</sub> production rate was also increased. H<sub>2</sub> production by *Aldh* strain did not reach to the upper-level estimated stoichiometrically as 1 mol increase in H<sub>2</sub> with 1 mol decrease in L-lactic acid in *Aldh* strain, implying that there is a room to improve H<sub>2</sub> production by further metabolic modification.

## Acknowledgement

I would like to thank Professor Yukako Hihara in Department of Biochemistry and Molecular Biology, Graduate School of Science and Engineering, Saitama University for kindly giving us a research material, *Synechocystis* sp. strain PCC 6803 GT.

## References

- Appel J, Phunpruch S, Steinmiller K, Schulz R (2000) The bidirectional hydrogenase of *Synechocystis* sp. PCC 6803 works as an electron valve during photosynthesis. *Arch Microbiol* 173:333-338
- Appel J, Schulz R. (1996) Sequence analysis of an operon of a NAD(P)-reducing nickel hydrogenase from the cyanobacterium *Synechocystis* sp. PCC 6803 gives additional evidence for direct coupling of the enzyme to NAD(P)H-dehydrogenase (complex I). *Biochim Biophys Acta* 1298:141-147
- Baebprasert W, Lindblad P, Incharoensakdi A (2010) Response of H<sub>2</sub> production and Hox-hydrogenase activity to external factors in the unicellular cyanobacterium *Synechocystis* sp. strain PCC 6803. *Int J Hydrog Energy* 35:6611-6616
- Cournac L, Geneviève G, Gilles P, Vignais PM (2004) Sustained photoevolution of molecular hydrogen in a mutant of *Synechocystis* sp. strain PCC 6803 deficient in the type I NADPH-dehydrogenase complex. *J Bacteriol* 186:1737-1746
- Gutthann F, Egert M, Marques A, Appel J (2007) Inhibition of respiration and nitrate assimilation enhances photohydrogen evolution under low oxygen concentrations in *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta* 1767:161-169
- Kahlon S, Beerli K, Ohkawa H, Hihara Y, Murik O, Suzuki I, Ogawa T, Kaplan A. (2006) A putative sensor kinase, Hik31, is involved in the response of *Synechocystis* sp. strain PCC 6803 to the presence of glucose. *Microbiol* 152:647-655
- Kaneko T, Sato S, Kotani H, Tanaka A, Asamizu E, Nakamura Y, Miyajima N, Hirose M, Sugiura M, Sasamoto S, Kimura T, Hosouchi T, Matsuno A, Muraki A, Nakazaki N, Naruo K, Okumura S, Shimpo S, Takeuchi C, Wada T, Watanabe A, Yamada M, Yasuda M, Tabata S (1996) Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res* 3:109-136
- McNeely K, Xu Y, Bennette N, Bryant DA, Dismukes GC (2010) Redirecting reductant flux into hydrogen production via metabolic engineering of fermentative carbon metabolism in a cyanobacterium. *Appl Environ Microbiol* 76:5032-5038
- Schmitz O, Boison G, Hilscher R, Hundeshagen B, Zimmer W, Lottspeich F, Bothe H (1995) Molecular biological analysis of a bidirectional hydrogenase from cyanobacteria. *Eur J Biochem* 233:266-276
- Smith AJ (1983) Modes of cyanobacterial carbon metabolism. *Ann Microbiol* 134:93-113
- Williams JGK (1998) Construction of specific mutations in photosystem II photosynthetic reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol* 167:766-778
- Yamamoto T, Asami K, Ohtaguchi K (2012) Anaerobic production of hydrogen in the dark by *Synechocystis* sp. strain PCC 6803:

effect of photosynthesis media for cell preparation. *J Biochem Tech* 4(1):464-468.