Study of the Thermodynamic Parameters for Interaction of Ciprofloxacin Antibiotic with Bulk and Nanocopper Sulfate

Elsayed M. AbouElleef* and Saleh D. Mekkey

Received: 28 October 2018 / Received in revised form: 092 March 2019, Accepted: 08 March 2019, Published online: 23 March 2019 © Biochemical Technology Society 2014-2019 © Society 2014-2019

© Sevas Educational Society 2008

Abstract

The association parameters of both bulk and nano-CuSO₄ salts in (EtOH-H₂O) mixed solvents were estimated at room temperature (293.15K) using conductivity method by applying Fuoss-Shedlovsky calculation. The association parameters of nano-CuSO₄ was greater than bulk CuSO₄ salt. The interaction between the cation (Cu²⁺) with ligand (ciprofloxacin, CFL) in (EtOH-H₂O) mixed solvents at room temperature (293.15K) was determined by the conduct metric method. The formation Gibbs free energies change (ΔG_f) and transfer Gibbs free energies change (ΔG_t) were increased in negative signs with increasing in EtOH content in the binary solvent. Two stoichiometric complexes 1:2 and 1:1 (M:L) are formed and the formation constant and Gibbs free energy of the formed complexes follow the order: $K_f(1:1) > K_f(1:2)$ and ΔG_f (1:1) > ΔG_f (1:2) for (M:L) (in negative values), which indicate more favorable formation of (1:1) complex compared with (1:2) complex. Also there are a decrease in values of Kf and $\Delta G_{\rm f}$ in case of using nano-CuSO₄ than that of using bulk-CuSO₄ but in small difference due to the more solvation effect in case of using nanoparticles.

Keywords: Copper Sulfate, Ciprofloxacin, Association Parameters, Complexation, Formation Constant, Gibbs Free Energy. Mixed Solvents.

Introduction

The investigation of the complexation formation reactions of the ligand with various cations in organic solvent is important for understanding the mechanism of biological transport, molecular recognition, and other analytical applications (Jose et al. 2009). Transition metal cations have a vital role in redox enzyme systems and bio-inorganic chemistry and also provide the basis of

Elsayed M. AbouElleef*

Chemistry Department, Faculty of Arts and Science, Northern Border University, 91911 Rafha, Kingdom of Saudi Arabia. Basic Science Department, Delta Higher Institute for Engineering & Technology, 35681-Dakhlia, Mansoura, Egypt.

Saleh D. Mekkey

Chemistry Department, Faculty of Arts and Science, Northern Border University, 91911 Rafha, Kingdom of Saudi Arabia. Department of Chemistry, Faculty of Science, Al-Azhar University, Cairo, Egypt.

*Email: s.abouelleef@yahoo.com

models biological systems (Angelusiu et al. 2010). It is helpful to consider the free energy, enthalpy and entropy change contributions to complexation reactions to obtain a good understanding of the thermodynamics of the complexation formation reactions (Gomaa et al. 2014a; Helmy et al. 2016).

The different techniques like polarography, spectrophotometry and conduct metric depends on physical and chemical properties of a material solution have been used to determine the complex formation between different metal cations within the ligand in the solutions (Nezhadali et al. 2009 & 2008). The conduct metric is an electrochemical technique, which is an extremely widespread and useful tool with a highly sensitive, simple and inexpensive experimental arrangement, that is used for studying the formation of complex between transition metal cations and ligand in solution (Harwood et al. 1989). It is very important to study complexation formation between different metal cations and ligand in a various solvent systems and their effect on solute's solvation by the solvent mixtures (Christopher et al. 2004; Nezhadali et al. 2011).

Thermodynamics parameters were used in understanding many systems and have vital importance for learning about the spontaneity of a given process at a certain temperature (Gomaa et al. 2013 a; Gomaa et al. 2013 b; El-Hady et al. 2012). Determination of stability constant is fundamental for understanding the behavior of the metal cations in a solution in the presence of some chelating agents and is best explained in terms of thermodynamics. Conductivity technique can be used to estimate the stability constant values, and also makes available the thermodynamics parameters of metal-ion complex formation.

Ciprofloxacin (generic name) is an antibiotic prescribed for treating a number of infections caused by susceptible bacteria including infections of bone, joint, lung, skin, intra-abdominal infections, certain type of infectious diarrhea, typhoid fever, and urinary tract infections, among others. It is used in addition to other antibiotics for some infections (Grabe et al. 2013; Baddour et al. 2005).

It is therefore necessary to study the effect of ciprofloxacin antibiotic on the properties of copper sulfate salt by determining the formation constant and the free energy change of interaction between them to find the extent of benefit of the ciprofloxacin antibiotic.

In this work, the association parameter for bulk and nano-copper sulphate salt in (EtOH-H₂O) mixed solvents at room temperature (293.15K) was determined by using the conduct metric method. The complex formation constant and stoichiometry of the complexation were elucidated between the metal cation (Cu²⁺) and the ligand (ciprofloxacin, CFL) in (EtOH-H₂O) mixed solvents at room temperature (293.15K) by using the conduct metric method to study the effect of the solvent mixture properties on the stabilities of the complexes.

Experimental

Materials and solutions

Copper sulfate salt (CuSO4.5H₂O) and ethanol were obtained from Merck as analytical reagent grade materials and used without further purification while ciprofloxacin antibiotic (CFL) was from ["SANDOZ" a Novartis company].



Figure 1. Ciprofloxacinhydrochloride antibiotic (CFL). Formula: C₁₇H₁₉ClFN₃O₃ Molar mass: 367.805 g/mol

The water and the binary mixed solvents of (EtOH-H₂O) 0%, 20%, 40%, 60%, and 80% EtOH were used as the solvents under the study and were prepared by mixing EtOH and H₂O (with error $\pm 0.01\%$).

Apparatus

The conductance measurements were carried out with a conduct meter (OAKTON WD-35607-10,-20,-30) with platinum electrode and a cell constant equal 0.99. The temperature was adjusted on an ultra-thermostat (Kottermann 4130) with a constant temperature maintained within $\pm 0.05^{\circ}$ C.

Procedure

The nano-CuSO₄ salt was prepared by using ball milling method, by shaking CuSO₄ salt in ball-milling apparatus of the type Retsch MM 2000 swing mill at 20225 Hz at room temperature for a period of one hour. The mill has 10 cm³ stainless steel double-walled tubes. Two balls made from stainless steel of 12 mm diameter were used.

Transmission electron microscopy (TEM) analysis

The TEM images of Nano copper sulphate sample in distilled water were measured using electron microscope model Joel 2010.

Preparation of solutions

Solutions of Cu^{2+} all at 10^{-3} M were prepared by dissolving the accurately weighed amount of salt in proper volume of binary solvents. 10^{-3} M solutions of ligand (CFL) were prepared by dissolving the accurately weighed amount of (CFL) in the proper volume of binary solvents.

Conduct metric measurement

-To calculate the association parameters of bulk and nano-CuSO₄.5H₂O in the (EtOH-H₂O) mixed solvent, a solution of metal cation (10^{-3} M, 20 mL) was placed in a conductance cell, and the conductance was measured. The conductance of solution was measured after each addition of the mixed solvent using a microburret after stirring. The conductance reading, taken after 1.0 min of each addition, was corrected for dilution.

-To calculate the formation constants between $CuSO_4.5H_2O$ and a ligand (Ciprofloxacin, CFL) in (EtOH-H₂O) mixed solvents, a solution of CuSO₄.5H₂O salt (10⁻³ M, 20 mL) was placed in a conductance cell, and the conductance was measured. The ligand (10⁻³ M) (CFL) was transferred step by step to the conductance cell using a micropipette and the conductance was measured after each addition. The temperature used is room temperature (293.15K).

Results and Discussion

X-ray diffraction (XRD)

X-ray diffraction of investigated bulk and nano-CuSO₄ samples where nano samples prepared by ball mill method are shown in Figures 2 & 3. This new peaks at 2θ = 15.508, 16.322, 18.734, 22.364, 24.102, 27.035, 29.225, 32.825, 36.757, 47.096, 49.815, 56.364, 60.717, 67.086 may be due to elimination of water molecules by ball mill method. The nano-CuSO₄.5H₂O is better crystallized than bulk-CuSO₄.5H₂O and new faces are formed as clear in X-ray figs (3). The observed peaks are related to triclinic structure of CuSO₄.5H₂O as indexed in the standard data (Varghese et al. 1985).

By using Scherrer's Formula, the crystalline sizes of the samples have been estimated.

$d_{XRD} = k\lambda/\beta \cos\theta$ (1)

Where k is a shape factor which is taken to be 0.9, λ is the used wavelength of x-rays used (1.15405 nm), β is the full width at half maximum and $\boldsymbol{\theta}$ is the angle of diffraction. The obtained crystalline sizes are found to be in the nanometer scale range. The average crystal size is obtained in Table (1).

Table 1: XRD data of bulk and nano-CuSO₄.5H₂O samples.

Sample	Crystal size d _{XRD} (nm)
Bulk CuSO ₄ .5H ₂ O	141.45
Nano CuSO ₄ .5H ₂ O	163.76







Figure 3: X-ray diffraction of nano-CuSO₄.5H₂O.

Scanning & Transmission electron microscope (SEM) & (TEM)

Transmission electron microscope (TEM) is a special kind of electron microscope for imaging of different objects. In TEM the electrons pass through with atoms of the sample and interact with the atoms and the electrons are therefore being scattered.

Their size and morphology of nano-CuSO₄.5H₂O are studied by TEM and CEM analysis. Fig. (4) shows the nanoparticles that obtained in ethanol are sphere like. The Nano particles are in the size ranging 5.027 - 49.33 nm. The all shapes are separated with defined boundaries. The diffraction image of TEM shown in Fig. (5) supported the X-ray diffraction; the nano-CuSO₄.5H₂O is better crystallized. The images are also indicated by SEM measurements in Fig. (6).



Figure 4: TEM images of nano-CuSO₄.5H₂O.



Figure 5: TEM diffraction image of nano-CuSO₄.5H₂O.



Figure 6: SEM images of nano-CuSO₄.5H₂O.

Calculation of association parameters for CuSO₄.5H₂O

The values of specific conductance (K_s) for different concentrations of CuSO₄.5H₂O solution in (EtOH-H₂O) mixed solvents were measured experimentally at room temperature (293.15 K). Also, the values of molar conductance (Λ_m) can be calculated using equation (2) (Gryzybkowski et al. 1980; Gomaa et al. 2014b):

$$\Lambda_m = \frac{(K_s - K_{solv})K_{cell} \times 1000}{C}$$
(2)

where, K_s and K_{solv} are the specific conductance of the solution and the solvent, respectively; K_{cell} is the cell constant, equal to 0.99 and C is the concentration of the CuSO₄.5H₂O solution in molar.

The limiting molar conductance (Λ_0) at infinite dilutions was estimated by extrapolating the relation between Λ_m and $C_m^{\nu_2}$ to zero concentration (Fig. 7).

The experimental data for conductance measurements were analyzed using Fuoss-Shedlovsky (Shdlovsky et al. 1956) extrapolation techniques which follows equation (3-12): -

$$\frac{1}{\Lambda S_{(z)}} = \frac{1}{\Lambda_o} + \left(\frac{K_A}{\Lambda_o^2}\right) \left(C\Lambda \gamma_{\pm}^2 S_{(z)}\right)$$
(3)

where, $S_{(Z)} = 1 + Z + Z^2/2 + Z^3/8 + \dots$ etc. and

$$Z = \frac{S(\Lambda C)^{1/2}}{\Lambda_o^{3/2}}$$
(4)

The Onsager slope (S) can be calculated by knowing the value of (Λ_0) from the equation (5):

$$S = a\Lambda_0 + b$$
 (5)
where, $a = 8.2 \times 10^5 / (\epsilon T)^{3/2}$ (6)
 $b = 82.4/n((\epsilon T)^{1/2}$ (7)

where, (ϵ) is the dielectric constant of the solvent, (η_0) is the viscosity of the a binary solvent and (T) is the temperature. Using the values of (ϵ) and (η_0), the value of (S) were easily estimated. Using the data of (Λ), S_(z) and (Λ_0), the degree of dissociation (α) was calculated by equation (8):

$$(\alpha) = \Lambda S_{(Z)} / \Lambda_0 \tag{8}$$

Using these (α) and (ϵ) values, the mean activity coefficients (γ_{\pm}) were calculated by equation (9) (Debye et al. 1923):

$$\log \gamma_{\pm} = -A(\alpha C)^{1/2} / [1 + B r^{o}(\alpha C)^{1/2})] (9)$$

where, (z^{\cdot}, z^{+}) are the charges of ions A & B are the Debye-Hückel constant.

 $A=1.824~X~10^6~(\epsilon T)^{-3/2}~$; $B=50.29~X~10^8~(\epsilon T)^{-1/2}~$ and (r^o) is the solvated radius.

The association constant (K_A) is given by equation (10):

$$K_{A} = \frac{C_{[MX_{n}]} \cdot \gamma_{[MX_{n}]}}{C_{M^{n+}} \cdot \gamma_{M^{n+}} \cdot C_{X^{-}}^{n} \cdot \gamma_{X^{-}}^{n}} (10)$$

Using the values of association constant (K_A), the values of the dissociation constant (K_D) were easily calculated by means of the following equation (11):

$$\mathbf{K}_{\mathrm{D}} = 1/\mathbf{K}_{\mathrm{A}} \tag{11}$$

The triple ion association constant (K_3) were calculated (Hirsch et al. 1956; Fuoss et al. 1959 a) by using the equation (12):

$$\frac{\Lambda C^{1/2}}{(1-\frac{\Lambda}{\Lambda_{o}})^{1/2}} = \frac{\Lambda_{o}}{(K_{A})^{1/2}} + \frac{\lambda_{3}^{o}C}{K_{3}(K_{A})^{1/2}}(1-\frac{\Lambda}{\Lambda_{o}})$$
(12)

Equation (12) was derived by Fuoss and using Walden approximation ($\Lambda_0=3\lambda_0$).

The values of free energy change of association (ΔG_A) of CuSO₄.5H₂O solution in (EtOH-H₂O) mixed solvents at room temperature (293.15 K) were calculated (Popuych et al. 1972; Isabel et al. 1996) from the association constant (K_A) values by using equation (13) (Atkins 1978; Ives 1971).

$$\Delta G_A = -2.303 \text{ RT} \log K_A \qquad (13)$$

From the calculated (ΔG_A) values, the free energies change of transfer (ΔG_t) from water (w) as a reference solvent to the (EtOH-H₂O) mixed solvent (s) were calculated using equation (14).

$$\Delta G_t = \Delta G_{A(s)} - \Delta G_{A(w)} \tag{14}$$

Where $\Delta G_{A(s)}$ and $\Delta G_{A(w)}$ are the free energies of association in mixed solvents and in water, receptively.

The Walden product $(\Lambda_0\eta)$ values (Walden, 1906; Kay, 1973; Kay, 1991; Kay et al., 1968 and Fuoss, 1959 b) were calculated for the solutions of CuSO₄.5H₂O in pure (EtOH) solvent and also

in (EtOH-H₂O) mixtures. The Walden product values as computed from the limiting molar conductance values (Λ_{o}) would be informative from the point of view of ion-solvent interactions.

The fluidity ratios (Bishopa et. al 1958) were also calculated by using equation (15).

$$R_{x} = \frac{\Lambda_{0} \eta_{0} (\text{in mixed solvent})}{\Lambda_{0} \eta_{0} (\text{in water})}$$
(15)

It was observed that the fluidity ratios (R_x) values were decreased with increasing the organic solvent (EtOH) content in the (EtOH-H₂O) mixtures; this is an indication of selective solvation of CuSO₄.5H₂O or ion by the organic solvent molecules in this mixed solvent and is ascribed mainly to the decrease in the Zwanzig solvent relaxation effect (Zwanzig, 1963). In the other words, this decrease in Walden product ($\Lambda_0\eta$) indicates strong solvent-solute interaction, as a result of which the solvent fails to decrease the mobility of ions and increases the viscosity of the medium. The strength of such interaction however increases with increase of the proportion of EtOH in (EtOH-H2O) mixtures. We see that Walden product is lower in water medium than in (EtOH-H2O) mixtures. This is explained by Frank and Wen (Franck et. al 1957) as follows: ions like quaternary ammonium salts that possess large hydrophobic side chains, a water molecule at the surface of these large ion is influenced very little by either the ionic change or the inert hydrocarbon chain on its one side. Consequently, the water molecules on the hydrophobic surface can be oriented to a greater extent by their nearest neighbors and can, in effect, be oriented into cages about the hydrocarbon side chains that one similar in structure to polyhedral clathrate hydrates of this ion (Beurskens, et. al 1964). This clathrate like structures can be considered as a type of hydration, since increase the size of the moving entity. Ions experiencing this effect should have lower mobilities in aqueous solution than in non-aqueous solution, since only water appears able to form any appreciable amount of three-dimensional structure.

The calculated values of association parameters and Gibbs free energies change for the solution of bulk and nano-CuSO₄.5H₂O in (EtOH-H₂O) mixed solvents at room temperature (293.15 K), and were calculated and reported in Tables 2 & 3 and illustrated in Fig. (8).



Figure 7: Variation of the molar conductance Λ_m (S.cm².mol⁻¹) with C^{1/2} for bulk CuSO₄.5H₂O in (EtOH-H₂O) mixed solvents at (293.15 K).

Table 2: The values of limiting molar conductance (Λ_0), degree of dissociation (α), activity coefficient (γ_{\pm}), dissociation constant (K_D), triple ion association constant (K₃), association constant (log K_A), Gibbs free energy change of association (ΔG_A), transfer free energy change (ΔG_t), Walden product ($\Lambda_0\eta$) and fluidity ratios (R_x) for bulk-CuSO₄.5H₂O in (EtOH-H₂O) mixed solvents at room temperature (293.15 K).

EtOH vol. %	Xs	Λo	α	γ_{\pm}	$10^3 K_D$	$10^{3} K_{3}$	$\log K_{\text{A}}$	ΔG_{A}	$\Delta \; G_t$	Λ οη	Rx
0	0	257.2	0.9209	0.9599	3.2953	1.8112	2.4821	-13.9320	0.0000	2.5761	1.0000
20	0.0716	207.3	0.7845	0.9572	0.8723	3.6223	3.0593	-16.0004	-2.0684	2.1022	0.8161
40	0.1706	135.6	0.7757	0.9592	0.8226	3.7044	3.0848	-17.3151	-3.3831	1.3988	0.5430
60	0.3163	101.6	0.6480	0.9583	0.3652	5.5745	3.4374	-19.2942	-5.3622	1.0732	0.4166
80	0.5523	48.2	0.5320	0.9514	0.1825	7.6645	3.7389	-20.9860	-7.0540	0.5281	0.2050
100	1	6.87	0.8452	0.7997	0.9835	1.3328	3.0072	-16.8795	-2.9475	0.0806	0.0313

Table 3: The values of limiting molar conductance (Λ_0), degree of dissociation (α), activity coefficient (Y_{\pm}), dissociation constant (K_D), triple ion association constant (K_3), association constant ($\log K_A$), Gibbs free energy change of association (ΔG_A), transfer free energy change (ΔG_t), Walden product ($\Lambda_0\eta$) and fluidity ratios (R_x) for nano-CuSO₄.5H₂O in (EtOH-H₂O) mixed solvents at room temperature.(293.15 K).

EtOH vol. %	X _s	Λo	α	γ±	$10^3 K_D$	$10^{3} K_{3}$	$\logK_{\rm A}$	ΔG_{A}	ΔG_t	Λ οη	Rx
0	0	248.7	0.9291	0.9597	3.7401	1.6921	2.4271	-13.6234	0.0000	2.4909	1.0000
20	0.0716	202.1	0.8035	0.9567	1.0027	3.3691	2.9988	-15.6839	-2.0604	2.0490	0.07641
40	0.1706	103.3	0.8180	0.9581	1.1258	3.1459	2.9485	-16.5501	-2.9266	1.3432	0.0501
60	0.3163	96.4	0.7116	0.9563	0.5354	4.5764	3.2712	-18.3615	-4.7381	1.0178	0.0379
80	0.5523	42.7	0.6777	0.9453	0.4246	4.9418	3.3720	-18.9270	-5.3036	0.4682	0.0174
100	1	4.2	0.7000	0.7726	0.5218	3.3472	3.2837	-18.4382	-4.8148	0.0728	0.0282

 Λ_{\circ} in (Ω^{-1} .cm².mol⁻¹), Λ_{\circ} in (Ω^{-1} .cm².mol⁻¹) and Δ_{\circ} G_A in (kJ mol⁻¹).



Figure 8: Variation of the association parameters (log K_A, $\Lambda_{0\eta}$, R_x) and Gibbs free energy change ($\Delta G_{A,..} \Delta G_{1}$) for bulk and nano-CuSO₄.5H₂O in (EtOH-H₂O) mixed solvents at (293.15 K).

It is obvious that the values of the association constant (KA) and the triple ion association constant (K₃) were found to be increased in increasing the EtOH content in the binary solvent. Gibbs free energies change of association (ΔG_A) and transfer Gibbs free energies change of association (ΔG_t) were increased with negative signs in increasing in EtOH content. On contrary the values of Walden product $(\Lambda_0\eta)$, fluidity ratio (R_x) , activity coefficient (γ_{\pm}), dissociation constant (K_D) of the CuSO₄.5H₂O solutions in the EtOH is less than in H2O. The decrease in the values of activity coefficient, limiting molar conductance, association constant, Gibbs free energy change of association, Walden product and fluidity ratios for nano-CuSO4 in comparison to bulk-CuSO4 indicate that the association of nano-CuSO4 is greater than bulk CuSO₄ salt, due to the high surface to volume ratio of nanoparticles which lead to greater ability for ion-pair formation.

Calculation of formation constants for interaction of Ciprofloxacin antibiotic and CuSO4.5H₂O solution in (EtOH-H₂O) mixed solvents

The binding of transition metal cation (Cu^{2+}) with the ligand ciprofloxacin antibiotic (CFL) can be represented by following equilibrium equation:

$$M^{2+} + L \leftrightarrow ML^{2+}$$
 (14)

where M^{2+} , L and ML^{2+} are the cation, ligand (CFL) and complex, respectively.

The formation constants (K_f) for CuSO₄complexes in terms of molar conductivity were estimated by using equation (15) (Takeda et al. 1983 a; Nasrabadi et al. 2009).

$$K_{f} = \frac{[ML^{2+}]}{[M^{2+}][L]} = \frac{\Lambda_{M} - \Lambda_{obs}}{(\Lambda_{obs} - \Lambda_{ML})[L]}$$
(15)

and

$$[L] = [L]_{t} - \left\{ [M]_{t} \frac{\Lambda_{M} - \Lambda_{obs}}{(\Lambda_{M} - \Lambda_{ML})} \right\}$$
(16)

Where, [L] is the ligand concentration, Λ_M is the limiting molar conductance of the CuSO₄ alone, Λ_{obs} is the apparent molar conductance of the solution during titration, Λ_{ML} is the molar conductance of the complex. The value of Λ_{ML} is determined from the value at point of large [L]_t to [M]_t ratios (Hakimi et al. 2008).

The variations of molar conductance (Λ_m) versus the $[L]_t / [M]_t$ mole ratio for $(Cu-CFL)^{2+}$ complex in binary EtOH-H₂O mixture were studied at room temperature Fig. (9). Different lines were obtained with breaks indicating the formation of 1:2 and 1:1 (M:L) stoichiometric complexes (Zollinger et al. 1987). The molar conductance value (Λ_m) always decreases with an increase in $[L]_t/[M]_t$ mole ratio, showing that the metal cation (Cu^{2+}) becomes less mobile upon complexation with ligand.

Valuable chemical information can be obtained between metal cations (Cu^{2+}) and ligand (CFL) in in binary EtOH-H₂O mixture. As it is shown in Table (4), the K_f value of the (Cu-CFL)²⁺ complexes increases with increase in EtOH ratios. Insolvents with medium donicites, such as EtOH, a more stable complex is formed, while in strong solvating solvents, such as H₂O, the complex formation is poor. In general, in higher donor solvent, the metal ion is more strongly solvated.

The Gibbs free energy change of formation (ΔG_f) for each stoichiometric complex was calculated from equation (17) (Takeda et al. 1983 b):

$$\Delta G_{\rm f} = -2.303 \text{ RT} \log K_{\rm f} \tag{17}$$

From the calculated (ΔG_f) values, the free energies change of transfer (ΔG_t) from water (w) as a reference solvent to the (EtOH-H₂O) mixed solvent (s) were calculated using equation (14).

$$\Delta G_t = \Delta G_{f(s)} - \Delta G_{f(w)}$$
(18)

Where $\Delta G_{A(s)}$ and $\Delta G_{A(w)}$ are the free energies of association in mixed solvents and in water, receptively.

The obtained values (K_f) for bulk and nano-CuSO₄ stoichiometric complexes and their calculated ΔG_f and ΔG_t values are presented in Tables (4 & 5).

It was found that formation Gibbs free energies change (ΔG_f) and transfer Gibbs free energies change (ΔG_t) were increased in negative signs with increasing in EtOH content in the binary solvent. Two stoichiometric complexes 1:2 and 1:1 (M:L) are formed and the formation constants and Gibbs free energies of the formed complexes follow the order: K_f (1:1) > K_f (1:2) for (M:L) and ΔG_f (1:1) > ΔG_f (1:2) for (M:L) (in negative values), which indicated the more favorable of formation of (1:1) complex compared with (1:2) complex. Also there are decrease in values of K_f and ΔG_f in case of using nano-CuSO₄ than that of using bulk-CuSO₄ but in small difference due to the more solvation effect in case of using nanoparticles.

Table 4: Formation constants (log K_f), Gibbs free energies change of formation (ΔG_f) and transfer free energies change (ΔG_t) for 1:2 and 1:1 (M/L) complex of bulk-CuSO₄ .5H₂O with Ciprofloxacin, in (EtOH-H₂O) mixed solvents at room temperature (293.15 K).

EtOH vol. %	X _s	Complex ratio (M:L)	log K _f	$\Delta G_{\rm f}$	ΔG_t
0	0	(1:2)	2.384	-13.381	0.000
		(1:1)	3.022	-16.965	0.000
20	0.0716	(1:2)	2.453	-13.771	-0.390

		(1:1)	3.123	-17.530	-0.565
40	0.1706	(1:2)	2.466	-13.844	-0.463
40	0.1706	(1:1)	3.206	-17.993	-1.028
60	0.3163	(1:2)	2.469	-13.860	-0.488
00		(1:1)	3.204	-17.985	-1.038
80	0.5523	(1:2)	2.559	-14.363	-0.982
		(1:1)	3.332	-18.702	-1.737
100	1	(1:2)	2.606	-14.628	-1.247
		(1:1)	3.450	-19.365	-2.400

Table 5: Formation constants (log K_f), Gibbs free energies change of formation (ΔG_f) and transfer free energies change (ΔG_t) for 1:2 and 1:1 (M/L) complex of nano-CuSO₄ .5H₂O with Ciprofloxacin, in (EtOH-H₂O) mixed solvents at room temperature (293.15 K).

EtOH vol. %	X_s	Complex ratio (M:L)	log K _f	$\Delta G_{\rm f}$	ΔG_t
0	0	(1:2)	2.382	-13.369	0.000
	0	(1:1)	3.020	-16.951	0.000
20	0.0716	(1:2)	2.450	-13.752	-0.383
20	0.0716	(1:1)	3.119	-17.507	-0.556
40	0.1706	(1:2)	2.492	-13.989	-0.620
		(1:1)	3.207	-18.002	-1.051
60	0.3163	(1:2)	2.492	-13.988	-0.620
		(1:1)	3.188	-17.896	-0.945
80	0 5523	(1:2)	2.584	-14.506	-1.137
	0.3325	(1:1)	3.325	-18.665	-1.714
100	1	(1:2)	2.603	-14.612	-1.243
	1	(1:1)	3.447	-19.347	-2.396

Figure 9: Variation of the molar conductance (Λ_m) with [M]/[L] for CuSO₄.5H₂O with CFL in (EtOH-H₂O) mixed solvents at (293.15 K).

Conclusion

The association constants of both bulk and nano-CuSO₄ salts in (EtOH-H₂O) mixed solvents were calculated using Fuoss-Shedlovsky method. It shows that the association parameters of nano-CuSO₄ is greater than bulk CuSO₄ salt due to the high surface to volume ratio of nanoparticles which leads to greater ability for ion – pair formation.

The thermodynamic parameters of complexation reactions between bulk, nano-CuSO₄ salts and ciprofloxacin antibiotic in (EtOH-H₂O) mixed solvents were calculated from conductance measurement. It was found that the formation Gibbs free energies change (ΔG_f) and transfer Gibbs free energies change (ΔG_f) were increased in negative signs with increasing in EtOH content in the binary solvent. Two stoichiometric complexes 1:2 and 1:1 (M:L) are formed and the formation constant and Gibbs free energy of the formed complexes follow the order: K_f (1:1) > K_f (1:2) for (M:L) and ΔG_f (1:1) > ΔG_f (1:2) for (M:L) (in negative values), which indicate the more favorable formation of (1:1) complex than (1:2) complex. Also there are decrease in values of K_f and ΔG_f in case of using nano-CuSO₄ than that of using bulk-CuSO₄ but in small difference due to the more solvation effect in case of using nanoparticles.

Acknowledgement

The authors gratefully acknowledge the approval and the support of this research study by the grant No. **7320-SAR-2017-1-8-F** from the Deanship of Scientific Research at Northern Border University, Arar, K.S.A.

References

- Angelusiu, M.V., Barbuceanu, S.F., Draghici, C., Almajan, G.L. (2010) Eur. J. Med. Chem. 2055–2062.
- Atkins, P.W.; (1978) Physical chemistry; Oxford University Press.
- Baddour LM, Wilson WR, Bayer AS, Fowler VG, Bolger AF, Levison ME, et al. (June 2005). "Infective endocarditis: diagnosis, antimicrobial therapy, and management of complications: a statement for healthcare professionals from the Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease, Council on Cardiovascular Disease in the Young, and the Councils on Clinical Cardiology, Stroke, and Cardiovascular Surgery and Anesthesia, American Heart Association: endorsed by the Infectious Diseases Society of America". Circulation. 111 (23): e394– 434.
- Beurskens, P. T and Jeffrey, G. A. (1964) J. Chem. Phys., 40, 906.
- Bishop, E.and Jennings, V. J. (1958) Talanta, 1, 197.
- Christopher, R.K. Glasson, A. Lindoy, L.F. George, V. Meehana, M. (2004) Developments in the d-block metallosupramolecular chemistry of polypyridyls, Coor. Chem. Rev. 252 940–948.
- Debye., P. Huckel, H.: (1923) The theory of electrolyte II The border law for electrical conductivity. Phys. Z. 24, 305.
- El-Hady, M.N.A. Zaky, R.R. Ibrahim K.M., Gomaa EA, (2012), (E)-3-(2-(furan-ylmethylene) hydrazinyl)-3-oxo-N-(thiazol-2yl) propanamide complexes: Synthesis, characterization and antimicrobial studies. Journal of Molecular Structure 1016 (Complete), 169-180.
- Franck, H. S.and Wen, Y. W. (1957) Discuss. Faraday Soc., 24, 133.
- Fuoss R. M. and Accascina F.; (1959 a) Electrolytic conductance, Interscience, New York.
- Fuoss, R. M. Proc. Nat. Acad. Sci. (1959 b). 45, 807.
- Gomaa EA, Abou Elleef EM, Abdel-Razek MG (2013 a) Thermodynamics of the Solvation of CaSO4 in Mixed DMF-H2O at 301.15 K. International Research Journal of Pure & Applied Chemistry. 3 (4): 320-329.
- Gomaa EA, Abou Elleef EM, Helmy ET (2014 b) Conductance Studies on Complex Formation between CaCl2 and Ampicillin As Ligand In Water And In Methanol Solvent at Different Temperatures. RESEARCH & REVIEWS:

JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES. 3 (3): 45.

- Gomaa EA, AbouElleef EM (2013 b) Thermodynamics of the solvation of lead nitrate in mixed DMF-H2O solvents at 301.15 K. American
- Gomaa, E. A., AbouelLeef, E. M., K. S Shalaby, S. E Salem. (2014 a). Thermodynamic Effect of Bulk and Nano-CuSO4 Salts on Tenoxicam Using a Variety of Different Techniques. Journal of Environments 1 (2), 44-53.
- Grabe M, Bjerklund-Johansen TE, Botto H, Çek M, Naber KG, Pickard RS, Tenke P, Wagenlehner F, Wullt B (2013). "Guidelines on Urological Infections"(PDF). European Association of Urology. Archived from the original(PDF) on 31 December 2013.
- Gryzybkowski W. and Pastewski R.; (1980) Electrochimica Acta; 25, 279.
- Hakimi, M. Nezhadali, A. Naemi, A. (2008) Conduct metric study of complex formation between Cu(II) ion and 4amino-3-ethyl- 1,2,4-triazol-5-thione in binary ethanol/water mixtures, E-J. Chem. 5 551–556.
- Harwood, L. M. and Moody C. J. (1989). Experimental organic chemistry: Principles and Practice (Illustrated ed.). Wiley-Blackwell. p. 292.
- Helmy, E.T., Gomaa, E. A., AbouelLeef, E. M., (2016) Gibbs Free Energy and Activation Free Energy of Complexation of Some Divalent Cations with Ampicillin in Methanol at Different Temperatures. American Journal of Applied Chemistry; 4(6): 256-259.
- Hirsch E. and Fuoss R.M.; (1956) J. Amer. Chem. Phys., 25; 1199.
- Isabel, M. S. Lampreia, L. and Ferreira, A.V. (1996) J. Chem. Soc. Faraday. Trans. T., 92; 1487.
- Ives D.J.G.; (1971) Chemical Thermodynamics, University Chemistry, Macdonald Technical and Scientific, London.
- Jose, N., Sengupta, S., and Basu, J.K. (2009) J Mol Catal A-Chem, 309 153-158.
- Kay, R. L. (1973) Water a Comprehensive Treatise, , 3, 173-209.

Kay, R. L. (1991) Pure Appl. Chem., 63, 1393.

- Kay, R. L., Cunninghan, G. P., Fennel, D. (1968) Hydrogen Bonded Solvent Systems, 249-260.
- Nasrabadi, M. R. Ahmadi, F. Pourmortazavi, S.M. Ganjali, M.R. Alizadeh, K.; (2009) Journal of Molecular Liquids; 144 97–101.
- Nezhadali, A. Rabani, N. (2011) Competitive bulk liquid membrane transport of Co(II), Ni(II), Zn(II), Cd(II), Ag(I), Cu(II) and Mn(II), cations using 2,20-dithio (bis) benzothiazole as carrier, Chin. Chem. Lett. 22 88–92.
- Nezhadali, H.A. Hosseini, P. Langara, (2009) Study of complex formation between iodoquinol (IQ) and Co2+, Mn2+, Cd2+, Pb2+ and Zn2+ cations in DMF/EtOH binary solvent mixtures using spectrophotometry, Polish J. Chem. 83573–580.
- Nezhadali, P. Langara, H.A. Hosseini, (2008) Study of complex formation between 5, 7-diiodo-8-hydroxyquinoline and Zn2+, Cd2+, Pb2+ and Tl+ cations in binary non-aqueous solvents using square wave polarography technique (SWP), J. Chin. Chem. Soc. 55 271–275.
- O. Popuych, A. Gibofsky and D.H. Berne. (1972) Analytic. Chem., 44; 811.
- Shdlovsky T. and Kay R.L.; (1956) J.Phys., Chem. 60; 151.
- Takeda Y.: (1983 a) Thermodynamic Study for Dibenzo-24crown-8 Complexes with Alkali Metal Ions in Nonaqueous Solvents. Bull. Chem. Soc. Jpn. 56, 3600–3602.
- Takeda Y.; (1983 b) Bull. Chem. Soc. Jpn 56, 3600.
- Varghese J. N. and Maslen, E. N. (1985) Acta Crystallogr.B41, 184.
- Walden, P. Z. (1906) Physik. Chem., 55, 207-246.
- Zollinger., D.P. Bulten, E. Christenhusz, A. Bos, M., Van der Linden, W.E: (1987) Computerized conduct metric determination of stability constants of complexes of crown ethers with alkali metal salts and with neutral molecules in polar solvents. Anal. Chim. Acta. 198, 207–222.
- Zwanzig, R., (1963). J. Phys. Chem., 38, 1903.