Anti-inflammatory activity and COX 2 inhibition potential of Trois in murine macrophage cell line

Manu Chaudhary, Santosh Kumar Patnaik, Anurag Payasi*

Received: 28 February 2015 / Received in revised form: 07 August 2015, Accepted: 28 October 2015, Published online: 25 December 2015 © Biochemical Technology Society 2014-2015

Abstract

The objective of present study was to investigate the antiinflammatory effects of Trois in lipopolysaccharide (LPS) induced macrophage RAW 264.7 cells. A biomembrane stabilisation assay was also conducted to evaluate its role in biomembrane fluidity during inflammation. Cytotoxicity of drugs on RAW 26-4.7 macrophages cell line was conducted using 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. mRNA expression of cyclooxygenase-2 (COX-2) was studied using polymerase chain reaction (PCR). Results of our study revealed that IC₅₀ of Trois and volini was 2.82 and 1.73 mg/ml, respectively, after 24 hrs treatment. Further, our data proved that Trois had excellent anti-inflammatory activity as it caused reduced production of nitric oxide (9-12%), myeloperoxidase (5-14%) and mRNA of COX-2 (12-33%) in comparision to volini. More-over, Trois had 5.44% more membrane stabilization potential compared to volini. From the above study it can be concluded that Trois can be an excellent option for the treatment of inflammation related diseases.

Keywords: Trois, membrane stabilization, inflammation, cyclo-oxygenase-2 inhibition.

Introduction

Inflammation is a common clinical condition, which is typically characterized by redness, swelling, pain and heat. Majority of the human populations are affected by inflammation related disorders at some point, across the world (Bhagyasri et al., 2015). A growing line of evidences have indicated that inflammation is involved in the pathogenesis of many diseases including arthritis and other lifethreatening and debilitating disorders (Finch, 2005; Caruso et al., 2004; Lee et al., 2014; Kumar et al., 2009; Manouri et al., 2015). Among the various causes, oxidative stress is the primary cause of inflammation. It can induce inflammatory cells to produce inflammatory mediators, such as cytokines and chemokines, which

Manu Chaudhary, Santosh Kumar Patnaik, Anurag Payasi*

Department of cell culture and molecular biology, Venus Medicine Research Centre, Baddi, H.P.,173205 India

*Email: ccmb@vmrcindia.com

enhance tissue damage from the recruitment of more inflammatory cells, eventually resulting in more oxidative stress (Mansouri et al., 2015; Kasama et al., 2005; Wolf et al., 2005; Odobasic et al., 2014). Induction of pro-inflammatory cytokines (IL-6, TNF-a, and NF-kB) can induce nitric oxide (NO) production and cyclo-oxygenase-1 (COX-1) which contribute in inflammation (Kirby et al., 2014; Saraswathi et al., 2015; Alhouayek and Muccioli, 2014). Another inflammatory process is the production of matrix metalloproteinases (MMPs). MMPs are extracellular matrix degrading enzymes that can be secreted by the induction of many pro-inflammatory cytokines and interleukins (Breckwoldt et al., 2008). Several reports have demonstrated the involvement of COX-2, NO and myeloperoxidase (MPO) in inflammation of rheumatoid arthritis or spondyloarthropathies (Sharma et al., 2008; Breckwoldt et al., 2008; Saraswathi et al., 2015; Alhouayek and Muccioli, 2014). Lipid peroxidation causes an auto-oxidation of polyunsaturated fatty acids in the membrane leading to reduced membrane fluidity (Chippada et al., 2011).

Inflammation related diseases are commonly treated by nonsteroi-dal anti-inflammatory drugs (NSAIDs). However NSAIDs are not always useful due to their side effects (Bhagyasri et al., 2015; Ki-rby et al., 2014). To avoid side effects of NSAIDs, natural reme-dies can be the alternative option. Natural remedies are known to have less side effects and toxicity (Arif et al., 2009; Chatterjee et al., 2015; Kapewangolo et al., 2015). Medicinal plants have been wide source for the treatment of various diseases and nearly 80% of the world populations rely on medicinal herbs for their primary health care (Jordan et al., 2010). There is increasing evidence that medicinal plants have many biological properties including anti-oxidant, anti-inflammatory, anti-cancer and antiatherosclerotic effects (Zhen et al., 2015; Bhagyasri et al., 2015; Chatterjee et al., 2015).

Trois is a nanotechnology based herbal product that contains wintergreen oil (*Gaultheria precumbens*) and eucalyptus oil (*Eucalyptus globulus*) as major ingredients. It is indicated for the treatment of pain and inflammation associated with different types of arthritis including osteoarthritis, gouty arthritis, rheum-atoid arthritis, ankylosing spondylitis, sprains and backache. To study its anti-inflammatory effects, murine macrophages cells i.e. RAW 264.7 cells have been used as in vitro inflammatory model in which inflammation was induced by lipopolysaccharides (LPS) (Mosmann, 1983; Seo et al., 2011; Achoui et al., 2010).

This study was aimed to assess the anti-inflammatory effect Trois in LPS induced macrophage RAW 264.7 cells. Biomembrane stabilisation assay was also conducted to evaluate its biomembra-ne fluidity ability during inflammation.

Materials and Methods

Drugs

Following two drugs were used in the study: Trois (Venus Remedies Limited, Baddi, India) and volini (Ranbaxy Laboratories Limited, Paonta Sahib, India). Trois contains wintergreen oil 18.9 % (v/v), eucalyptus oil 6.06 % (v/v), menthol 5.08 % (w/v), *Vit-ex negundo* leaves extract 3.24 % (w/v) and *Apium graveolens* fruit extract 3.24 % (w/v) and volini contains diclofenac diethyl amine 1.16 % (w/w) equivalent to diclofenac sodium (1 %), linseed oil BP (3 % w/w), methyl salicylate IP (10 % w/w) and menthol IP 5 % (w/w).

Cell viability

Viability of the cells was assessed using the 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) colorimetric assay as described earlier (Mosmann, 1983). The percentage of cell viability was calculated by the following for-mula:

Cell viability (%) = OD_{570sample} X 100 / OD_{570control}

IC50 determination of drugs

To evaluate the effect of drugs on LPS stimulated inflammation on macrophage RAW 264.7 cell, 5×10^3 cells/well seeded into 96 well plate and grouped as follows: A, B, C, D, E, F, G and H. Cells in group A were treated with PBS which served as negative control. Cells in group B to H were stimulated with LPS (1 µg/ml) and cells in group B served as positive control. Cells in C to H groups were simultaneously treated with various concentrations of Trois (0.25 to 8 µg/ml). Only cell culture media in another group I served as blank. The same drug treatment procedure was adopted for volini with drug concentrations ranged 0.25 to 8 µg/ml in another 96 wells plate. After drug treatment, plates were incubated for 24 hrs and processed for IC₅₀ determination. The cytotoxic



Figure 1: Flow chart of entire experiment.

Cell culture

RAW 264.7 mouse macrophages cell line was obtained from National Centre for Cell Sciences (NCCS), Pune, India. It was grown in Dulbecco's Modified Eagle's Medium (DMEM, HiMedia, Mumbai, India) supplemented with 10% fetal bovine serum (FBS, Bio Gene, India) and 1% antibiotic / antimyotic solution at 37°C in an atmosphere of 5% CO_2 in a humidified incubator. Cells of 70-90 % confluent flasks with >90% viability were used for further study. The flow chart of entire experiment is shown in Figure 1.

doses that kill cells by 50% (IC₅₀) was determined from cell viability (%) versus concentration curve. All the experiments were done in triplicate.

RNA isolation

Total mRNA from LPS stimulated RAW 264.7 cells (positive control; LPS+), without LPS stimulated cells (negative control; LPS-) and LPS stimulated and drugs treated (Trois and volini) cells with half of IC_{50} values was isolated using Trizole (Invitrogen, Banglore, India). The ratio of absorbance at 260 nm and 280 nm was used to assess the purity of RNA. A ratio

J Biochem Tech (2015) 6(2): 938-944

of ~2.0 is generally accepted as "pure" for RNA. RNA was stored at -70 $^{\circ}\mathrm{C}$ until use.

Complementary deoxyribonucleic acid (cDNA) synthesis

Total RNA (2 μ g) of each treatment group was converted to first strand cDNA as follows: Two microgram of RNA was added to 1 μ l of oligo dT primer and 9.2 μ l of water and the mixture was incubated at 65 °C for 5 min. After incubation, following reagents were added sequentially: 4.0 μ l of 5X RT buffer, 1.0 μ l of 0.1M DTT, 0.5 μ l of 10 mM dNTPs and 0.3 μ l of 20 U moloney murine leukaemia virus reverse transcriptase (MMLVRT), mixed well and the mixture was subsequently incubated at 37°C for 60 min. The reaction was stopped by heating at 70°C for 10 min. The resultant solution was cDNA which can be used for gene expression study by PCR.

PCR analysis

PCR was performed using the cDNA as template. The primers used in this study were obtained from sigma Aldrich (Chemicals Private Limited, Bangalore, India). The following sequences of primers were used. For COX-2-F-5-ATGCTCCTGCTTGAGTATGT-3; COX-2-R-ATGCTCCTG CTTGAGTATGT-3'. β-actin-F-5 GTGGGCCGCCCTAGGCA CC AG-3', for β-actin-R-5-GGA-GGAAGAGGATGCGGC AGT-3'. For PCR amplifications, about 3 µl of cDNA was added to 20 µl mixture containing 0.5 mM of dNTPs, 1.25 µM of each primer and 3.0 U of Taq polymerase (Banglore Genei) in 1x PCR buffer. Amplification was performed in an Eppendorf thermocycler (Germany) with following cycling parameters: initial denaturation at 94°C for 5 min followed by 35 cycles (45 seconds of denaturation at 94°C, 45 second of anneali-ng at 60-62 °C and 1 min of extension at 72°C) and final extension at 72°C for 5 min. PCR products were then electrophoresed on a 1.5 % agarose gel containing ethidium bromide. The expected amplicon size for COX-2 gene was 696 base pairs and β-actin gene was 603 base pairs. PCR products were then electrophoresed on a 1.0 % agarose gel containing ethidium bromide and intensity of each amplified band was analyzed using the image J software.

Measurement of myeloperooxidase (MPO)

MPO activity was measured using 3,3,5-5 tetramethylbenzidine (TMB) as described elsewhere (Suzuki et al., 1983). About 5×10^3 cells/well were seeded into 96 well plate and grouped as follows: negative control (without LPS stimulated cells; LPS-), positive control (LPS stimulated cells; LPS+), drug treatment groups (treated with drugs at half of IC₅₀ values following LPS stimulation) and blank (only media). The plates were then incubated for 24, 48 and 72 hrs in CO₂ incubator (37 °C, 5 % CO₂). After incubation at indicated time intervals, 10 µl of cell culture media from each well was withdrawn and used for MPO activity measurement. Cell culture media was used as a blank. The released MPO from cells was determined by comparing the values to a standard curve generated using known concentrations of 10 mM hypochlorous acid. The percentage of MPO inhibition were calculated as follows:

% Inhibition = MPO_{control} - MPO_{sample} / MPO_{control} X100

Measurement of nitric oxide (NO)

NO production was assayed by measuring the level of nitrite in the supernatant of cultured RAW264.7 cells as described earlier (Kang et al., 2008). Grouping and treatment of cells were made as described in the above MPO section. The plates were then incub-

ated for 24, 48 and 72 hrs at 37 °C, 5% CO₂. Following incubation at indicated time intervals, 100 μ l of cell culture medium was withdrawn and used for NO assay. Fresh culture medium was used as a blank in every experiment. The concentrations of nitrite were then determined by comparing the values of test samples to a standard curve generated using known concentrations of sodium nitrite (μ M). All experiments were performed in triplicate. The percentages of NO inhibition were calculated as follows:

% Inhibition = 100 X NO_{2control}-NO_{2sample} / NO_{2control}

Human red blood cells (HRBC) membrane stabilization assay

Erythrocytes are considered to be an excellent model for membrane stabilization assay. This assay was performed according to the method described earlier (Ananthi and Chitra, 2013). Briefly, 5 ml of blood from healthy human was collected in EDTA vacutainer tube. Then, the blood was centrifuged at 3000 rpm for 10 min and packed RBCs were washed two times with isosaline (0.9 % NaCl) and 10 % suspension of RBCs was made. For membrane stabilization assay, reaction system consisted of 88.9 µl of hyposaline (0.25 w/v NaCl), 44.4 µl of 0.15 M phosphate buffer (pH 7.4) 22.2 µl of 10 % RBC suspension and test solution (half of IC50) in isosaline. For positive control, 44.4 µl of distilled water was added into reaction system instead of hyposaline to produce 100 hemolysis while negative control was lacking RBC suspension. All the tubes were incubated at 37°C for 15, 30, 60 and 120 minutes. At every interval of time, the tubes were centrifuged at 3000 rpm for 20 minutes. The haemoglobin content in the suspension was then estimated using spectrophotometer at 560 nm. The percentage of HRBC membrane stabilization was calculated using following formula:

% Protection= 100 - (Optical density of sample/Optical density of control X 100)

Statistical analysis

Data was analyzed using Graph pad prism 5.01 and expressed as mean \pm standard deviation (SD). The continuous variables were tested with one-way analysis of variance (ANOVA) and turkey test. Values of P>0.05 were considered not to be significant.

Results

Cytotoxicity assay

To examine the cytotoxic effects of drugs, MTT assay was performed on RAW 264.7 cells. Results of our study revealed that IC_{50} of Trois and volini was 2.82 mg/ml and 1.73 mg/ml, respectively after 24 hrs treatment. For further study half of IC_{50} values were used.

NO assay

Stimulation of RAW264.7 cells with LPS caused 12 to 14 folds increase in nitrite concentration in comparison to LPS unstimulated cells (negative control; LPS-) at 24, 48 and 72 hrs. Following treatment of cells with half of IC50 of Trois led to 80 to 95 % inhibition of NO production in comparison to LPS stimulated cells (positive control; LPS+) at 24, 48 and 72

hrs. Similarly, volini produced 69 to 86 % inhibition of NO in comparison to positive control (LPS+ve) at the same time intervals. When we compared the inhibition percentage of Trois and volini, Trois inhibited 9 to 12 % more NO production (Figure 2).



Figure 2: Effect of Trois and volini on NO production in LPS induced RAW264.7 macrophages cell.

The production of NO was assayed in the culture medium of cells stimulated with LPS for 24, 48 and 72 hrs. Values are mean \pm SD of triplicate experiments.

MPO assay

Similar to NO, cells stimulated with LPS produced about 9 fold increase in MPO concentration compared to LPS unstimulated cells (negative control; LPS-) at 24, 48 and 72 hrs. Our results showed that Trois markedly inhibited LPS induced MPO production by 62 to 92 % in comparison to LPS stimulated cells (positive control; LPS+) whereas volini produced only 57 to 81 % inhibition of MPO. Trois inhibited 5 to 14 % more MPO production compared to volini (Figure 3).



Figure 3: Effect of Trois and volini on MPO production in LPS induced RAW264.7 macrophages cell.

The production of MPO was assayed in the culture medium of cells stimulated with LPS for 24, 48 and 72 hrs. Values are mean \pm SD of triplicate experiments.

HRBC assay

Treatment of cells with Trois produced stabilization percentage of 56.8, 85.8, 89.9 and 92.0 % where as volini showed 41.4, 71.0, 76.5 and 83.7% stabilization after 15, 30, 60 and 120 min respectively. From these results, it is clearly evident that Trois had 8 to 15% more membrane stabilization potential compared to volini. Trois showed 15.4% more membrane stabilization potential compared to volini after 15 min indicating onset of action and free radical neutralization is faster in case of Trois (Figure 4).





COX-2 gene expression study

The expression of COX-2 mRNA was elevated in cells treated with LPS (positive control; LPS+) compared to LPS unstimulated cells (negative control; LPS-). Following treatment of cells with



Figure 5: Relative COX-2 gene expression in various groups. Values are mean \pm SD of triplicate experiments. *, p<0.05; **, p<0.01; ***, p<0.001.

drugs, Trois produced 28.7, 38.7, 76.2 % reduction in COX-2 m-RNA expression after 24, 48 and 72 hrs of treatment, respectively which correspond to **, p<0.01;***, p<0.001 and ***, p<0.001 in comparision to positive control. On the other hand, volini produced 17.5, 27.5, and 42.8 % reduction in COX-2 mRNA expression after 24, 48 and 72 hrs of treatment, respectively which correspond to *, p<0.05;**, p<0.01 and ***, p<0.01 in comparison to positive control (Figure 5 and Figure 6, A,B &C). Further, a 33.4% more COX-2 inhibition by Trois in comparison to volini after 72 hrs indicating the long acting action of Trois.

Discussion

Increasing number of natural plants have been used across the globe to treat various kinds of acute and chronic diseases (Pan et al., 2006; Bhagyasri et al., 2015; Margaret et al., 2014; Kapewangolo et al., 2015). The natural products which are derived from plants reduce the risk of developing pathological conditions, including cancer, nervous system disorders, cardiovascular, genetic, and inflammatory diseases (Jurenka, 2009; Newman and Cragg, 2007; Chatterjee et al., 2015, Kelly et al., 2014). Various drugs which contain wintergreen oil (Gaultheria precumbens), camphor oil (Cinnamomum camphora), eucalyptus oil (Eucalyptus globulus) and menthol as major ingredients have shown to have anti-inflammatory and analgesic properties (Chippada et al., 2011; Hajhashemi et al., 2003; Salleh et al., 2012; Alsaad et al., 2015; Mota et al., 2015).



Figure 6: Effect of Trois and volini on the mRNA expression of COX-2 in RAW264.7 cells. A-(A) without stimulation (B) stimulated with LPS (C) stimulated with LPS in presence of volini for 24 hrs (D) stimulated with LPS in presence of Trois for 24 hrs. B- (A) without stimulation (B) stimulated with LPS (C) stimulated with LPS in presence of volini for 48 hrs (D) stimulated with LPS in presence of Trois for 48 hrs. C- (A) without stimulated with LPS in presence of Volini for 72 hrs (D) stimulated with LPS (C) stimulated with LPS in presence of Trois for 72 hrs.

In this study, we established invitro anti-inflammatory activity of Trois, a nanotechnology based herbal product in comparison to volini (a diclofenac and menthol based product) in murine macrophage RAW264.7 cells. It has been reported that LPS induces iNOS transcription and transduction with subsequent NO production (MacmMicking et al., 1997). NO production has been implicated in a variety of pathological processes including various forms of inflammation and inflammatory disorders (Pan et al., 2006; Kitisin et al., 2015). Our results revealed that Trois inhibited 9 to 12 % more LPS induced NO production in RAW cells compared to volini indicating higher anti-inflammatory mechanism of Trois may be through the greater NO production inhibition. Furthermore, Vigo et al. (2004), suggested the role of Eucalyptus globulus in NO inhibition. Moreover, it has been indicated that Apium graveolens which containe apiin inhibit the nitrite (NO) production and also iNOS expression (Mencherini et al., 2007). Naithani et al. (2011) also demonstrated that Trois strongly inhibited the NO production.

MPO plays a central role in oxidant production by neutrophils. Also, it is the most abundant pro-inflammatory enzyme (Pulli et al., 2013; Marlous et al., 2014) which is secreted during inflammatory response and contributing to arthritis development (Breckwoldt et al., 2008; Garcia et al., 2014). Our results clearly showed that Trois inhibited 5 to 14 % more MPO production in RAW cells compared to volini. Previous studies have also demonstrated that many plants may inhibit MPO activity (Nastasijevic et al., 2012; Vieira et al., 2011; Leskovac et al., 2013), therefore, anti-inflammatory effect of Trois may be due to inhibition of MPO. Data indicated that NO has profound effect on COX-2 catalytic activity (Wu 1995). However, the exact mechanism needs to be explored. One possibility is that lipid peroxidation initiated by peroxynitrite the product of the reaction of NO with superoxide, liberates arachidonic acid from the cell membrane which in turn activates COX-2 (Davidge et al., 1995; Esposito and Cuzzocrea 2007). Our data further indicated that Trois exhibited significant down regulation in COX-2 mRNA expression indicating that the action of Trois occurs at transcriptional level.

Several studies have demonstrated that drugs targeting COX2 enzyme might provide a better anti-inflammatory activity (Ronchetti et al., 2009; Turini and DuBois 2002).

Erythrocyte membrane is an analogue to the lysosomal membrane (Breckwoldt et al., 2008). Stabilization of membrane is important in limiting the inflammatory response by preventing the release of lysosomal constituents of activated neutrophil such as proteases which cause further membrane damage up on extracellular release (Murugaesan et al., 1981). Our data demonstrated that Trois has enhanced membrane stabilization potential which is evident by higher protection of erythrocyte membrane from damage.

Conclusions

The anti-inflammatory property of Trois is strongly supported by its ability to reduce the NO, MPO and mRNA of COX-2 production as these are regarded as potent inflammatory mediators at concentration that were not toxic to cells. Therefore, Trois can be termed as more safe topical preparation for management of inflammatory disorders.

Acknowledgement

Researchers are thankful to Venus Medicine Research Centre, Werne, Germany for providing financial assistance (Grant no. VPG-005-2013) to carry out this study.

References

- Achoui M, Appleton D, Abdulla MA et al. (2010) In vitro and in vivo anti-inflammatory activity of 17-Oacetylacuminolide through the inhibition of cytokines, NFκB translocation and IKKβ activity. PLoS One 5: e15105.
- Ananthi T, Chitra M (2013) Screening of in-vitro antiinflammat-ory activity of *Michelia champaca* L. lowers. Asian J Pharm-aceut Clin Res 6:71-72.
- Alhouayek M, Muccioli GG (2014) COX-2-derived endocannabi-noid metabolites as novel inflammatory mediators. Trends Pharmacol Sci 35:284-92.
- Alsaad AMS, Fox C, Koren G (2015) Toxicology and teratology of the active ingredients of professional therapy muscle care products during pregnancy and lactation: a systematic review. BMC Compl Altern Med 15:40.
- Arif T, Bhosale JD, Kumar N et al. (2009) Natural products anti-fungal agents derived from plants. J Asian Nat Prod Res 7:621-638.
- Breckwoldt MO, Chen JW, Stangenberg L et al. (2008) Tracking the inflammatory response in stroke in vivo by sensing the enzyme myeloperoxidase. Proc Nat Acad Sci 105:18584-18589.
- Bhagyasri Y, Lavakumar V, DivyaSree MS et al. (2015) An overview on anti-inflammatory activity of Indian herbal plan-ts. Int J Res Pharmaceut Nano Sci 4:1-9.
- Chippada SC, Volluri SS, Bammidi SR et al. (2011) In vitro anti inflammatory activity of methanolic extract of centella asia-tica by hrbc membrane stabilisation. Rasayan J Chem 2:457-460.
- Chatterjee A, Sen B, Das S et al. (2015) Anti-inflammatory and analgesic activity of methanolic xtract of medicinal plant *Rhodiola rosea* rhizomes. Int J Pharma Res Rev 4:1-8.
- Caruso C, Lio D, Cavallone L et al. (2004) Aging, longevity, infl ammation and cancer. Ann N Y Acad Sci 1028:1-13.

- Davidge ST, Baker PN, Laughlin MK et al. (1995) Nitric oxide produced by endothelial cells increases production of eicosanoids through activation of prostaglandin synthase. Circ Res 77:274-83.
- Esposito E, Cuzzocrea S (2007) The role of nitric oxide synthase in lung inflammation. Current Opin Invest Drugs 8:899-909.
- Finch CE (2005) Developmental origins of aging in brain and blood vessels: an overview. Neurobiol Aging 26:281-91.
- García JJ, López-Pingarrón L, Almeida-Souza P et al. (2014) Protective effects of melatonin in reducing oxidative stress and in preserving the fluidity of biological membranes: a rev-iew. J Pineal Res 56:225–237.
- Hajhashemi V, Ghannadi A, Sharif B (2003) Anti-inflammatory and analgesic properties of the leaf extracts and essential oil of *Lavandula angustifolia*. J Ethnopharmacol 89:67-71.
- Jordan SA, Cunningham DG, Marles RJ (2010) Assessment of herbal medicinal products: challenges, and opportunities to increase the knowledge base for safety assessment. Toxicol Appl Pharmacol 243:198-216.
- Jurenka JS (2009) Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: a review of preclinical and clinical research. Alternat Med Rev14:141-153.
- Kang JY, Khan MN, Park NH et al. (2008) Antipyretic, analgesic, and anti-inflammatory activities of the seaweed Sargassum fulvellum and Sargassum thunbergii in mice. J Ethanopharmacol 116:187-190.
- Kasama T, Miwa Y, Isozaki T et al. (2005) Neutrophil-derived cytokines: potential therapeutic targets in inflammation. Curr Drug Targets Inflam Allergy 4:273-279.
- Kumar KJS, Wang SY(2009) Lucidon inhibits iNOS and COX2 expression in LPS mediated RAW264.7 murine macrophage cells via NF-KB and MAPKs signalling pathways. Planta Medica 75:494-500.
- Kirby SK, Lundberg MH, Wright WR et al. (2014) COX-2 protects against atherosclerosis independently of local vascu-lar prostacyclin: identification of COX-2 associated pathw-ays implicate Rg11 and lymphocyte networks. 9:e98165.
- Kapewangolo P, Omolo J, Bruwer R et al. (2015). Antioxidant and anti-inflammatory activity of *Ocimum labiatum* extract and isolated labdane diterpenoid. J Inflam12:4.
- Kelly R, Souzaa D, Silvab MA et al. (2014) Ethnopharmacology of medicinal plants of carrasco, northeastern Brazil. J Ethnopharmacol 157:99-104.
- Kitisin T, Saewan N, Luplertlop N (2015) Potential antiinflammatory and anti-oxidative properties of thai colored rice extra cts. Plant J 8:69-77.
- Leskovac A, Joksić G, Pašti I et al. (2013) The antiradical, antiinflammatory and anti-genotoxic potential of herbal preparation chlamyfin. J Chem Chem Eng 32:227–237.
- Lee EB, Fleischmann R, Hall S et al. (2014) Tofacitinib versus methotrexate in rheumatoid arthritis. N Engl J Med 370:-2377-2386.
- Margaret O, Sofidiyaa N, Foluso O et al. (2014) Antiulcer activity of the ethanolic extract and ethyl acetate fraction of the leaves of *Markhamia tomentosa* in rats. J Ethnophar-macol 157: 01-06.
- Mota VS, Turrini RNT, Poveda VB (2015) Antimicrobial activity of *Eucalyptus globulus* oil, xylitol and papain: a pilot study. Rev Esc Enferm USP 49:215-219.
- Mencherini T, Cau A, Bianco G et al. (2007) An extract of Apium graveolens var. dulce leaves: structure of the major constituent, apiin, and its anti-inflammatory properties. J Pharm pharmacol 59:891-7.
- Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity ass-ays. J Immunol Methods 16:55-63.

- Marlous RD, Poelgeest EP, Malone KE et al. (2014) Characteriz-ation of inflammation and immune cell modulati-on induced by low dose LPS administration to healthy volun-teers. J Inf-lam11:28.
- Mansouri MT, Hemmati AS, Naghizadeh B et al. (2015) A study of the mechanisms underlying the anti-infl ammatory effect of ellagic acid in carrageenan-induced paw edema in rats. Indian J Pharmacol 47:292-298.
- Murugaesan N, Vember S, Damodharan C (1981) Studies on ery-throcyte membrane IV: in vitro hemolytic activity of Olean-der extract. Toxicol Lett 8:33-38.
- MacmMicking J, Xie QW, Nathan C (1997) Nitric oxide and macrophage function. Annu Rev Immunol 15:323-350.
- Nastasijevic B, Lazarevic-Pasti T, Dimitrijevic-Brankovic S, et al. (2012) Inhibition of myeloperoxidase and antioxidative activity of *Gentiana lutea* extracts. J Pharm Biomed Anal 66:191-196.
- Naithani V, Singhal AK, Chaudhary M (2011) Comparative evaluation of metal chelating, antioxidant and free radical scavenging activity of Trois and six products commonly used to control pain and inflammation associated with arthritis. Int Drug Dev Res 3:208-216.
- Newman DJ, Cragg GM (2007) Natural products as sources of new drugs over the last 25 years. J Nat Prod 70:461-77.
- Odobasic D, Yang Y, Muljadi RCM et al. (2014) Endogenous myeloperoxidase is a mediator of joint inflammation and da-mage in experimental arthritis. Arthr Rheumatol 66:907–917.
- Pan MH, Lai CS, Wang YJ et al. (2006) Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice. Biochem Pharmacol 71:1293-1303.
- Pulli B, Ali M, Forghani R et al. (2013) Measuring myeloperoxi-dase activity in biological samples. Plos One 8:e67976.
- Ronchetti D, Borghi V, Gaitan G et al. (2009) A novel NOreleasing derivative of ferulic acid, suppresses inflammatory and nociceptive responses in in vitro and in vivo models. Br J Pharmacol 158:569-79.
- Seo JM, Lee J-Y, Ji GE et al. (2011) Down-regulation of ATPbinding cassette transporter G1 expression by unmethylated CpG oligodeoxynucleotides in RAW 264.7 macrophages. Exp Mol Med 43:510-516.
- Suzuki K, Ota H, Sasagawa S et al. (1983) Assay method for myeloperoxidase in human polymorphonuclear leukocytes. Anal Biochem 132:345-352.
- Sharma JN, Al-Omran A, Parvathy SS (2008) Role of nitric oxide in inflammatory diseases. Inflammopharmacol 15:252-259.
- Salleh FM, Anuar TS, Yasin AM et al. (2012) Wintergreen oil: a novel method in wheatley's trichrome staining technique. J Microbiol Methods 91:174-178.
- Saraswathi V, Meenal K, Perriotte-Olson C (2015) Ablation of COX-2 attenuates adipose inflammation but impairs metabo-lic homeostasis in diet-induced obesity. The FASEB J 29:1 Supplement 818.8.
- Turini ME, DuBois R (2002) Cyclooxygenase-2: a therapeutic target. Annul Rev Med 53:35-57.
- Vigo E, Cepeda A, Gualillo O et al. (2004) In-vitro antiinflammatory effect of *Eucalyptus globulus* and *Thymus vulgar-is*: nitric oxide inhibition in J774A.1 murine macroph-ages. J Pharma Pharmacol 56:257-63.
- Vieira TO, Seifriz I, Charão CCT et al (2011) Antioxidant effects of crude extracts from baccharis species: inhibition of myel-operoxidase activity, protection against lipid

peroxidation, and action as oxidative species scavenger, Rev Bras Farmacogn 21:601-607.

- Wu KK (1995) Inducible cyclooxygenase and nitric oxide syntha-se. Adv Pharmacol 33:179-207.
- Wolf AM, Wolf D, Rumpold H et al. (2005) The kinase inhibitor imatinib mesylate inhibits TNF-{alpha} production in vitro and prevents TNF-dependent acute hepatic inflammation. Pr-oc Nat Acad Sci 102:13622-13627.
- Zhen J, Guo Y, Villani T et al. (2015) Phytochemical analysis and anti-inflammatory activity of the extracts of the African medicinal plant *Ximenia caffra*. J Analyt Meth Chem 20-15:ID 948262.